

INGUINAL HYDATID CYST: REPORT OF A RARE CASE

M.R. KOUHSARI, M.D., AND H.A. MANZAR, M.D.

From the Department of Surgery, Razi Hospital, Guilan University of Medical Sciences, Rasht, Islamic Republic of Iran.

ABSTRACT

Hydatid cyst, the larval stage of *Echinococcus spp.*, is a common parasitic disease in Iran. The liver and the lungs are the most commonly involved organs. Hydatid cysts have also been reported in bones and soft tissues such as muscle, brain tissue, spleen, thyroid and even in the heart. Hydatid cyst of the inguinal area, resembling a hernia is a rarity presenting at an unusual site. A hypersensitivity reaction to components of hydatid fluid may cause a serious or even fatal reaction if cyst rupture occurs. Sudden occurrence, lack of a good clinical history, and the severity and complexity of symptoms and signs makes the diagnosis difficult. This paper reports sudden rupture of an inguinal hydatid cyst in a patient while taking a bath, resulting in severe anaphylactic shock and acute pulmonary edema, wrongly described by the patient's family as "steam suffocation". This case report emphasizes the value of a good and detailed clinical history, the need to include hydatid cyst in the list of inguinal swellings in endemic areas, and to consider hydatid cyst rupture in the differential diagnosis of sudden shock, sudden LVF, pulmonary edema and severe allergic reactions.

MJIRI, Vol. 12, No. 2, 177-179, 1998.

INTRODUCTION

Human infection with echinococci has a world-wide distribution.¹ Although relatively uncommon in most of North America, echinococcal disease of the liver is found throughout the world, particularly in areas where sheep are raised. These areas include the Mediterranean, Australia, New Zealand and countries in the Middle East such as Iran.²

Almost every organ of the human body has been involved, including the brain, heart, spleen, biliary tract, thyroid, parotid, tongue, tooth, bone, broad ligament,

prostate, bladder, tonsil, pulmonary artery and inferior vena cava.³ Our case was present in the inguinal canal.

Daughter cysts may leak from the main cyst when punctured by a biopsy needle and these may seed the peritoneal cavity. This can also occur as a result of blunt abdominal trauma.⁴ Intraperitoneal rupture results in the showering of hydatid fluid, brood capsules and scolices into the peritoneum leading to transient peritoneal irritation.⁵ A hypersensitivity reaction to components of hydatid fluid may cause a serious or even fatal reaction, if there is rupture of cysts.⁶ Usually the reproductive elements survive and initiate the formation of new cysts.⁷

We report a case of secondary hydatid cysts resembling bilateral inguinal hernia. The left one ruptured due to blunt trauma resulting in acute hypersensitivity reaction and the right sided inguinal hydatid cyst was removed

Correspondence:

M.R. Kouhsari, M.D., Assistant Professor of Surgery, Razi Hospital, Rasht, I.R. Iran, 41448.




Fig. 1. Intraoperative view of the hydatid cyst.

surgically at a later date.

Case report

In March 1989, a 31 year old young man from Manjeel was admitted to the Medical Emergency Ward of Roodbar General Hospital of Guilan. His clinical presentation included dyspnea, sweating, left inguinal pain and shock. In the emergency room he was unable to describe his medical history. His wife reported that he had no cardiovascular or respiratory illness in the past. Four years back, he had undergone abdominal surgery for liver disease. Recently he complained of hernia-like swellings in both groin areas, one of which reduced while he was taking a bath. At the same time, symptoms of "steam suffocation" also appeared.

On physical examination, he was pale, anxious and breathless. Both ears were red and edematous. Heart rate was 108/min and regular. Blood pressure was 70 mmHg (systolic).

On cardiac auscultation, a normal S_1 & S_2 with an audible S_3 was present. Bronchial breath sounds with wheezing and rhonchi were scattered throughout both lungs and fine crepitations were audible at the base of both lungs. Signs of bronchospasm were more clear on auscultation of the lung apex.

A surgical scar was visible at the right upper quadrant. The liver and spleen and no mass were palpable in the abdomen. The left inguinal region was tender. A cystic, non-tender oval swelling of 10 cm was palpable in the right groin.

Aside from sinus tachycardia, no significant change could be found in the electrocardiogram. Hypersensitivity and allergic shock was successfully managed in the emergency room by appropriate parenteral drug treatment.

Routine blood tests showed 8% eosinophilia. Other hematology and blood chemistry tests were normal. Nothing abnormal was detected in the chest radiograph

and abdominal ultrasound study. Right inguinal sonography reported a cystic mass of 10 × 10 cm.

Surgical approach and outcome

Upon incision of the right inguinal region, an extra-abdominal, oval cyst was found adherent to the spermatic cord within a hernial sac. It was carefully dissected and removed. On exploration of the left inguinal canal, the cyst wall, daughter cysts, and the laminated membrane were found inside a sac. All these contents were removed from the inguinal canal. Pathology of the cysts revealed *Echinococcus multilocularis*.

The patient recovered and left the hospital in good general health. He was also given albendazole 300 mg daily for 30 days.⁸⁻¹⁰

DISCUSSION

Hydatid cyst of the inguinal canal is very rare. Authentic textbooks on parasitology⁴ and surgery^{5,7} have not reported such a case until today. To our knowledge and after an enthusiastic literature review, no clear-cut case has been described in which the site of disease was the inguinal canal.

This case illustrates the difficulty encountered in making a diagnosis of hydatid cyst at unusual sites when clues are not recognized. The differential diagnosis of such swellings includes inguinal hernia, enlarged lymph nodes, psoas abscess, encysted hydrocele of the spermatic cord, tumors of the cord, hydatid of Morgagni, aneurysm of the external iliac artery, distended psoas bursa and effusions into the hip joint. Sonography of scrotal swellings can differentiate hydatid cyst from other swellings of the cord. Lipomas of the spermatic cord can mimic an indirect hernia sac exactly.¹¹

This patient was previously operated for hepatic hydatid cyst, and the presence of a hernial sac in the inguinal canal indicates that in this case cystic fluid might have entered the inguinal hernial sac, which later caused the hydatid cyst.

Spillage of hydatid fluid and peritoneal infestation by its contents was considered a possibility in such a case. Therefore, an accurate and detailed search for daughter cysts was undertaken by an abdominal CT scan.

Drug therapy should be considered as an adjunct rather than an alternative to surgery and administered for at least a few weeks postoperatively. Drugs have been continued from a few months to several years in different reports.¹²

Prolonged use of albendazole, as the sole mode of therapy, has been advised only when there is disseminated hydatidosis with no chance of cure by surgery, the patient is not a good candidate for surgery, or the cysts are surgically unapproachable.

REFERENCES

1. Matson RM, Richard MD, Smyth JD: Hydatidosis, a global problem of increasing importance. *Bull WHO* 55: 499-507, 1977.
2. Doty E, Tompkins RK: Management of cystic disease. *Surgical Clinics of North America*, April, 290-294, 1989.
3. Radford AJ: Hydatid disease, In: Weatherall DJ, Ledinghom JGG, Warrel DA, (eds.), *Oxford Textbook of Medicine*. Second ed., Vol. 1, Oxford: Oxford University Press, p. 563, 1988.
4. John DT: *Medical Parasitology*. Philadelphia: W.B. Saunders, pp. 246-254, 1992.
5. Schwartz SI: Hydatid cysts, In: Schwartz SI, Shirès GT, Spencer FC, (eds.), *Principles of Surgery*. 6th ed., Vol. 2, New York: McGraw-Hill, p. 1332, 1994.
6. Bastani B, Dehdashti F: A review on hydatid cyst of the liver with a brief report of 126 cases from Fars province. *MJIRI* 6(4): 257-264, 1993.
7. Saidi F: *Surgery of Hydatid Disease*. Philadelphia: W.B. Saunders, p. 286, 1986.
8. Morris DL, Dykes PW, Dickson B, Marriner SE, Bogan JA, et al: Albendazole in hydatid disease. *Br Med J* 286: 103-104, 1983.
9. Schantz PM: Effective medical treatment of hydatid disease. *JAMA* 253: 2095, 1985.
10. Webster LT Jr: Drugs used in the chemotherapy of helminthiasis. In: Goodman Gilman A, Rall TW, Nies AS, Taylor P (eds.), *The Pharmacological Basis of Therapeutics*. 8th ed., Vol. II, New York: Pergamon Press, p. 964, 1991.
11. Wantz GE: Abdominal Wall Hernias, In: Schwartz SI, Shires GI, Spencer FC, (eds.), *Principles of Surgery*. 6th ed., New York: McGraw-Hill, pp. 1519-1520, 1994.
12. Eyail L, Zveibil F, Stamler B: Anaphylactic shock due to rupture of a hepatic hydatid cyst. *J Ped Surg* 260: 217-218, 1991.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200

201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300

301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400

401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500