

Effectiveness of adenoidectomy on tympanostomy tubes retention duration

Mohammad Farhadi¹, Hadi Ghanbari², Farzad Izadi³, Maxine S. Eikani⁴,
Seyed Kamaran Kamrava⁵

Department of Otolaryngology, Head and Neck Surgery, ENT&HNS Research Center. Rasoul Akram Hospital, Tehran University of Medical Sciences. Tehran, Iran.

Received: 22 Dec 2010

Revised: 10 Aug 2011

Accepted: 20 Sep 2011

Abstract

Background: The children with middle ear effusion need repeated re-tympanostomies. Adenoidectomy is an effective surgical intervention in the management of chronic otitis media with effusion in conjunction with insertion of tympanostomy tubes (TTs). To find out whether TTs in different positions decrease the rate of re-tympanostomies study was done.

Methods: The present study retrospectively evaluated the effectiveness of adenoidectomy on retention of Shepard TTs in antero-inferior quadrant (AIQ) and postero-inferior quadrant (PIQ) with chronic, persistent or recurrent otitis media. Eighty-five children (one-hundred and seventy ears) underwent bilateral myringotomy and TTs placement with and without adenoidectomy with informed consent.

Results: According to the TTs retention duration rate, there was a significant difference between adenoidectomy and non-adenoidectomy groups in AIQ.

Conclusion: It was concluded that TTs placement in the AIQ in conjunction with adenoidectomy showed better improvement and prolonged ventilation. This study suggests that adenoidectomy is an effective surgical intervention in the management of otitis media especially when it is performed in conjunction with insertion of TTs. This significantly decreases tube extrusion rate especially in an AIQ, which might be due to improving eustachian tube function that consequently reduces repeated otitis media.

Keywords: chronic secretory otitis media, adenoidectomy, tympanostomy tube.

Introduction

Obstruction of the Eustachian tube orifices is an important indication for adenoidectomy [1]. Thus resultant middle ear effusion is prevented by the elimination of the tubal obstruction. Conductive deafness associated

with refractory serous or secretory otitis media is likewise an indication for adenoid surgery in conjunction with TTs insertion [2]. Myringotomy with insertion of TTs is currently the most common surgical procedure performed in children [3]. The length of time the middle ear is ventilated by functioning

1. Professor of Otolaryngology-Head and Neck surgery. Department of Otolaryngology Head and Neck Surgery, ENT&HNS Research Center. Rasoul Akram Hospital, Tehran University of Medical Sciences. Tehran, Iran. farhadi@ent-hns.org

2. (**Corresponding author**) MD. Otolaryngologist, Head and Neck Surgeon. Department of Otolaryngology Head and Neck Surgery, ENT&HNS Research Center. Rasoul Akram Hospital, Tehran University of Medical Sciences. Sattarkhan Avenue, Tehran, Iran. Tel: +98 21 66504294 ghanbari_md@tums.ac.ir

3. Associated Professor of Otolaryngology, Head and Neck Surgery. Department of Otolaryngology Head and Neck Surgery, ENT&HNS Research Center. Rasoul Akram Hospital, Tehran University of Medical Sciences. Tehran, Iran. izadi@ent-hns.org

4. Pediatrics resident, Department of Pediatrics, Mercer University School of Medicine (Medical Center of Central Georgia). maxineseikani@yahoo.com

5. Assistant Professor of Otolaryngology, Head and Neck Surgery. Department of Otolaryngology Head and Neck Surgery, ENT&HNS Research Center. Rasoul Akram Hospital, Tehran University of Medical Sciences. Tehran, Iran. kamrava@ent-hns.org

TTs varies on the basis of the epithelial migration pattern of the tympanic membrane [4]. Children with middle ear effusion are at higher risk of significantly increases in both the tube extrusion rate and the need for subsequent re-tympanostomies [5]. This study explored the efficacy of adenoidectomy on the duration of tympanostomy tube vs. location in children with persistent or recurrent otitis media who had not previously undergone tube placement [6].

Methods

Eighty-five children underwent bilateral TTs insertion in conjunction with and without adenoidectomy and TTs insertion; 42, 43 respectively, in Rasoul Akram hospital (Tehran university of medical sciences) with informed consent. The indication for surgery was bilateral symmetric chronic otitis media with effusion that documented by Type B tympanogram with hearing level less than 35db. Effusion was not been responsive to medical treatment for more than 3 months. In adenoidectomy and non-adenoidectomy groups 15, 11 patients with serous or non-secretory were excluded and 27, 32 patients were mucoid and TTs inserted in different position respectively. Patient’s age ranges were 2-12 year. General anesthesia used in both groups, and all external ear canals were mechanically cleaned and irrigated with saline solution followed by a radial incision, and aspiration of the middle-ear mucoid effusions which were inclusion criteria for TTs

insertion. The Shepard TTs with an internal diameter of 1mm had been used. Patients were to examined 2-4 weeks after tube placement and then at 3 months intervals, also tympanometry and otoscopy were performed. The date of extrusion was considered to be the date of the examination on which the tube was not open or extruded by tympanometry.

Statistical Analysis

Qualitative and quantitative variables are reported as frequency percentage and mean± standard deviation (SD), respectively. Kolmogorov-Smirnov Test was used to investigate if the variables had normal distribution and then Paired and independent sample t-tests were applied where appropriate. One way ANOVA for comparison between age-groups was used. A two-tailed P-value was considered statistically significant.

Results

Eighty-five children underwent bilateral myringotomy and TTs insertion in conjunction with and without adenoidectomy; 42 and 43 cases respectively, and aspiration of the middle-ear effusion. In adenoidectomy and non-adenoidectomy groups 15, 11 patients who were serous or non-secretory were excluded, and 27, 32 were mucoid and TTs inserted in different positions respectively. Thirty-five patients with seventy ears were included for evaluation; eighteen patients of adenoidectomy group and seventeen pa-

Table 1. The number of the ears operated in right (AIQ) and left (PIQ) both in adenoidectomy and non-adenoidectomy groups.

Duration	Adeno	Number of patients	Mean	Std.Deviation	Independent t-test
Left	Adenoidectomy	18	10	4.102	t=1.497 P=0.144
	Non-adenoidectomy	17	7.53	5.591	
Right	Adenoidectomy	18	20.39	4.104	t=5.105 p<0.0001
	Non-adenoidectomy	17	10.71	6.854	

Table 2. The effect of adenoidectomy on TTs duration in left ears in postero-inferior quadrant (PIQ) group in comparison with right ears in antero-inferior quadrant (AIQ) group showed more prolongation in the later.

	N	Mean duration	Std Deviation	Paired T-test
TTs Left	35	2.628571	1.086974	t= -4.65
TTs right	35	3.714286	1.226459	P<0.0001

Duration	Age range	Number	Mean	Std.Deviation	One Way ANOVA test
Left	<2year	1	14		F:0.356
	2-5year	15	8.6	5.289	
	6-12year	18	8.67	4.971	P:0.785
	>12year	1	9		
	Total	35	8.8	4.969	
Right	<2year	1	3		F:1.377
	2-5year	15	14.67	6.532	
	6-12year	18	17.11	7.828	P:0.268
	>12year	1	18		
	Total	35	15.69	7.391	

tients of non-adenoidectomy group. Patients' age range was between 2-12 year including 18 (51.4%) female and 17 (48.6%) male. Sixteen patients were 2-5 year and nineteen patients were 6-12 year, 45.7% and 54.3% respectively. The number of left ears with TTs duration (PIQ) in adenoidectomy group were 18 with mean: $10(\pm 4.102)$ months and 17 in non-adenoidectomy group with mean: $7.53(\pm 5.591)$ which was not statistically significant (Fig 1). The numbers of right ears TTs duration (AIQ) in adenoidectomy group were 18 with mean $20.39(\pm 4.104)$ months, and 17 in non-adenoidectomy group with mean $10.71(\pm 6.854)$ months was statistically significant ($p < 0.05$, Fig. 2). The patients' age range were <2 year, 2-5 year, 6-12 year, > 12 year and including cases in groups were 1, 15, 18, 1 respectively, there was no difference between female and male in right and left ears TTs du-

ration and also no difference between age groups (Table 3). Independent t-test showed significant difference between duration in AIQ and PIQ ears groups for adenoidectomy in comparison to non-adenoidectomy patients ($p < 0.05$) (Fig1, Table1).

Discussion

Obstructed airway by enlarged adenoid has already been mentioned as an indication for adenoidectomy [2]. Eustachian tube orifices obstruction by adenoid tissue is another important indication for adenoidectomy [1]. Conductive deafness associated with refractory secretory otitis media due to obstruction of the eustachian tube orifices is likewise to indicate for adenoid surgery in conjunction with TTs insertion [2]. Sustained middle ear ventilation could be achieved by selecting the appropriate tube, site and incision [7].




Fig.1. The posteroinferior quadrant (PIQ) TTs placement retention in adenoidectomy and non-adenoidectomy groups was almost the same.

Stinson found clues to the premature extrusion of a foreign body in the tympanic membrane; information went essentially unrecognized for many years [8]. Turner found

Conclusion

The location of a myringotomy for insertion of a (TTs) should not be determined to expect increase duration of action, and dif-

The effect of adenoidectomy on PETs duration time in anteroinferior quadrant (AIQ) right ears group




Fig.2. The antero-inferior quadrant (AIQ) TTs placement retention in the adenoidectomy group is more prolonged than non-adenoidectomy group.

that the epithelium of the anteroinferioer quadrant and of both posterior quadrants migrated toward the posterior border of the tympanic membrane [9]. To evaluate the efficacy of adenoidectomy on the duration of TTs placement in children with persistent or recurrent otitis media who had not previously undergone tube placement was determined [6]. The children with middle ear effusion are at higher risk of repeated post-TTs otorrhea episodes during the first insertion of TTs significantly increases both the tube extrusion rate and the need for subsequent re-tympanostomies [5]. The different tube designs should be considered when prolonged middle ear ventilation is desired [10]. This study concluded that TTs placement on the AIQ in conjunction with adenoidectomy remained more functional than the left (Fig.2), and showed better improvement and prolonged ventilated and decreased episode of otorrhea. Although this significantly decreased tube extrusion rate especially in an AIQ (Fig.2), which might be due to improving eustachian tube function [11], but it needs further investigations.

ferent tube designs should be considered when prolonged middle ear ventilation is desired. This study concluded that the TTs placement in the AIQ in conjunction with adenoidectomy showed better improvement and prolonged ventilation and the difference was statistically significant $p < 0.05$ Kolmogorov-Smirnov Test. It is concluded that the TTs placement on the AIQ in conjunction with adenoidectomy remained more functional than the PIQ.

Acknowledgement

Authors wish to thank Mrs. Fatemeh Hosseini for her helpful statistical analysis and Dr ghavami for his efforts in edition.

References

1. Bluestone, C.D, Cantekin, EI, and Beery, Q C: Certain effects of Adenoidectomy on Eustachian tube function, *Laryngoscope* 85:113-127, 1975.
2. Kornblut A, Kornblut AD, Paparella, Shumbrick, Gluckman, Meyerhoff, *Otolaryngology Head and neck surgery*, WB. Saunders Company, 1991; Vol 3, pp. 2150.
3. Sedlmair B; Jivanjee A; Gutzler R; Huscher D; Jovanovic S. Duration of middle ear ventilation after

laser myringotomy with CO2 laser otoscope. *Otoscan HNS* 2001; 49(6):447-453.

4. Valtonen H, Tuomilehto H, Qvarnberg Y, Nuutinen J. A 14-year prospective follow-up study of children treated early in life with tympanostomy tubes: Part 1: Clinical outcomes. *Arch Otolaryngol Head Neck Surg.* 2005 Apr;131(4):293-8.

5. Boston M, McCook J, Burke B, Derkay C. Incidence of and risk factors for additional tympanostomy tube insertion in children. *Arch Otolaryngol Head and neck surg* 2003 mar; 129(3) 293-6.

6. Casselbrant ML, Mandel EM, Rockette HE, Kurs-Lasky M, Fall PA, Bluestone CD, Adenoidec-tomy for otitis media with effusion in 2-3-year-old children. *Int J Pediatr Otorhinolaryngol* 2009 Dec;73(12):1718-24. Epub 2009 Oct 12.

7. Armstrong BW, MD. Prolonged middle ear ven-tilation: The right tube in the right Place. *Ann Otol*

Rhinol Laryngol 1983; 92:582-6.

8. April MM, Portella RR, Orobello PW Jr, Naclerio RMTympanostomy tube insertion: anterosuperior vs. anteroinferior quadrant. *Otolaryngol Head Neck Surg.* 1992 Mar;106(3):241-2.

9. van den Aardweg MT, Schilder AG, Herkert E, Boonacker CW, Rovers MM. Adenoidectomy for otitis media in children. *Cochrane Database Syst Rev.* 2010 Jan 20;(1):CD007810.

10. McDonald S, Langton Hower CD, Nunez DA. Grommets (ventilation tubes) for recurrent acute otitis media in children. *Cochrane Database Syst Rev.* 2008 Oct 8;(4):CD004741.

11. Song CM, Park MH, Kim YH, Lee JH, Factors affecting the extrusion rate of ventilation tubes. *Clin Exp Otorhinolaryngol.* 2010 Jun;3(2):70-5. Epub 2010 Jun 30.