

Effects of capsaicin in low doses and high doses on the small intestine motor response in wild-type and TRPV1 knockout mice

Brief Communication

Contractile activity of the small intestine is coordinated by interplay of myogenic, neural, and chemical controls [1]. Both primary afferent neurons (extrinsic) and intrinsic enteric neurons are present in the small intestine and capsaicin (vanilloid, 8-methyl-N-vanillyl-6-noneamide), the pungent ingredient of chili peppers, specifically acts on extrinsic primary afferent but not intrinsic enteric neurons [2-5]. The transient receptor potential V1 (TRPV1) or formerly known as VR1, is activated by capsaicin [3]. The aim of this study was to examine

the role of TRPV1 receptors in the generation of intestinal motor activity and its reflex modulation by intraluminal distension and capsaicin. Experiments were performed on mice in which the TRPV1 gene had been disrupted (KO) using standard gene targeting techniques [6] and wild-type (WT) littermates. Jejunal contractile activity was recorded from in vitro segments of jejunum 4-5 cm in length. When distended to 2-3 cm H₂O the segments generated regular migrating motor complexes (MMCs) recorded as changes in intraluminal pressure. Motility of isolated segments of mouse small intestine was recorded using a technique described previously [7,8]. Distension of isolated segments caused




Fig. 1A. Representative traces show motor complexes (MCs) in the isolated jejunum from wild-type and TRPV1 knockout (-/-) mice. Motor pattern was similar in both animals. There was no difference in the features of the MCs between two groups. B. Bar charts illustrate the absence of any difference in the baseline interval and amplitude MCs between wild type (WT) and knockout (KO) jejunum mice (n=10).




Fig. 2 A. Representative traces show the effects of low and high doses of TRPV1 agonist, capsaicin on typical MCs evoked by distension in the wild-type mice jejunum. B. Representative traces show effects of low and high doses of capsaicin on MCs evoked by distension in the TRPV1 KO. C. Concentration response curves versus the effects of TRPV1 receptor agonist, capsaicin on interval MCs in the WT and KO mice. Capsaicin responses are expressed as change in intervals and each point represents the mean (\pm SEM) of values. There was significant difference between capsaicin effects in low dose (1-100nM) in the WT and KO mice (* $P < 0.05$, $n=5$) but no significant difference between WT and KO mice found in the higher concentrations of capsaicin ($n=5$).

regular, propagating motor complexes (MMCs) at intervals of approximately 3 minutes, separated by periods of quiescence during which inhibitory motor activity occurs (Fig.1). The MMCs were characterized by periods of phasic contractions that migrate in an aboral direction. It seems that the pattern and the pharmacological properties of motor activity described here in the TRPV1 wild-type (WT) and TRPV1 knockout (KO) mice are equivalent to the migrating motor complexes (MCs) in other species described by others [7-11]. The TRPV1 KO mice had normal contractions and showed

no obvious difference with WT animals. Thus the periodic generation of regular MMCs in isolated mouse jejunum was not causally related to the TRPV1 receptors.

Capsaicin (1-100nM), a ligand for the TRPV1 receptor, caused a dose dependent inhibition of motility manifested as an increase in the interval between MMCs in the wild-type animal only (Fig.2). This indicates that capsaicin acts via activation of TRPV1 receptors that are possibly located on the peripheral terminals of afferent nerves in the mice jejunum. This observation is consistent with the findings that TR-

PV1 receptor has been identified on both intrinsic and extrinsic neurons of the GI tract and both vagal and spinal afferents of the GI tract can be activated by capsaicin [12-15].

At higher doses of capsaicin (1-100 μ M), the periodic MCs were replaced by tonic increases in pressure upon which were superimposed continuous phasic contractions. This stimulation occurred in both KO and WT mice. It can be suggested that capsaicin acts probably by sensitizing sensory neurons through an unknown site, probably a subtypes of an as yet unknown TRP receptors. However, the relative lack of knowledge about the properties of other TRP receptor family members makes it difficult to make inferences regarding the role of these receptors in the high-dose capsaicin effects described here.

TRPV1s are expressed in sensory neurons and also in nonneuronal tissues [15-20]. Although; there has as yet been no direct evidence that functional TRPV1 receptor is expressed in non-neural tissues in gut. Some results suggest that the capsaicin-induced relaxation is associated with the direct inhibitory action intracellularly on the voltage-operated Ca⁺⁺ channels [21]. Studies of the effect of capsaicin on smooth muscle contractions are controversial and have shown both contraction and relaxation in various preparations [17-23]. Therefore, on the basis of these results, it appears to be a fundamental difference may exist between the mechanism of capsaicin in the low dose and high dose and suggested that additional mechanisms rather than the TRPV1 were involved in the mouse.

Acknowledgement

The author wish to thank Professor David Grundy from Department of the Biomedical Science at University of Sheffield in UK for his kind support and supervision and also the Ministry of Health and Medical Education of Iran for financial support of this work.

Reza Rahmati¹, MSc, PhD

Department of Physiology, Gorgan Faculty of Medicine, Golestan University of Medical Science, Gorgan, Iran

References

- Otterson MF, Sarr MG. Normal physiology of small intestinal motility. *Surg Clin North Am* 1993. 73(6): 1173-1192.
- Bevan S, Szolcsanyi J. Sensory neuron-specific actions of capsaicin: mechanisms and applications. *Trends Pharmacol Sci* 1990. 11(8): 330-333.
- Holzer P. Capsaicin: cellular targets, mechanisms of action, and selectivity for thin sensory neurons. *Pharmacol Rev* 1991. 43 (2): 143-201.
- Jancso G. Pathobiological reactions of C-fibre primary sensory neurones to peripheral nerve injury. *Exp Physiol* 1992. 77(3): 405-431.
- Holzer P, Lippe IT, Tabrizi AL, Lenard LJr, Bartho L. Dual excitatory and inhibitory effect of nitric oxide on peristalsis in the guinea pig intestine. *J Pharmacol Exp Ther* 1997. 280 (1): 154-161.
- Davis JB, Gray J, Gunthorpe MJ, Hatcher JP, Davy PT, Overend P, Harries MH, Latcham J, Clapham C, Atkinson K, Hughes SA, Rance K, Grau E, Harper AJ, Pugh PL, Rogers DC, Bingham S, Randall A, Sheardown SA. Vanilloid receptor-1 is essential for inflammatory thermal hyperalgesia. *Nature* 2000; 405(6783): 183-187.
- Abdu F, Hicks GA, Hennig G, Allen JP, Grundy D. Somatostatin sst (2) receptors inhibit peristalsis in the rat and mouse jejunum. *Am J Physiol Gastrointest. Liver Physiol* 2002. 282 (4) G624-G633.
- Bian X, Ren J, Vries MD, Birthe S, Cockayne DA, Ford APDW, Galligan JJ. Peristalsis is impaired in the small intestine of mice lacking the P2X3 subunit. *J Physiol* 2003. 551: 309-322.
- Mule F, D'Angelo S, Tabacchi G, Amato A, Serio R. Mechanical activity of small and large intestine in normal and mdx mice: a comparative analysis. *Neurogastroenterol Motil* 1999. 11(2): 133-139.
- Bercik P, Bouley L, Dutoit P, Blum A. L, Kucera P. Quantitative analysis of intestinal motor patterns: spatiotemporal organization of nonneural pacemaker sites in the rat ileum. *Gastroenterology* 2000. 119 (2): 386-394.
- Bush TG, Spencer NJ, Watters N, Sanders KM, Smith TK. Spontaneous migrating motor complexes oc-

cur in both the terminal ileum and colon of the C57BL/6 mouse in vitro. *Auton. Neurosci* 84(3): 162-168, 2000.

12. MacNaughton WK, Van Sickle MD, Keenan, CM, Cushing K, Mackie K, Sharkey KA. Distribution and function of the cannabinoid-1 receptor in the modulation of ion transport in the guinea pig ileum: relationship to capsaicin-sensitive nerves. *Am. J Physiol Gastrointest. Liver Physiol* 2004.286(5): G863-G871.

13. Ward SM, Bayguinov J, Won KJ, Grundy D, Berthoud HR. Distribution of the vanilloid receptor (VR1) in the gastrointestinal tract. *J Comp Neurol* 2003. 465 (1):121-135.

14. Ichikawa H, Sugimoto, T. The co-expression of VR1 and VRL-1 in the rat vagal sensory ganglia. *Brain Res* 2003. Vol. 980 (2): 293-296.

15. Knotkova H, Pappagallo M, Szallasi A. Capsaicin (TRPV1 Agonist) therapy for pain relief: farewell or revival? *Clin J Pain* 2008. 24(2):142-54.

16. Poli-Neto OB, Filho AA, Rosa e Silva JC, Barbosa Hde F, Candido Dos Reis FJ, Nogueira AA. Increased capsaicin receptor TRPV1 in the peritoneum of women with chronic pelvic pain. *Clin J Pain*.25 (3):218-22.

17. Sim JH, Kim YC, Kim SJ, Lee SJ, Suh SH, Jun JY, So I, Kim KW. Capsaicin inhibits the voltage-operated calcium channels intracellularly in the antral circular myocytes of guinea-pig stomach. *Life Sci* 2009. 68(21): 2347-2360, 2001.

18. Kullmann FA, Shah MA, Birder LA, de Groat WC. Functional TRP and ASIC-like channels in cultured urothelial cells from the rat. *Am J Physiol Renal Physiol* 2009. 296(4):F892-901.

19. Yamada T, Ugawa S, Ueda T, Ishida Y, Kajita K, Shimada S. Differential localizations of the transient receptor potential channels TRPV4 and TRPV1 in the mouse urinary bladder. *J Histochem Cytochem* 2009. 57(3):277-87.

20. Spencer NJ, Kerrin A, Singer CA, Hennig GW, Gerthoffer WT, McDonnell O. Identification of capsaicin-sensitive rectal mechanoreceptors activated by rectal distension in mice. *Neuroscience* 2008. 153(2):518-34.

21. Sim JH, Kim YC, Kim SJ, Lee SJ, Suh SH, Jun JY, So I, Kim KW. Capsaicin inhibits the voltage-operated calcium channels intracellularly in the antral circular myocytes of guinea-pig stomach. *Life Sci* 2001. 68(21): 2347-2360.

22. Kark T, Bagi Z, Lizanecz E, Posztor ET, Erdei N, Czikora A, Papp Z, Edes I, Porszasz R, Tóth A. Tissue-specific regulation of microvascular diameter: opposite functional roles of neuronal and smooth muscle located vanilloid receptor-1. *Mol Pharmacol* 2008. 73(5):1405-12.

23. Huang W, Wang H, Galligan JJ, Wang DH. Transient receptor potential vanilloid subtype 1 channel me-

diated neuropeptide secretion and depressor effects: role of endoplasmic reticulum associated Ca²⁺ release receptors in rat dorsal root ganglion neurons. *J Hypertens* 2008. 26(10):1966-75.