

## ENDEMIC GOITER AND HYPOTHYROIDISM IN ORANG ASLI AND MALAYS IN PENINSULAR MALAYSIA

OSMAN ALI, B.A.K. KHALID\*, T.T. TAN\*, O. SAKINAH\*,  
L.L. WU<sup>+</sup>, AND M.L. NG<sup>‡</sup>

*From the Department of Community Health, \*Department of Medicine,  
\*Department of Paediatrics and ‡Department of Biochemistry,  
Medical Faculty, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia.*

### ABSTRACT

The study was conducted to determine the prevalence of goiter and hypothyroidism in two groups of the population at various stages of social development and to determine its association with malnutrition and dietary intake. Aboriginal inhabitants (Orang Asli), a resettlement rural village (Betau Post), a traditional village in the jungle (Lanai Post) and a modern village settlement near Kuala Lumpur City (Bukit Lanjan) were selected. Three Malay villages with almost similar environments were selected for comparison. The study included house to house interviews, anthropometric, clinical and biochemical assessment of 1419 samples and estimation of iodine concentration in the drinking water of the areas studied. The World Health Organization criteria for classification of goiter were used. It was found that all Orang Asli settlements in rural areas were iodine deficient. The prevalence of goiter and hypothyroidism were higher among Orang Asli at all ages compared to Malays, and increased with remoteness of the areas (20-70% and 20-30% respectively). Apart from iodine deficiency, a high intake of cassava and deficient intake of sea foods and protein diets were also significantly related with high prevalence of goiter. Among children, goiter was associated with the level of thyroid stimulating hormone (TSH) whereas among adults, body mass index (BMI), triiodothyronine and thyroxine levels were predictors for the presence of goiter. In conclusion, goiter and hypothyroidism were more common among Orang Asli compared to Malays, and were associated with rural location and poverty.

**Key Words:** Goiter, Hypothyroidism, Malay, Orang Asli, Malaysia.

*MJIRI, Vol. 9, No. 1, 19-25, 1995.*

### INTRODUCTION

The World Health Assembly in 1990 passed a resolution to prevent and eliminate iodine deficiency disorders (IDD), especially endemic cretinism.<sup>1</sup> Iodine deficiency retards growth and mental development.<sup>2</sup> Because iodine deficiency occurs mostly in developing countries, it is a major problem

#### Correspondence:

Dr. Osman Ali,  
Department of Community Health, Medical Faculty, University  
Kebangsaan Malaysia, Jalan Raja Muda Abdul Aziz, 53000, Kuala  
Lumpur, Malaysia.  
Tel: (603) 4405272, 2923066  
Fax: (603) 2912659

## Endemic Goiter and Hypothyroidism in Malaysia




Fig. 1. Study areas

for their manpower development.

The prevalence of goiter and hypothyroidism varies with geographical, socioeconomic and environmental factors. The prevalence is highest in isolated communities, especially those living in mountainous areas.<sup>3,4,5,6</sup> Approximately 400 million people in Asia are afflicted by iodine deficiency disorders.<sup>7</sup> Dietary intake, age, and racial type were the factors found to be important in predicting the occurrence of endemic goiter in the tropics.<sup>2</sup> Because endemic goiter is not considered a public health problem in Peninsular Malaysia, little attention has been given to it. However, published epidemiological data on the prevalence of goiter and hypothyroidism in Peninsular Malaysia is very limited. Most of the studies were performed in East Malaysia.<sup>3,8,9</sup> A community-based study in a rural population in Sarawak found a goiter prevalence of 4.5-21.4% and 3.1-55.3% in 10-14 year olds.<sup>9</sup> It was also found that the prevalence among the population living in other parts of the country was twice that living in urban areas. However, none of the studies done so far have investigated the prevalence among the aborigines (Orang Asli) of Peninsular Malaysia, nor the Malay people living in various locations ranging from jungle fringes to urban towns.

Thus, the aims of this study were to determine the prevalence of goiter and hypothyroidism among the Orang Asli and Malays from different locations and environmental

conditions with varying stages of modernization and development, in order to examine the factors associated with the occurrence of the disease.

## MATERIAL AND METHODS

### Sampling

A total of 1419 subjects aged 2 years and above were chosen using cluster sampling from different locations in Peninsular Malaysia (Fig. 1). The areas were selected based on degrees of social development and lifestyles. Different sampling strategies were chosen depending on the homogeneity of the population. For Orang Asli subjects, the areas chosen were Lanai Post, Betau Post and Bukit Lanjan. In Lanai Post, almost all eligible subjects from three randomly selected villages were invited to participate in the study. In Betau Post, three clusters comprising 6 villages were randomly selected with a participation rate of more than 85%. In Bukit Lanjan, all eligible subjects were invited, with a participation rate of more than 80%. For Malays, the areas selected were Ulu Sungai, Sungai Koyan and Kg. Kerinci. In Ulu Sungai traditional village, the response rate was more than 95%. In the Sungai Koyan resettlement scheme and Kampong Kerinci urban village, random

Table I: Prevalence of goiter according to age and stage of development.

Stage of development	Age group (years)	Number of subjects	prevalence	Standard error	95% confidence interval
Remote rural:					
Lanai Post	2 - 6	39	20.3	$4.1 \times 10^{-3}$	19.5 - 21.1
	7 - 17	61	55.7	$4.0 \times 10^{-3}$	54.9 - 56.5
	> 17	112	67.3	$2.0 \times 10^{-3}$	66.9 - 67.7
Hulu Sungai	2 - 6	34	2.9	$8.3 \times 10^{-4}$	2.74 - 3.06
	7 - 17	56	17.2	$2.4 \times 10^{-3}$	16.7 - 17.7
	>17	154	25.0	$1.2 \times 10^{-3}$	24.8 - 25.2
Rural:					
Betau Post	2 - 6	55	21.2	$3.0 \times 10^{-3}$	20.6 - 21.8
	7 - 17	68	55.2	$8.6 \times 10^{-3}$	53.5 - 56.9
	>17	137	48.2	$1.8 \times 10^{-3}$	47.8 - 48.6
FELDA Koyan	2 - 6	55	0	0	0
	7 - 17	119	1.7	$1.4 \times 10^{-4}$	1.67 - 1.73
	>17	133	8.1	$5.6 \times 10^{-4}$	7.99 - 8.21
Urban:					
Lanjan	2 - 6	57	0	0	0
	7 - 17	68	4.4	$3.6 \times 10^{-4}$	4.33 - 4.47
	>17	78	24.4	$2.3 \times 10^{-3}$	23.9 - 24.9
Kg. Kerinci	2 - 6	39	0	0	0
	7 - 17	59	1.7	$2.8 \times 10^{-4}$	1.64 - 1.76
	>17	92	6.6	$6.7 \times 10^{-4}$	6.47 - 6.73

samplings of the households were applied and more than 80% participated. Description of study areas and their sociodemographic profiles has been published elsewhere.<sup>10</sup>

### Survey procedures

The survey was conducted between March and April 1991. All the subjects selected were invited to attend the clinic. Neck examination for goiter detection and grading was performed by an experienced endocrinologist in our team. Twenty ml of blood was drawn from subjects aged 7 and above (school children and adults), centrifuged and serum kept frozen at -40°C for biochemical testing. Anthropometric assessment and blood pressure were measured and recorded. Dietary information was gathered using a 24-hour dietary recall and food frequency questionnaire.

Thyroid gland size was classified according to criteria recommended by WHO.<sup>11</sup> Thyroid stimulating hormone (TSH) levels of more than 5 mIU/L were taken as indicative of hypothyroidism.

Thyroxine ( $T_4$ ) and triiodothyronine ( $T_3$ ) were measured by radioimmunoassay and TSH by an immunoradiometric assay using reagents from NETRIA (North East Thames

Regional Immuno Assays).

Random urine samples were collected from 30 subjects in different locations. Samples of drinking water from various sources such as pipe water, river, gravity feed systems and spring water were also collected. The level of iodine was determined using an ashing method based on the Sandell-Kolthoff reaction.<sup>12</sup>

### Statistical Analysis

Statistical analyses were performed using the SAS statistical software release 6.3 (SAS Institute Inc.). In order to compare groups, Chi-square tests and Student's t-tests were used. In both age categories (children and adults), the common factors tested were racial group (Orang Asli), rural location, sex (female), body mass index (BMI) of less than  $15 \text{ kg/m}^2$ , serum albumin level of more than 4.5 g/L, serum cholesterol level of less than 5 mmol/L and presence of splenomegaly. Specifically for children, the factors tested were age of more than 13 years, energy intake of 1200 calories, protein intake of more than 45 g, carbohydrate intake of more than 200 g and fat intake of more than 25 g per day. For adults, age of less than 40 years, energy intake of 1500 calories, protein intake of more than 50 g,

## Endemic Goiter and Hypothyroidism in Malaysia

**Table II: Association between goiter and socio-demographic and nutrition factors among children and adults.**

	Children		Adults	
	Chi-sq	P	Chi-sq	P
Ethnic	62.6	<0.0001	102.9	<0.0001
Urbanity	40.2	<0.0001	31.5	<0.0001
Age (years)	22.6	<0.0001	7.4	<0.025
Sex	8.3	0.016	34.6	<0.0001
BMI(kg/m <sup>2</sup> )	3.3	0.19	23.1	<0.0001
Albumin (g/l)	7.2	0.028	2.1	0.354
Cholesterol(mmol/l)	17.8	<0.0001	32.9	<0.0001
Splenomegaly	17.4	<0.0001	21.6	<0.0001
Energy intake (Cal)	1.0	0.59	7.2	0.027
Protein intake (gm)	0.6	0.76	27.3	<0.0001
CHO intake (gm)	1.7	0.43	0.01	0.99
Fat intake (gm)	1.9	0.38	31.8	<0.0001

Chi-sq = chi square, P is significant at 0.05

CHO = Carbohydrate

carbohydrate intake of more than 250 g and fat intake of more than 25 g per day. Criterion of significance was at  $P < 0.05$ .

### RESULTS

The prevalence of goiter was higher among the Orang Asli compared to Malays in all types of environment or lifestyles (Table I). In remote rural areas, the difference in prevalence of goiter was 3 times in preschool and school-age children, and twice in adults. No difference was observed between remote rural (Lanai Post) and rural (Betau Post) Orang Asli settlements. However, the prevalence was low among Orang Asli children in an urban settlement (Bukit Lanjan) and Malays in FELDA (Sungai Koyan) and urban village (Kg. Kerinci).

Hypothyroidism (TSH levels of more than 5 mIU/L) was found in 28.4% of Orang Asli compared to 5.3% among Malays. The highest prevalence was observed in Lanai Post of which 40.4% (23/57) of school-age children and 20.5% of adults were suffering from the illness (Fig. 1). Hypothyroidism was found among 28% (21/75) of all children with goiter and 75% of children with visible goiter. In adults, only 16.4% (36/219) of goitrous patients had hypothyroidism and the proportion was also higher in patients with visible goiters (19.7%). Hypothyroidism among non-goitrous subjects was 11.8% (34/287) and 8.8% (43/488).

There was little variability in the levels of  $T_3$  between Orang Asli and Malays. However, there was a significant difference in the median level of thyroxine in children, young adults and elderly between the two ethnic groups. For TSH, a marked significant difference was observed between

Orang Asli and Malays at all age groups. Orang Asli had higher TSH levels compared to Malays.

Goiter and hypothyroidism were significantly associated with ethnic type, being more common in Orang Asli (Table II and III). It was also associated with remoteness (rural) of the area. The overall prevalence of goiter in rural areas was 25.4% in children and 35.8% in adults compared to 1.9% and 15.5% respectively in urban areas. Low education, income of less than RM250 per month and light daily activities were also risk factors associated with the occurrence of goiter.

Goiter and hypothyroidism were not associated with BMI in children but were strongly related in adults (Table II and III). Goiter was found in 48.8% of malnourished adults compared to 29.3% in well nourished ones. The prevalence was 22.5% and 10.6% respectively for hypothyroidism.

Nutrient intakes were not associated with goiter in children. However, energy, protein and fat intake were significantly related with occurrence of goiter in adults. Low intake of these nutrients were associated with a high risk of developing goiter (Table II). A high serum albumin was significantly associated with the occurrence of goiter in children but a low serum cholesterol showed a significant relationship with the occurrence of goiter in children as well as in adults (Table II). In terms of food frequency, the study found that frequent consumption of cassava and deficient intake of sea foods and proteins in the diet were significantly associated with goiter (Table IV).

Goiter and hypothyroidism were significantly associated with age and gender. The study found that the prevalence was higher in females compared to males in both children and adults. For goiter, the prevalence was 19.4% vs 14.3% in children and 41.7% vs 20.7% in adults.

Using multiple regression, goiter in children was asso-

**Table III: Association between hypothyroidism and socio-demographic and nutrition factors among children and adults.**

	Children		Adults	
	Chi-sq	P	Chi-sq	P
Ethnic	62.6	<0.0001	102.9	<0.0001
Age (years)	22.6	<0.0001	7.4	<0.025
Sex	8.3	0.016	34.6	<0.0001
BMI(kg/m <sup>2</sup> )	3.3	0.19	23.1	<0.0001

**Table IV: Association between goiter and frequency of food intake among children and adults.**

Food items	Children		Adults	
	Chi-sq	P	Chi-sq	P
cassava	54.8	<0.0001	83.0	<0.0001
squid	23.4	<0.0001	24.7	<0.0001
cockles	22.6	<0.0001	30.6	<0.0001
chicken	14.3	<0.0002	25.5	<0.0001
eggs	37.3	<0.0001	56.8	<0.0001

\*for cassava the tested categories were from the most frequent to the least frequent intake.

ciated with the level of thyroid stimulating hormone (TSH) whereas body mass index (BMI), T<sub>3</sub> and T<sub>4</sub> levels were predictors for the presence of goiter among adults. The regression model for predicting the occurrence of goiter in children was GOITER= 0.06 AGE + 0.19 GENDER - 0.45 ETHNIC - 0.11 LOG TSH - 0.02 (F= 31.4, P= 0.0001, r<sup>2</sup>= 0.28), whereas in adults GOITER= 4.47 + 0.37 GENDER - 0.48 ETHNIC - 0.03 BMI - 0.51 LOG T<sub>3</sub> - 0.76 LOG T<sub>4</sub> (F= 43.1, P= 0.0001, r<sup>2</sup>= 0.25).

There were significant differences in the levels of urinary iodine excretion in different areas. The levels of urinary iodine in the remote areas of Lanai Post was 2.9 ± 7.6 µg/dl and Betau Post was 1.9 ± 1.2 µg/dl compared to 5.4 ± 1.7 µg/dl in Hulu Sungai village, 2.9 ± 2.0 µg/dl in FELDA Sungai Koyan and 7.3 ± 3.4 µg/dl in urban areas. The iodine level of drinking water in remote areas was 2.3 µg/dl compared to 2.8 µg/dl in Hulu Sungai village, 4.2 µg/dl in FELDA and 1.5 µg/dl in urban areas.

## DISCUSSION

This study found the prevalence of goiter to be higher among Orang Asli aborigines compared to Malays. The prevalence among aborigines increased with age and was highest among females. The findings were similar to that found by Ogiharain Sarawak.<sup>13</sup> The prevalence was maximal between the age groups 7-12 and 13-17 years old. This

corresponded to maximum secretion of TSH and thyroid gland growth.<sup>14</sup> The highest prevalence rates were in the interior aborigine settlements and Malay traditional village; the prevalence decreased as the level of social development increased. Similar findings were observed in other studies.<sup>15,16</sup> This also correlated well with the urinary iodine levels. In most areas endemic goiter, insufficient intake of iodine through food and drinking water has been shown to play a part and most of these areas were iodine deficient areas.<sup>2,7</sup> In the present study, the aboriginal settlements of Lanai Post and Betau Post had moderate endemic goiter (prevalence of goiter 20-50% with urinary iodine levels of 2.0-5.0 µg/dl) whereas the traditional Malay village had mild endemic goiter (prevalence of goiter 10-30% and urinary iodine levels of 5.0-10.0 µg/dl). Thus, the prevalence of goiter is in inverse ratio to the urinary iodine. Changes in prevalence had been shown in some communities to be due to an increase in the dietary iodine intake through improved social and economic conditions even before any iodine prophylaxis had been started.<sup>2</sup> This was supported in the present study. Even though the iodine level in drinking water was low in the city, there was no endemic goiter (even among aborigines living near the city), probably because town dwellers have access to manufactured and processed foods and an iodine rich diet such as sea foods whereas the rural people depend entirely on local products.

There was no association between malnutrition and goiter among preschool children and children 7-17 years

## Endemic Goiter and Hypothyroidism in Malaysia

old. Among school children, age, gender, ethnic type and TSH levels were predictors for the occurrence of goiter after controlling the other factors. Goiter is a consequence of excessive TSH secretion and the effect of TSH being reinforced at the level of thyroid cells by the depletion of thyroidal iodine stores.<sup>11,17,18</sup> However, among adults, apart from gender, ethnic type, and BMI, levels of  $T_3$  and  $T_4$  were significantly associated with goiter. Intake of energy, protein and fat was also associated with goiter in adults. TSH levels were not associated with goiter prevalence in adults. In adults, the chronic state of iodine deficiency leads to functional alterations due to thyroid hyperplasia.<sup>19</sup> The level of circulating TSH is enhanced and the thyroidal iodine stores are progressively depleted. The body maintains homeostasis by adaptive mechanisms, synthesizing and secreting  $T_3$  at the expense of  $T_4$ .<sup>20,21,22</sup> Serum  $T_3$  and  $T_4$  levels were also influenced by malnutrition. In chronic malnutrition, the thyroid gland is not responsive to TSH stimulation.<sup>23</sup> In this study, chronic iodine deficiency and malnutrition were both affecting the size of the thyroid gland and the synthesis of thyroid hormones.

The prevalence of hypothyroidism (TSH of more than 5 mIU/L) was higher among aborigines and corresponded with the prevalence of goiter and urinary iodine levels. However, no cretin patients were found. This finding was also similar to that found by Ogihara in Sarawak.<sup>13</sup> The question is whether aborigines and other indigenous peoples in Malaysia are resistant to cretinism. The distribution patterns of hypothyroidism were similar to that of goiter. Among children, the prevalence was not associated with age, sex, or nutritional status. Among adults, the prevalence of hypothyroidism was higher among males and malnourished subjects. This probably indicated the relative unresponsiveness of the male thyroid gland to TSH stimulation.

This study also found that other important factors may have contributed to endemic goiter. Apart from lack of iodine in drinking water, frequent intake of cassava root leaves (a main staple food in rural communities, especially aborigines), less intake of iodine-rich foods such as sea foods, and a low protein diet were various factors significantly contributing to the occurrence of goiter especially in rural areas. Malnutrition and malabsorption may potentiate the deficiency. Iodine intervention may not be effective in this situation because of poor iodine absorption from the gut, inhibition by thiocyanates from cassava, poor thyroid hormone production due to malnutrition, etc. The use of thyroxine as supplement therapy as a short-term measure may be beneficial. The ultimate objective is still to eliminate the condition through integrated primary health care by proper nutrition and elevation of health status.

### ACKNOWLEDGEMENT

This work was supported by a grant from the IRPA

program no.03-07-03-051. The authors gratefully acknowledge the support of the Department of Aboriginal Affairs and cooperation of the RIA laboratory technicians for analysing the samples.

### REFERENCES

1. ICCIDD: World Health Assembly urges elimination of IDD by the year 2000. IDD Newsletter 6(2): 1-8, 1990.
2. Stanbury JB, Hetzel BS: Endemic Goiter and Endemic Cretinism. Wiley Medical, New York, 1980.
3. Maberly GF: The significance of endemic goiter in Sarawak, Borneo. A thesis submitted to the University of New South Wales for the degree of Bachelor of Science, 1975.
4. Djokomoeljanto, R: Iodine deficiency in Indonesia: reassessment of its control program. In: Tanphaichitr V, Dahlan V, Suphakarn V, Valyasevi (eds). Human Nutrition-Better Nutrition Better Life. Arkonsmai Press Bangkok 318-23, 1984.
5. Hetzel BS, Hales IB: New Zealand, Australia and New Guinea. In: Stanbury JB, Hetzel BS (eds.) Endemic Goiter and Endemic Cretinism. Wiley Medical, New York, 1980.
6. Boyages SC, Maberly GF, Eastman CJ, Halpern, Collin, J, Jupp TC, ZH Wong, dan Jin C: Features of endemic cretinism in Qinghai, China. The Thyroid. Elsevier Science Publishers B. V. (Biomedical division) 311-410; 1988.
7. Hetzel BS: Iodine deficiency disorder (IDD) and their eradication. The Lancet: 1126, 1983.
8. Chen PC: The prevalence of endemic goiter among Penans of the Baram. Med J Malaysia 43(2): 159-61, 1988.
9. Polunin IV: Goiter control; West and East Malaysia. Internal WHO report. Malaysia, 1971.
10. Osman Ali, Khalid BAK, Tan TT, Sakinah O, Wu LL and Ng ML: The prevalence of diabetes mellitus among Malaysian aborigines and Malays and its relationship with socioeconomic, health and nutrition factors. Diabetes Care 16(1): 68-75, 1993.
11. Delange F, Hershman JM, Ermans AM: Relationship between the serum thyrotropin level, the prevalence of goiter and the pattern of iodine metabolism in Idjwi Island. J Clin Endocrinol. 33: 261-268, 1971.
12. Belling GB: Further studies on the recovery of iodine as iodine-125 after alkaline ashing prior to assay. Analyst 108: 763-765, 1983.
13. Ogihara T, Oki K, Iida Y, Hayashi S: Endemic goiter in Sarawak, Borneo Island: prevalence and pathogenesis. Endocrinol Jap 19(3): 285, 1972.
14. Bachtarzi H, Benmiloud M: TSH and goitrogen in severe iodine deficiency. Acta Endocrinol 103: 21-27, 1983.
15. Maberly GF: The aetiology, treatment and prevention of endemic goiter in Sarawak, Malaysia. A thesis submitted for the degree of Doctor of Medicine, The University of New South Wales, 1982.
16. Suwanik R, Nondasuta A, Nondusuta A: Endemic goiter in Thailand. J Med Assoc Thailand 60: 79, 1977.
17. Adam DD, Kennedy TH, Choufoer JC, Querido A: Endemic goiter in Western New Guinea III. Thyroid stimulating activity of serum from severely iodine-deficient people. J Clin

- Endocrinol 28: 685-692, 1988.
18. Pisarev MA, Utiger RD, Salvaneschi JP, Altschuler N, De Groot LJ: Serum TSH and thyroxine in goitrous subjects in Argentina. *J Clin Endocrinol* 30: 680-681: 1970.
  19. Wollman SH, Breitman TR: Changes in DNA and weight of thyroid glands during hyperplasia and involution. *Endocrinology* 86: 322-327, 1970.
  20. Delange F, Camus M, Erman AM: Circulating thyroid hormones in endemic goiter. *J Clin Endocrinol* 34: 891-895, 1972.
  21. Greer MA, Grimm Y, Studer H: Qualitative changes in the secretion of thyroid hormones induced by iodine deficiency. *Endocrinology* 83: 1193-1198.
  22. Vagenakis AG, Koutras DA, Burger A, Malamos B, Ingbar SH, Braverman LE: Studies of serum triiodothyronine, thyroxine and thyrotropine concentrations in endemic goiter in Greece. *J Clin Endocrinol* 37: 485-488, 1973.
  23. Becker DJ: The endocrine responses to protein calorie malnutrition. *Clin Endocrinol* 3: 187-222, 1983.