

HEART RATE: A PREDICTOR OF EARLY MORTALITY IN PATIENTS WITH MYOCARDIAL INFARCTION

NIZAL SARRAF-ZADEGAN, M.D.

From the Cardiovascular Research Center, Isfahan University of Medical Sciences, Isfahan, Islamic Republic of Iran.

ABSTRACT

A number of epidemiologic studies have reported a positive relationship between heart rate, cardiovascular disease and mortality. To examine the correlation between heart rate and mortality after acute myocardial infarction (AMI), 2147 patients hospitalized in coronary care units in Isfahan were investigated in a cross-sectional study. Their heart rate was measured according to an electrocardiograph obtained during the 2nd, 5th and final days of hospitalization, and all patients were followed for cardiovascular mortality. The means of 3 measurements were divided into four fractions.

A strong positive relationship between heart rate and cardiovascular mortality was obtained. Confounding variables such as age, sex, type of infarction and drugs were adjusted by regression models. The nonlinear regression relationship was due primarily to a sharp increase in mortality in the highest quartile of heart rate.

We conclude that although sinus tachycardia can be considered as a predictive independent risk factor for mortality after AMI, further longitudinal studies are required.

Keywords: heart rate, myocardial infarction, mortality, prognosis, sinus tachycardia, cardiovascular diseases, risk factors.

MJIRI, Vol. 11, No. 4, 295-298, 1998.

INTRODUCTION

It is believed that a low heart rate is associated with longevity, largely because of lower rates of cardiovascular mortality in patients with low heart rates. Increased heart rate is associated with increased risk of cardiovascular mortality.¹⁻⁶

There is also evidence of a positive relation between resting heart rate and non-cardiovascular mortality,^{2,7,8} with a significant association between heart rate and cancer mortality.^{8,9} New data from the Framingham study showed

that this association is seen in normotensive and hypertensive patients, and also indicate an association between heart rate and sudden cardiac death.^{4,10}

Sinus tachycardia following AMI (acute myocardial infarction) is common and is frequently an unfavorable prognostic sign. Patients with a large area of infarcted myocardium may have sinus tachycardia due to left ventricular dysfunction; other causes are anxiety, fever, pain, pulmonary embolism, hypovolemia, anemia or drugs such as dobutamine, dopamine, etc.¹⁶ Commonly, the initial episode of tachyarrhythmia and sinus tachycardia begin on the 1st day of AMI.¹⁶

MATERIALS AND METHODS

This study was a cross-sectional one with 2147 patients

* Address correspondence and reprint requests to: N. Sarraf-zadegan, M.D., Associate Professor of Cardiology, Cardiovascular Research Center, Isfahan University of Medical Sciences, P.O. Box: 81465-1148, Isfahan, Islamic Republic of Iran.

Heart Rate and Mortality in MI Patients

aged 35-70 years hospitalized in all coronary care units in Isfahan between 1993-1994. These patients suffered from acute myocardial infarction (AMI) which was diagnosed according to WHO criteria. Patients suffering from complicated AMI such as those with arrhythmia, heart failure, cardiogenic shock, pulmonary edema, pericarditis and persistent chest pain were excluded according to physical examination and chest x-ray. Each completed a questionnaire for cardiovascular disease risk factors and, to determine the heart rate, an electrocardiogram was recorded at paper speed of 2.5 cm/s on the 2nd, 5th and final days of hospitalization.

All patients were followed for cardiovascular deaths during their hospitalization period. Cardiovascular death events include sudden coronary death, stroke death and death from arrhythmia or cardiac failure.

All data were coded, checked and entered using the software package Epi 6 data sets at the Computer Unit of the Cardiovascular Research Center in Isfahan. Calculations were done on a personal computer using the statistical software package SPSS. The effect of some variables such as age, sex, infarction type or drugs used were adjusted. All patients were distributed into four groups according to their heart rate (mean of three readings) and mortality rate was determined. The mortality rate was computed for each approximate quartile.

RESULTS

A total of 2147 patients aged 35-70 years suffering from acute myocardial infarction (AMI), hospitalized in coronary care units in 1993-1994 were enrolled and followed for cardiovascular death during their hospitalization period. Patients suffering from complicated AMI were excluded. The association between heart rate (which was the mean of 3 measurements on different days) and mortality was examined (Table I). For the cross classification analysis, heart rate was divided into approximate quartiles, then mortality rates were determined for each quartile. The results indicated that the mortality rates were significantly different for the four heart rate groups, and the relationship was strongly positive. Fig. 1 indicates that there is a non-linear regression relationship between heart rate and cardiovascular death. This non-linear association may be due primarily to the sharp increase in cardiovascular death in the highest quartile of heart rate.

DISCUSSION

Some studies suggest that high heart rate may be an independent risk factor for sudden death from coronary heart disease (CHD) and death from causes other than

Table I. The mortality rate according to heart rate quartiles.

Heart rate group * (bpm)	No. of patients	No. expired	Mortality (%)
41-60	132	2	1.5
61-80	960	31	3.2
81-100	982	45	4.6
101-120	73	8	10.9

* Mean of three measurements on the 2nd, 5th and final day of hospitalization.




Fig. 1. Regression relationship between heart rate and mortality rate of 2147 CCU patients.

cardiovascular disease.²⁻⁶ In contrast to these positive findings, three large studies found no significant association between heart rate and incidence of CHD.¹¹⁻¹³ Some reported that cardiovascular-related mortality rate is higher in men than in women irrespective of heart rate,² while others consider sinus tachycardia to be a predictor of overall mortality in women only.²⁸

The heart rate in patients with acute myocardial infarction (AMI) may vary from marked bradycardia to rapid regular or irregular tachycardia, depending on the underlying rhythm and the degree of left ventricular failure.¹⁴

In this study, as mentioned before, although patients with heart failure, pericarditis, arrhythmia, and persistent chest pain were excluded, mortality still increased as heart rates increased. Therefore it seems that sinus tachycardia can be an independent predictor of mortality in patients with AMI.

Initially after AMI the pulse is rapid and regular (sinus tachycardia at 100 to 110 beats/min), but as the patient's pain and anxiety are relieved, the pulse rate tends to decrease.^{14,15} In order to relieve this initial effect, we measured the heart rate on the 2nd, 5th and last day before discharge, and mean values were categorized in one of 4 quartiles.

The presence of an increased heart rate following AMI is common and considered as an unfavorable prognostic sign because increased heart rate enhances myocardial oxygen demand, while the decrease in diastole decreases

diastolic coronary flow.¹⁶ There was no significant difference in the frequency of atrial tachyarrhythmia in relation to the area of AMI.¹⁶ The obtained results from this study were similar, because the type of infarction was adjusted initially.

Several etiologies for atrial tachyarrhythmias after AMI have been proposed, including congestive heart failure and cardiogenic shock, infarct size, compromise of the sinoatrial node blood supply and pericarditis.^{17-20, 32} These causes were excluded in our study and non-complicated patients were enrolled. Another explanation for sinus tachycardia after AMI which was not considered in our study was acute systolic hypertension (ASH). Some studies have reported an increased incidence of acute left ventricular failure and of arrhythmias such as sinus and ventricular tachycardia in patients with ASH in comparison with those with normal blood pressures.³⁸

In the coronary care unit, some forms of supraventricular tachyarrhythmias (i.e., sinus tachycardia) are exhibited by nearly two-thirds of patients.^{16, 23-24} On hospital discharge 2 or 3 weeks after the acute event, 30 to 50% of patients still show some type of supraventricular tachyarrhythmia.^{25, 26} Supraventricular tachycardia usually develops in the presence of severe hemodynamic decompensation and is associated with an increased risk of subsequent cardiac death.^{16, 24, 27} The nonlinear regression relationship obtained in our study indicates that when heart rate was between 101-120 bpm, the mortality rate increases to 10.9%

Some reports have shown that sinus tachycardia and some types of arrhythmia are the most common clinical and electrocardiographic prodromes of ventricular fibrillation.^{29, 34, 36} Unfortunately, this was not investigated in our study.

Sinus tachycardia has been considered as a risk factor for mortality after using thrombolytic therapy in patients with AMI,³⁰ and for ventricular tachycardia with higher rates (181 to 220 bpm). Some investigators consider sinus tachycardia as a complication of AMI, just as ventricular tachycardia or fibrillation, pulmonary edema, cardiogenic shock and advanced heart block are others.³³ Previous findings from longitudinal studies suggested that persistent sinus tachycardia occurring during the first four hospital days in patients with AMI can predict subsequent complications requiring urgent medical attention,³⁵ but we could find no similar effect due to the type of our study which was cross-sectional and without patient follow-up.

Sinus tachycardia is considered as one of the later arrhythmias (occurring after the 15th day of AMI) which is controlled by beta-adrenergic blocking agents as well as amiodarone and verapamil.³⁷ One explanation for the association between heart rate and mortality is the effect of major risk factors.¹ Heart rate was positively correlated with systolic and diastolic blood pressure³⁹⁻⁴³ and cigarette use,⁴⁴⁻⁴⁷ and showed a small or inconsistent association with age, serum cholesterol and relative weight.^{13, 40-48} Unfortunately

this subject was not investigated in our study.

We conclude that although a strong positive relationship between heart rate and mortality after AMI was obtained from this study, further longitudinal studies with patient follow-up are required in order to improve the predictive ability of sinus tachycardia as an independent predictor of mortality after AMI.

ACKNOWLEDGEMENT

We wish to express our appreciation to Miss Nasrin Jamadi and Miss Noshin Mohammadifard for their assistance in the preparation of this manuscript.

REFERENCES

1. Dyer AR, Persky V, Stamper J, et al: Heart rate as a prognostic factor for coronary heart disease and mortality: findings in three Chicago epidemiologic studies. *Am J Epidemiol* 112: 749-63, 1980.
2. Kannel WB, Kannel C, Paffen Barger RS, et al: Heart rate and cardiovascular mortality: the Framingham study. *Am Heart J* 113: 1489-94, 1987.
3. Schroll M, Hangerup LM: Risk factors of myocardial infarction and death in men aged 50 at entry: a ten-year prospective study from the Glostrup population studies. *Dan Med Bull* 24: 252-5, 1977.
4. Friedman GD, Klatsky AL, Siegel AB: Predictors of sudden cardiac death. *Circulation* 51 (suppl 3): 164-9, 1975.
5. Medalie JH, Kahn HA, Neufeld HN, et al: Five year myocardial infarction incidence. II. Association of single variables to age and birth place. *J Chronic Dis* 26: 329-349, 1973.
6. Shurtleff D: Some characteristics related to the incidence of cardiovascular disease and death: Framingham study, 18 year follow-up. Section 30. In: Kannel WB, Gordon T (eds.), *The Framingham study—an epidemiological investigation of cardiovascular disease*. Washington DC, DHEW, 1974.
7. Wilhelmsen L, Berglund G, Elmfeldt D, et al: The multifactor primary prevention trial in Goteborg, Sweden. *Eur Heart J* 7: 279-88, 1986.
8. Persky V, Dyer AR, Leonas J, et al: Heart rate: a risk factor for cancer? *Am J Epidemiol* 114: 477-87, 1981.
9. Wamamethee G, Shaper G, MacFarlane P: Heart rate, physical activity, and mortality from cancer and other non-cardiovascular diseases. *Am J Epidemiol* 134: 735-748, 1993.
10. Pepine CJ: Heart rate and mortality. *J Myocardial Ischemia* 4(6): 8-13, 1992.
11. Tibblin G, Wilhelmsen L, et al: Risk factors for myocardial infarction and death due to ischemic heart disease and other causes. *A J Cardiol* 35: 514-523, 1975.
12. Paffen Barger RS, Wolf PA, Notkin J, et al: Chronic disease in former college students. I. Early precursors of fatal coronary

Heart Rate and Mortality in MI Patients

- heart disease. *Am J Epidemiol* 83: 314-328, 1966.
13. Keys A, Taylor HL, Blackburn H, et al: Mortality and coronary heart disease among men studied for 23 years. *Arch Intern Med* 128: 201-214, 1971.
 14. Paspernak R, Braunwald E, Sobel B: Acute myocardial infarction. In: Braunwald E (ed.), *Heart Disease, A Textbook of Cardiovascular Medicine*. Philadelphia: W.B. Saunders Company, Vol 2, p. 1216, 1992.
 15. Webb SW, Adgey AA, Pantridge JF: Autonomic disturbance at onset of acute myocardial infarction. *Br Med J* 818: 89, 1982.
 16. Libberthson RR, Salisbury KW, Hutter AM, DeSanctis RW: Atrial tachyarrhythmias in acute myocardial infarction. *Am J Med* 60: 956-960, 1976.
 17. Lown B, Klein MD, Hershberg PI: Coronary and precoronary care. *Am J Med* 46: 705, 1969.
 18. Lown B, Vasaux C, Hood WB Jr: Unresolved problems in coronary care. *Am J Cardiol* 20: 494, 1967.
 19. James TN: The coronary circulation and conduction system in acute myocardial infarction. *Progr Cardiovascular Dis* 10: 410, 1968.
 20. James TN: Pericarditis and the sinus node. *Arch Intern Med* 110: 305, 1962.
 21. James TN: Arrhythmia associated with acute myocardial infarction. *Heart Bull* 15: 87, 1966.
 22. James TN: Myocardial infarction and atrial arrhythmias. *Circulation* 24: 761, 1961.
 23. DeSanctis RW, Block P, Hutter A: Tachyarrhythmia in myocardial infarction. *Circulation* 45: 681-702, 1972.
 24. Cristall N, Szwarcberg J, Gueron M: Supraventricular arrhythmias in acute myocardial infarction. Prognostic importance of clinical setting: mechanism of production. *Ann Intern Med* 82: 35-39, 1975.
 25. Moss A, Schnitzler R, Greem R: Ventricular arrhythmias three weeks after acute myocardial infarction. *Ann Intern Med* 75: 837-841, 1971.
 26. Zoni Berisso M, Ferroni A, De Caro E: Clinical significance of supraventricular tachyarrhythmias after acute myocardial infarction. *Eur Heart J* 7: 743-748, 1986.
 27. Luria MH, Keake JD: Acute myocardial infarction: prognosis after recoronary. *Ann Intern Med* 85: 561-565, 1976.
 28. Casiglia E, Spolaore P, Ginocchio G: Mortality in relation to Minnesota code interns in elderly subjects. Sex-related differences in a cardiovascular study in the elderly. *Jpn Heart J* 34(5): 567-77, 1993.
 29. Denisiuk VI, Brovko EA, Lipnitskii TN: The clinico-electrocardiographic prodromes of fatal cardiac arrhythmias in patients with an acute myocardial infarction. *Kardiologia* 33(10): 43-5, 1993.
 30. Hillis LD, Forman S, Braunwald E: Risk stratification before thrombolytic therapy in patients with acute myocardial infarction. *J Am Coll Cardiol* 16(2): 313-5, 1990.
 31. Bluzhas J, Lukshien D: Ventricular tachycardia in myocardial infarction: relation to heart rate and premature ventricular contractions. *Eur Heart J* 6(9): 745-50, 1985.
 32. Grand P, Pedersen A: Myocardial infarction size: correlation with cardiac arrhythmias and sudden death. *Eur Heart J* 5(8): 622-7, 1984.
 33. Pryor DB, Hindman MC, Wagner GS: Early discharge after myocardial infarction. *Ann Intern Med* 99 (4): 528-38, 1983.
 34. Olsson G, Rehnqvist N: Sudden death precipitated by psychological stress. A case report. *Acta Med Scand* 212(6): 437-41, 1982.
 35. Severance HW Jr, Morris KG, Wagner GS: Criteria for early discharge after acute myocardial infarction: validation in a community hospital. *Arch Intern Med* 149: 39-41, 1982.
 36. Brignole M, Barra M, Sartore B, Prato R: Sudden death during Holter monitoring in later phase of hospitalization after myocardial infarction. *G Ital Cardiol* 12(7): 536-8, 1982.
 37. Zhivoderov VM, Doshchitsin VL, Dunaeva ZI: Late arrhythmias in myocardial infarction. *Kardiologia* 20(1): 22-5, 1980.
 38. Buyukozturk K, Gultekin N, Deligonul U: Acute systolic hypertension after acute myocardial infarction: prognostic and therapeutic significance. *Acta Cardiol* 35(5): 341-7, 1986.
 39. Stamler J, Rhomberg P, Schoenberger JA: Multivariate analysis of the relationship of seven variables to blood pressure: findings from the Chicago Heart Association Detection Project in industry, 1967-1972. *J Chronic Dis* 28: 527-548, 1975.
 40. Stamler J, Stamler R, Rhomberg P, et al: Multivariate analysis of the relationship of six variables to blood pressure: findings from Chicago community surveys, 1965-1971. *J Chronic Dis* 28: 499-525, 1975.
 41. Paffenbarger RS, Thome MC, Wing AL: Chronic disease in former college students. VIII. Characteristics in youth predisposing to hypertension in later years. *Am J Epidemiol* 88: 25-32, 1968.
 42. Kannel WB, Sorlie P: Hypertension in Framingham. In: Paul O (ed), *Epidemiology and Control of Hypertension*. Miami, FL, Symposium Specialists, pp. 553-590, 1975.
 43. Kahn HA, Medalie JH, Neufeld HN, et al: The incidence of hypertension and associated factors: the Israeli ischemic heart disease study. *Am Heart J* 84: 171-182, 1972.
 44. Blackburn H, Brozek J, Taylor HL, et al: Comparison of cardiovascular and related characteristics in habitual smokers and non-smokers. *Ann NY Acad Sci* 90: 277-289, 1960.
 45. Higgins MW, Kjelsberg M: Characteristics of smokers and non-smokers in Michigan. II. The distribution of selected physical measurements and physiologic variables and the prevalence of diseases in smokers and non-smokers. *Am J Epidemiol* 86: 60-77, 1967.
 46. Thomas CB: Characteristics of smokers compared with non-smokers in a population of healthy young adults, including observations of family history, blood pressure, heart rate, body weight, cholesterol and certain physiologic traits. *Ann Intern Med* 53: 697-718, 1960.
 47. Goldbourt U, Medalie JH: Characteristics of smokers, non-smokers and ex-smokers among 10,000 adult males in Israel. II. Physiologic, biochemical and genetic characteristics. *Am J Epidemiol* 105: 75-86, 1977.
 48. Goldbourt U, Medalie JH, Neufeld HN: Clinical myocardial infarction over a five-year period. III A multivariate analysis of incidence, the Israeli ischemic heart disease study. *J Chronic Dis* 28: 217-237, 1975.