# **Determinants of maternal mortality in Eastern Mediterranean** region: A panel data analysis

Mohsen Bayati<sup>1</sup>, Sajad Vahedi<sup>2</sup>, Firooz Esmaeilzadeh<sup>3</sup>, Zahra Kavosi<sup>4</sup> Zahra Jamali<sup>5</sup>, Abdolhalim Rajabi<sup>6</sup>, Yousef Alimohamadi<sup>\*7</sup>

Received: 29 May 2015

Accepted: 1 December 2015

Published: 23 April 2016

#### Abstract

Background: As one of the main criteria of health outcomes, maternal mortality indicates the socioeconomic development level of countries. The present study aimed at identifying and analyzing the effective factors on maternal mortality in Eastern Mediterranean Region (EMR) of the World Health Organization (WHO).

Methods: Analytical model was developed based on the literature review. Panel data of 2004-2011 periods for 22 EMR countries was used. Required data were collected from WHO online database. Based on results of diagnostic tests for panel data model, parameters of model were estimated by fixed effects method.

**Results**: Descriptive statistics demonstrated the large disparities in social, economic, and health indicators among EMRO countries. Findings obtained from evaluating the model showed a negative, significant relationship between GDP per capita ( $\beta$ =-0.869, p<0.01), health expenditure)  $\beta$ =-0.525, p<0.01 (female literacy rate)  $\beta$ =-1.045, <0.01 (skilled birth attendance)  $\beta$ =-0.899, p<0.05) and maternal mortality rate.

Conclusion: Improved income and economic development, increased resources allocated to the health sector, improved delivery services particularly the increased use of trained staff in the delivery, improve quality of primary care centers, mitigating the risks of marginalization and its dangers, and especially improving the level of women's education and knowledge are the key factors in policy making related to maternal health promotion.

Keywords: Maternal mortality rate, Maternal health, Panel data model, EMRO.

Cite this article as: Bayati M, Vahedi S, Esmaeilzadeh F, Kavosi Z, Jamali Z, RajabiA, Alimohamadi Y. Determinants of maternal mortality in Eastern Mediterranean region: A panel data analysis. Med J Islam Repub Iran 2016 (23 April). Vol. 30:360.

### Introduction

Maternal mortality rate is a very important indicator, reflecting a country's economic, cultural or health system development, also recognized worldwide (1). Perhaps one of the sensitive indicators of inequality in health is maternal and child health. This indicator also shows the quality and effectiveness of health policies (2). Maternal mortality includes the deaths during pregnancies or the ones 42 days after deliveries. Traditionally, this indicator includes direct maternal deaths from pregnancy, indirect maternal deaths from pregnancy, and maternal deaths that occur during pregnancy, but do not relate to it (3,4). Hundreds of thousands of women in lowincome countries die each year from complications of pregnancy, abortion and childbirth. It is estimated that every minute

<sup>1</sup>. PhD Candidate of Health Economics, Department of Health Economics & Management, School of Public Health, Tehran University of Medical Sciences, & Health Management and Human Resources Research Center, Faculty of Management and Medical Information, Shiraz University of Medical Sciences, Shiraz, Iran. bayatim66@gmail.com

<sup>&</sup>lt;sup>2</sup>. PhD Student of Health Economics, Health Management and Economics Research Center, Iran University of Medical Sciences, Tehran, Iran. vahedi.s3@gmail.com

<sup>&</sup>lt;sup>3</sup>. PhD Candidate of Health Economics, Department of Health Economics & Management, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran. firooz.esmaeilzadeh@gmail.com

Associate Professor, Department of Health Services Management, Faculty of Management and Medical Information, Shiraz University of Medical Sciences, Shiraz, Iran. zhr.kavosi@gmail.com <sup>5</sup>. BSc, Department of Environmental Health, School of Public Health, Shiraz University of Medical Sciences, Shiraz, Iran.

zjamali923@gmail.com PhD Student of Epidemiology, Department of Epidemiology, School of Public Health, Iran University of Medical Sciences, Tehran, Iran. rajabiepid@gmail.com

<sup>. (</sup>Corresponding author) PhD student of Epidemiology, Department of Epidemiology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran. y.alimohamadi67@gmail.com

a woman and every day 16,000 women in the world die from pregnancy complications (5-7). According to other statistics, the maternal mortality rate is about 402 per 100000 live births which is approximately half million deaths a year (8). One of the Millennium Development Goals is maternal health improvement (MDG5). It aims at reducing the maternal mortality rate by 75% between 1990 and 2015 and the universal access to reproductive health by 2015 (9).

It has been globally reported that maternal mortality between 1990 and 2008 has reduced by 1.03% on average, which is lower than the 5.5% recommended by MDG5 (10). Failure to reduce the maternal mortality over the last 20 years has been one of the world's health problems (11). Hence, women's health is one of the most important priorities of the world health assemblies, and identifying the main factors of women's mortality as well as improving it depends on a correct definition of required priorities for appropriate prevention, diagnosis and treatment (12). Analyzing the factors affecting maternal mortality in ecological and macro levels may provide some credible information and evidence to health sector manager in order to make appropriate policies to improve maternal health. A few general studies on nonclinical determinants of maternal mortality have analyzed different factors including access to safe drinking water, government corruption, out-of-pocket payment, fertility rate, education level, delivery done by skill birth attendants, GDP per capita, health expenditures, life expectancy at birth, access to health services, and access to food (13-17). In the present study, the economic, social and healthcare system-related factors affecting maternal mortality in the EMR of WHO will be evaluated.

# Methods

### *Study setting*

The present study was an ecological cross-sectional and retrospective one that has used the panel data of the 22 countries

located in the Eastern Mediterranean Region (EMR) of the WHO. No sampling was done in this study. The study population was the countries located in the EMR and with regard to the availability of the intended data, such countries as Afghanistan, Bahrain, Djibouti, Egypt, Islamic Republic of Iran, Iraq, Jordan, Kuwait, Lebanon, Libya, Palestine, Oman, Pakistan, Qatar, Saudi Arabia, Somalia, Sudan, Syria, Tunisia, United Arabic Emirates, Yemen, and Morocco were studied. The research period was from 2004 to 2011.

# Data and Variables

The data about the intended variables was collected from the WHO website where the information about the EMR was provided. The variables studied in this research were as follows:

Maternal mortality tare (the number of maternal deaths per 100000 live births); Gross domestic product per capita (in international dollar); primary health care center ratio; the total fertility rate; health expenditure share of gross domestic product (total health expenditure to GDP); urbanization (the proportion of urban population to total population); women's education (the 15-year-old and older women's literacy rate); and percentage of births attended by skilled health personnel.

# Econometrics analysis

The effects of economic, social and health system factors on maternal mortality rate can be studied using the model below.

The initial model is: MMR=f(S, E, H)

Where MMR, S, E, and H are maternal mortality, social factors, economic factors, and health system-related factors, respectively. The model developed as follows using previous studies (e.g. 7,14,16) and accessing reliable data.

In the present study, the social factors included total fertility rate (TFR), urbanization rate (URB), and female's education level (EDU). The only economic determinant was the gross domestic product (GDP) per capita. Finally, the factors related to the health system were health expenditure share of income (HE), percentage of births attended by skilled health personnel (BA), and percentage of primary healthcare centers (PHC). Therefore, the final econometric model presented as follows:

# $\label{eq:limit} \begin{array}{ll} LMMRit=& \beta_0+\beta_1+LGDPit+\beta_2LHEit+\beta_3LE\\ DUit+\beta_4LTFRit+\beta_5LBAit+\beta_6LPHCit+\beta_7LU\\ RBit+Uit \end{array}$

The variables are entered in the model as natural logarithms. Therefore, the coefficients  $\beta_1$  to  $\beta_7$  indicate the maternal mortality rate elasticity with respect to explanatory variables. U<sub>it</sub> shows the error term with classic assumptions. To determine the method of estimation the parameters of model, diagnostic tests such as the Chow and Hausman tests were applied. Eviews5 software was also used for data analysis.

# Results

Table1 shows the descriptive findings including the mean, median, maximum, minimum, and standard deviation of the studied variables for EMR from 2004 to 2011. On average, in the EMR 181.6 MMR has occurred per 100,000 live births. Other findings on maternal mortality indicate the disparity of this variable between the countries of the region so that the minimum and maximum rates have been between 0 and 1600. The EDU was almost low in this region and it was also remarkable disparate distribution. During pregnancies, on average woman in this region gives birth to more than 3 children.

The HE has been approximately 5% for the region. Generally, descriptive findings show a significant disparity or inequality in economic, social, health systems-related, and maternal mortality variables. To select the best estimation method for the model, Chow and Hausman diagnostic tests were applied (Table 2). According to the results from Chow Test (F and Chi-Sq statistics), the null hypothesis based on using Pool effect method is not accepted at 90% confidence level. According to the results from Hausman Test Chi-Sq statistics, the null hypothesis based on using Random Effects (RE) method is not accepted at 99% confidence level. Consequently, the Fixed Effects (FE) method is selected to estimate the model of this study. Results of model estimation through the FE method are shown in Table 3. Since the independent variables and the dependent variable in the model are logarithmic, the estimated coefficients show the dependent variable (MMR) elasticity with respect to the explanatory variables. For example, the GDP coefficient (0.869) shows the MMR elasticity with respect to the GDP; this means that 1% increase in the MMR would lead to 0.86% MMR reduction. Coefficients of other variables can be interpreted the same way.

GDP, HE, EDU, and BA have had a negative, significant relationship with MMR in this region. Variables such as URB and PHC showed a positive, significant relationship with MMR, while the TFR did not have a significant effect on MMR.

Overall, significance of the model based on the F statistic (p<0.001) can be confirmed. In addition, the estimated value of the adjusted R<sup>2</sup> (0.795) shows that almost 80% of the changes in maternal deaths can

Table 1. Descriptive statistics of the studied variables in the Eastern Mediterranean countries (2004-2011)								
	Mean	SD	Min	Max	Median			
Maternal mortality ratio per 100 000 live	181.60	337.95	0	1600	40			
Gross Domestic Product Per Capita (GDP)	10702.7	17444.11	158	88990	2720.41			
Health Expenditure as % of GDP (HE)	5.01	2.71	1.5	16	4.2			
Literacy rate among Female adults aged 15+	67.34	22.05	5	95.6	74			
Total Fertility Rate (TFR)	3.44	1.34	0.9	6.3	3.1			
BA (% Births attended by skilled health) per-	82.90	23.88	14	100	95.5			
Primary Healthcare units and Centers (PHC)	1.65	1.27	0.2	8	1.6			
Urbanization Rate (URB)	67.27	23.23	22	100	71			

Table 2. Results of diagnostic tests for model estimation method					
Test	Statistic	р			
Chow test	$\chi^2 = 15.204$	0.033			
	F=2.032	0.057			
Hausman test	$\chi^2 = 29.380$	<0.001			

Table 3. Results of model estimation with fixed effects method for the Eastern Mediterranean countries (2004-2011)

Variable	Coefficients (Elasticity)	t-statistic	р		
С	14.267	12.898	< 0.001		
LGDP	-0.869	-7.626	< 0.001		
LHE	-0.525	-2.957	0.003		
LEDU	-1.045	-2.894	0.004		
LTFR	0.394	1.618	0.108		
LBA	-0.899	-2.002	0.047		
LPHC	0.197	2.014	0.046		
LURB	1.282	2.867	0.005		
Goodness of fit	$R^2 = 0.818$				
	Adjusted R <sup>2</sup> =0.795				
Overall significance of model	F-statistic=35.361				
-	p≤0.001				

be explained by the economic, social and health system variables used in this study.

### Discussion

In the present study, it was seen that the GDP, HE, EDU, BA have had a negative, significant relationship with MMR in the EMR. That is to say, the increase of each abovementioned indicators would lead to maternal mortality reduction. This emphasizes on the importance of government investments in the health sector and the share of the health sector of the total government expenditures so that by increasing budgets for the health sector, great steps might be taken to improve the health of population especially mothers. Moreover, improving women's literacy and knowledge can significantly contribute to their health. Findings of a similar study by Luis Alvarez in Africa were similar to the present study so that there was a negative, significant relationship between maternal mortality rate and the GDP per capita, health expenditure, women literacy rate, and the number of deliveries by skilled birth attendance in this region (13). Another study concluded in Sub-Saharan African countries, birth in the presence of health professionals and the life expectancy at birth are highly correlated to maternal mortality. Furthermore, a very effective relationship was found between GDP per capita and health expenditure per capita and maternal mortality (14). In other study in Pakistan villages, it was seen that mothers with a previous history of abortion were among those at high risk for mortality. Besides, a significant relationship was observed between having access to health serprofessional health staff vices. and healthcare during pregnancy and MMR so that maternal mortality reduced considerably as these situations improved (15). The positive impacts of GDP, HE, EDU on health outcome also, were reported in the other studies in EMR (18,19).

The present study suggested that variables such as URB and the PHC have a positive, significant relationship with MMR. The relationship between URB and MMR could be interpreted as follows: since the majority of the EMR are developing countries, migration from rural to urban areas and creating the marginalization phenomenon is one serious problem in these countries. Theredevelopment of marginalization fore. around big cities has caused high population areas that lack the minimum level of sanitary and welfare facilities. There were different findings from other studies; the positive (20) and also negative impact of urbanization on health outcomes reported in previous research (21).

The imbalance between population and resources in city fringes has caused a lot of problems, including the spread of different

Downloaded from mjiri.iums.ac.ir on 2025-07-06

diseases, increased mortality due to diseases, increased crime, and other healthrelated problems (22). Perhaps the reason for a positive relationship between the health centers ratio and MMR in this study is that this ratio is a little indicator of the primary care centers' conditions. Therefore, this positive relationship probably shows the lack of good performance and quality of these centers. In his study, Hertz evaluated the social and environmental factors affecting life expectancy as well as maternal and child mortality. Results showed a strong correlation between the factors related to food, available medical supplies, sanitation facilities and safe drinking water resources and maternal mortality so that improving them would significantly reduce maternal mortality (17).

In this study, the TFR had no significant effect on MMR. However, Muldoon (2009) found that fertility rates higher than 1 would bring about significant risk of maternal death (16).

As stated in descriptive findings, there exists considerable disparity between different countries of the region in terms of the studied variables. Therefore, one reason for the lack of compliance of some results such as the positive impact of urbanization or primary care centers on maternal mortality, with that of other studies is the heterogeneity of the data. Furthermore, the interpretation of the variable's coefficients has to be done with caution; results obtained from the entire EMR are considered as a unit of analysis and should not be extended to individual countries within the region.

# Conclusion

Mothers belong to the vulnerable groups of societies are seriously influenced by adverse conditions, especially in relation to health sector deficit. However, the maternal health is one of the most important indicators of economic, social and health development in countries. Improved earnings and economic development in the region, increasing the resources allocated to the health sector, improving delivery service particularly the increased use of trained staff in deliveries, enhancing the quality of primary health care, reducing marginalization phenomenon and other urban hazards, and especially, improving the quantity and quality of women's knowledge and literacy are the most important factors in promoting women's health and reducing maternal mortality in the EMR.

# Acknowledgements

This research resulted from a research project approved by the Student Research Committee of Shiraz University of Medical Sciences. Hereby, they are sincerely thanked.

### References

1. Liang J, Zhu J, Dai L, Li X, Li M, Wang Y. Maternal mortality in China, 1996-2005. Obstet Gynecol Int J 2010;110(2):93-6.

2. Marin GH, Silberman M, Uriarte A, Sarijulis M, Ozaeta B, Henen J. Healthcare model based on personalised attention: impact on maternal mortality and health system quality. Qual Prim Care 2011; 19(5):311-6.

3. Khan KS, Wojdyla D, Say L, Gulmezoglu AM, Van Look PF. WHO analysis of causes of maternal death: a systematic review. Lancet 2006;367 (9516):1066-74.

4. Campbell OM, Graham WJ. Strategies for reducing maternal mortality: getting on with what works. Lancet 2006;368(9543):1284-99.

5. McTavish S, Moore S, Harper S, Lynch J. National female literacy, individual socio-economic status, and maternal health care use in sub-Saharan Africa. Soc Sci Med (1982) 2010;71(11):1958-63.

6. McCarthy J, Maine D. A framework for analyzing the determinants of maternal mortality. Stud Fam Plan 1992;23(1):23-33.

7. Robinson JJ, Wharrad H. The relationship between attendance at birth and maternal mortality rates: an exploration of United Nations' data sets including the ratios of physicians and nurses to population, GNP per capita and female literacy. JAN 2001;34(4):445-55.

8. Almerie MQ, Matar HE, Almerie Y. A 20-year (1989-2008) audit of maternal mortality in Damascus, Syria. Obstet Gynecol Int J 2011; 112(1):70-1.

9. Hogan MC, Foreman KJ, Naghavi M, Ahn SY, Wang M, Makela SM, et al. Maternal mortality for 181 countries, 1980-2008: a systematic analysis of progress towards Millennium Development Goal 5. Lancet 2010;375(9726):1609-23.

Med J Islam Repub Iran 2016 (?23 April). Vol. 30:360.

10. Amarin Z, Khader Y, Okour A, Jaddou H, Al-Qutob R. National maternal mortality ratio for Jordan, 2007-2008. Obstet Gynecol Int J 2010; 111(2):152-6.

11. Horton R. Maternal mortality: surprise, hope, and urgent action. Lancet 2010;375(9726):1581-2.

12. Ribeiro PS, Jacobsen KH, Mathers CD, Garcia-Moreno C. Priorities for women's health from the Global Burden of Disease study. Obstet Gynecol Int J 2008;102(1):82-90

13. Alvarez JL, Gil R, Hernandez V, Gil A. Factors associated with maternal mortality in Sub-Saharan Africa: an ecological study. BMC public health 2009;9:462.

14. Buor D, Bream K. An analysis of the determinants of maternal mortality in sub-Saharan Africa. J Womens Health (Larchmt) 2004;13(8): 926-38.

15. Midhet F, Becker S, Berendes HW. Contextual determinants of maternal mortality in rural Pakistan. Soc Sci Med 1998;46(12):1587-98.

16. Muldoon KA, Galway LP, Nakajima M, Kanters S, Hogg RS, Bendavid E, et al. Health system determinants of infant, child and maternal mortality: A cross-sectional study of UN member countries. Global and health 2011;7:42.

17. Hertz E, Hebert JR, Landon J. Social and environmental factors and life expectancy, infant

mortality, and maternal mortality rates: results of a cross-national comparison. Soc Sci Med 1994;39(1): 105-14.

18. Bayati M, Akbarian R, Kavosi Z. Determinants of life expectancy in eastern mediterranean region: a health production function. Int J Health Policy Manag 2013;1(1):57-61.

19. Homaie Rad E, Vahedi S, Teimourizad A, Esmaeilzadeh F, Hadian M, Torabi Pour A. Comparison of the effects of public and private health expenditures on the health status: a panel data analysis in eastern mediterranean countries. Int J Health Policy Manag 2013;1(2):163-7.

20. Bayati M, Akbarian R, Kavosi Z, Sadraei Javaheri A, Amini Rarani M, Delavari S. Socioeconomic Determinants of Health in Western Pacific Region: A Panel Data Analysis. Social welfare quarterly 2013;12(47):111-30.

21. Samadi AH, Keshtkaran A, Kavosi Z, Vahedi S. The Effect of Fiscal Decentralization on Underfive Mortality in Iran: A Panel Data Analysis. Int J Health Policy Manag 2013;1(4):301-6.

22. Esmaielnasb N, Amirkhani A, Ansari F, Bahonar A, Bolouki Z, Chaman R, et al. Epidemiology Textbook of prevalent disease in Iran. frist ed. Yavari P, editor. Tehran: Gap Publisher; 2014.