



Investigation of the relationship between end-tidal carbon dioxide and partial arterial carbon dioxide pressure in patients with respiratory distress

Gholamreza Masoumi¹, Amir Noyani², Arezoo Dehghani³, Ali Afrasiabi⁴, Nahid Kianmehr*⁵ 

Received: 11 Jan 2019

Published: 24 Jun 2020

Conflicts of Interest: None declared

Funding: Iran University of Medical Science

*This work has been published under CC BY-NC-SA 1.0 license.

Copyright© Iran University of Medical Sciences

Cite this article as: Masoumi Gh, Noyani A, Dehghani A, Afrasiabi A, Kianmehr N. Investigation of the relationship between end-tidal carbon dioxide and partial arterial carbon dioxide pressure in patients with respiratory distress. *Med J Islam Repub Iran.* 2020 (24 Jun);34:67. <https://doi.org/10.47176/mjiri.34.67>

Brief Communication

Blood gas test is requested when a person shows either signs of imbalance in oxygen/carbon dioxide or pH, such as difficulty in breathing, shortness of breath, vomiting, suffering from respiratory illness, metabolic disorder, kidney disease, and experiencing respiratory failure, or injuries that could affect breathing, including head or neck trauma. Therefore, measuring blood gas is highly important for assessing oxygenation and acid/base situation (1-3). Patients admitted to the emergency wards with respiratory distress as their main complaint need careful examination for oxygenation status, ventilation, and acid/base balance (4, 5).

Taking the arterial blood gas (ABG) test from the patient can provide valuable information for the physician. Unfortunately, ABG apparatus may not be available in all emergency wards (6-8). Arterial blood taking from the patient is time-consuming and very painful, has error probability, and needs to be repeated several times in some cases. Thus, using noninvasive methods such as pulse oximetry and capnography is necessary (9, 10). On the other hand, ABG test provides steady information on the patients' oxygenation instead of providing intermittent findings. In sum, ABG is not ideal to monitor critically-ill patients (11, 12). However, end-tidal carbon dioxide pressure can be measured using capnography (13).

End-tidal CO₂ can be a noninvasive, quick, and reliable technique which predicts PaCO₂ in patients with respiratory distress (14). Monitoring end-tidal CO₂ could be a suitable

substitute for measuring PaCO₂ in many emergency wards and operating rooms in developed countries (15). However, to date, exact correlation with PaCO₂ has not been confirmed. This study aimed to evaluate the correlation between PaCO₂ and EtCO₂ in patients with respiratory distress admitted to the emergency wards.

In this cross sectional study, the ABG test was taken and simultaneously the EtCO₂ was measured with capnograph in patients admitted to the emergency ward of Hazrat-e Rasool hospital with one or more following symptoms: difficulty breathing, grunting, tachypnea, orthopnea, costal retraction, bluish color around mouth, wheezing, and nasal flaring described as respiratory distress symptoms (16). Also, blood pressure and body temperature of the patients were recorded. Then, patients diagnosed as newborn respiratory distress syndrome were excluded. In this study, the sample volume with 30% probability was 120 cases. All patients agreed to provide their information to the researcher.

The collected data were analyzed using SPSS-22. The recorded variables were collected using a predesigned questionnaire and analyzed using regression method.

A total of 120 patients (62 men (51.66%) and 58 women (48.33%)), with the age range of 10-90 years (mean: 48.3 years), entered the study. The mean value of their PaCO₂ and EtCO₂ was 47.45 and 26.9 mmHg, respectively. The mean number of their respiratory rate (RR) was 37.4 bpm, diastolic pressure 89.9 mmHg, and systolic pressure 124.9 mmHg. All demographic data are presented in [Table 1](#).

Corresponding author: Dr Nahid Kianmehr, kianmehr.n@iums.ac.ir

¹ Department of Emergency Medicine, Trauma and Injury Research Center, Iran University of Medical Sciences, Tehran, Iran

² Clinical Research Development Unit, Imam Hossein Hospital, Shahroud University of Medical Sciences, Shahroud, Iran

³ School of Public Health and Safety, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁴ Hazrat Rasool-e-Akram Hospital, Iran University of Medical Sciences and Health Services, Tehran, Iran

⁵ Department of Internal Medicine, Hazrat Rasool-e-Akram Hospital, Iran University of Medical Sciences and Health Services, Tehran, Iran

Table 1. Demographic data

Variable	Value
Male	62 (51.66%)
Female	58 (48.33%)
Mean age of patients	48.33years (SD=22.53)
Asthma	22 (18.33%)
COPD	28 (23.33%)
Pulmonary Edema	12 (10.0%)
Sepsis	8 (6.67%)
PTE*	14 (11.67%)
Others	36 (30.0%)
Mean Respiratory Rate	37.47bpm (SD=9.4)
Mean Systolic BP**	124.9mmHg (SD=14)
Mean Dystolic BP	89.9mmHg (SD=19)

*PTE: Pulmonary thrombosis emboli, **BP: Blood pressure

Based on the results, there was a significant statistical difference between PaCO₂ and EtCO₂ (Spearman correlation, p=0.001, CC=0.436).

In this study, no significant difference was found between systolic and diastolic blood pressure with regards to

PaCO₂ and EtCO₂. However, there was a significant difference between EtCO₂ and PaCO₂ in chronic obstructive pulmonary disease (COPD) and sepsis (Table 2). The statistical difference between EtCO₂ and PaCO₂ was mentioned separately for each disease.

According to the results, there was no significant statistical difference between EtCO₂ and PaCO₂ in age (Table 3). Based on the study results, no significant difference was seen between blood pressure as regards to EtCO₂ and PaCO₂ (Table 4).

In the present study, using Spearman test, no significant difference was found between the respiratory rate as regards to EtCO₂ and PaCO₂ (Table 5). Also, no significant difference existed between the 2 genders in age, respiratory rate, EtCO₂, and PaCO₂. In the linear regression analysis, EtCO₂, with R=0.424, predicted the level of PaCO₂ (p=0.001, Linear Regression).

Blood gas analysis with EtCO₂ can be a noninvasive, quick, and reliable method to predict PaCO₂ in patients

Table 2. Pearson correlation coefficient of Log PaCO₂ and EtCO₂ mentioned separately for each disease

	EtCO ₂	Correlation-coefficient	Log PaCO ₂
Asthma	EtCO ₂	Correlation-coefficient	-0.228
		P-Value	0.308
		N	22
COPD	EtCO ₂	Correlation-coefficient	0.665**
		P-Value	<0.001
		N	28
Pulmonary edema	EtCO ₂	Correlation-coefficient	0.200
		P-Value	0.533
		N	12
Sepsis	EtCO ₂	Correlation-coefficient	0.894**
		P-Value	0.003
		N	8
PTE	EtCO ₂	Correlation-coefficient	0.025
		P-Value	0.933
		N	14

** Correlation is significant at 0.01 level (2-tailed)

Table 3. Spearman correlation coefficient of Log PaCO₂ and EtCO₂ with age

		Correlations	
Spearman rank correlation	EtCO ₂	Correlation coefficient	Age -0.015
		P value	0.869
		n	120
	Log PaCO ₂	Correlation coefficient	0.137
		P value	0.135
		n	120

Table 4. Spearman correlation coefficient of Log PaCO₂ and EtCO₂ with blood pressure

		Correlations Variable		
Spearman rank correlation	EtCO ₂	Correlation coefficient	Diastolic BP -0.034	Systolic BP -0.047
		P value	0.713**	0.609
		n	120	120
	Log PaCO ₂	Correlation coefficient	-0.010	-0.059
		P value	0.918**	0.522
		n	120	120

*BP: Blood Pressure, ** Correlation is significant at 0.01 level (2-tailed)

Table 5. Spearman correlation coefficient of Log PaCO₂ and EtCO₂ with respiratory rate

		Correlations		Variable
Spearman rank correlation	EtCO ₂	Correlation coefficient		Respiratory Rate -0.022
		P value		0.809
		n		120
	Log PaCO ₂	Correlation coefficient		0.109
		P value		0.237
		n		120

with respiratory distress. The linear correlation between EtCO₂ and PaCO₂ was determined and its correlation coefficient was found to be 0.436. The disparity between EtCO₂ and PaCO₂ was an indicative of the difference between ventilated alveoli and perfuse ones. The increase in anatomical and physiological dead space and disruption in pulmonary blood circulation lead to a reduction in EtCO₂ value and an increase in the proportion of EtCO₂/PaCO₂. Also, in hemodynamically stable patients, the gradient of 5-6 is normal (17-19). Pulmonary embolism and shock cause EtCO₂ reduction and increase in the gradient of PaCO₂/EtCO₂ (17, 20).

In the study by Yosefy (19), a linear correlation was found between EtCO₂ and PaCO₂ and the correlation coefficient was reported as 0.736 and 0.772, respectively. These findings confirmed our results. Also, Yosefy showed that aging leads to an increase in the gradient of PaCO₂/EtCO₂ through the increase of dead space in the lungs.

In this study, it was found that as age increased, the PaCO₂/EtCO₂ gradient also increased, but the increase was not significant, which was consistent with other studies. According to Scaman et al (20), capnometer is not able to accurately predict the changes in EtCO₂ when respiratory rate increases and will lead to low correlation coefficient. Likewise, in the present study and Yosefy study, the correlation coefficient reduced when the respiratory rate was above 30, but this reduction was not statistically significant.

The highest correlation coefficient of EtCO₂ and PaCO₂, according to the pathogenesis, was seen between sepsis and COPD and their correlation coefficients were 0.894 and 0.665, respectively. This statistically significant difference indicates that in these 2 diseases the probability to predict PaCO₂ is more than EtCO₂. In the study by Yosefy, these comparisons have not been concluded.

A good correlation was found between PaCO₂ and EtCO₂; however, this correlation was higher in such diseases as sepsis and COPD. Variables such as age, gender, and blood pressure did not affect this correlation. Nevertheless, further studies are needed to confirm these findings in healthy individuals.

Acknowledgments

This study belongs to Dr. Gholamreza Masoumi thesis and was supported by a grant of Iran University of Medical Science (IUMS) to DR. Nahid Kianmehr.

Conflict of Interests

The authors declare that they have no competing interests.

References

1. Godara H. The Washington manual of medical therapeutics: Lippincott Williams & Wilkins; 2013.

2. Siegler EL. The evolving medical record. *Ann Intern Med.* 2010;153(10):671-7.

3. Roberts JR, Hedges JR, Sener S. Clinical procedures in emergency medicine. Publisher: Elsevier; 7 edition; 2018.32

4. Nicholson P, San Miguel C, Kaide CG, Murphy-Lavoie H. Hyperbaric Oxygen Therapy in Emergency Medicine. *Emerg Med Rep.* 2018;39(13).

5. Stacey M. General anesthesia and failure to ventilate. Crises in Childbirth-Why Mothers Survive: CRC Press; 2018. p. 19-46.

6. Long B, Koyfman A, Vivirito MA. Capnography in the Emergency Department: A Review of Uses, Waveforms, and Limitations. *J Emerg Med.* 2017;53(6):829-42.

7. Schober A, Feiner JR, Bickler PE, Rollins MD. Effects of Changes in Arterial Carbon Dioxide and Oxygen Partial Pressures on Cerebral Oximeter Performance. *Anesthesiology.* 2018;128(1):97-108.

8. Goonasekera CD, Goodwin A, Wang Y, Goodman J, Deep A. Arterial and end-tidal carbon dioxide difference in pediatric intensive care. *Indian J Crit Care Med.* 2014;18(11):711-5.

9. Piquilloud L, Thevoz D, Jolliet P, Revelly JP. End-tidal carbon dioxide monitoring using a naso-buccal sensor is not appropriate to monitor capnia during non-invasive ventilation. *Ann Intensive Care.* 2015;5(1):2.

10. Ebert TJ, Novalija J, Uhrich TD, Barney JA. The effectiveness of oxygen delivery and reliability of carbon dioxide waveforms: a crossover comparison of 4 nasal cannulae. *Anesth Analg.* 2015;120(2):342-8.

11. Kodali BS, Urman RD. Capnography during cardiopulmonary resuscitation: Current evidence and future directions. *J Emerg Trauma Shock.* 2014;7(4):332-40.

12. Inoue S, Egi M, Kotani J, Morita K. Accuracy of blood-glucose measurements using glucose meters and arterial blood gas analyzers in critically ill adult patients: systematic review. *J Critic.* 2013Apr;17(2):R48.

13. Lorente J, Marañón R, Vázquez P, Míguez C, Mora A, Rivas A. Capnografía no invasiva: valoración del estado ventilatorio en niños diagnosticados de neumonía. *J Pediatr.* 2015 Sep;73(8):180-3.

14. Ickx B, Dolomie JO, Benalouch M, Melot C, Lingier P. Arterial to End-Tidal Carbon Dioxide Tension Differences in Infants and Children. *J Anesthesia Clin Res.* 2015;6(2). doi:10.4172/2155-6148.1000511

15. Sawe HR, Mfinanga JA, Lidenge SJ, Mpondo BC, Msangi S, Lugazia E, et al. Disease patterns and clinical outcomes of patients admitted in intensive care units of tertiary referral hospitals of Tanzania. *BMC Int Health Hum Rights.* 2014;14(1):26.

16. Bernard GR, Artigas A, Brigham KL, Carlet J, Falke K, Hudson L, et al. The American-European Consensus Conference on ARDS. Definitions, mechanisms, relevant outcomes, and clinical trial coordination. *Am J Respir Crit Care Med.* 1994 Mar;149(3):818-24.

17. Segal N, Metzger AK, Moore JC, India L, Lick MC, Berger PS, et al. Correlation of end tidal carbon dioxide, amplitude spectrum area, and coronary perfusion pressure in a porcine model of cardiac arrest. *Physiol Rep.* 2017 Sep;5(17):e13401.

18. Weinreich UM, Thomsen LP, Brock C, Karbing DS, Rees SE. Diffusion capacity of the lung for carbon monoxide—A potential marker of impaired gas exchange or systemic deconditioning in chronic obstructive lung disease? *Chron Respir Dis.* 2015;12(4):357-64.

19. Yosefy C, Hay E, Nasri Y, Magen E, Reisin L. End tidal carbon dioxide as a predictor of the arterial PCO₂ in the emergency department setting. *J Emerg Med.* 2004;21(5):557-9.

20. From RP, Scamman FL. Ventilatory frequency influences accuracy of end-tidal CO₂ measurements: analysis of seven capnometers. *Anesth Analg.* 1988;67:884-6.