

DOPPLER-DERIVED RIGHT VENTRICULAR MYOCARDIAL PERFORMANCE INDEX IN NEONATES: NORMAL VALUES

E. MALAKAN RAD,* M.D., AND
N. MOMTAZMANESH,** M.D.

*From the Department of Pediatric Cardiology, Kashan University of Medical Sciences,
Kashan, I.R. Iran.*

ABSTRACT

Doppler-derived myocardial performance index (MPI), defined as the sum of isovolumetric contraction and relaxation durations divided by ejection time, is an easily measured and reproducible index that shows both systolic and diastolic myocardial function. The goal of this study was to define normal values of right ventricular MPI in neonates in the first 48 to 72 hours of life.

Fifty-one quiet or sleeping healthy and term neonates underwent complete M-mode, two-dimensional color-Doppler echocardiographic examination and RV MPI was calculated in them. Statistical analysis was performed using SPSS software. A P value less than 0.05 was considered as significant.

RV MPI was shown to be 0.23 ± 0.14 in healthy neonates. There was no correlation between RV MPI and either age or weight of the newborn infants. While this value closely resembles the results of some researchers, it is lower than that previously reported by other investigators in healthy children. The lower value of RV MPI in neonates may be possibly related to the higher pulmonary arterial pressure or right ventricular mass in the first few days of life in neonates. Further study to evaluate the effect of pulmonary arterial pressure and right ventricular mass on RV MPI is recommended.

MJIRI, Vol. 16, No. 3, 129-132, 2002.

Keywords: RV MPI, neonates, normal values.

INTRODUCTION

Traditionally assessment of ventricular systolic function by two-dimensional echocardiography has been based on geometric models of ventricular shape. However this kind of assessment of ventricular ejection frac-

tion and volume may be difficult to obtain because of poorly defined ventricular endocardial borders and complex ventricular geometry in congenital heart diseases.^{1,2} On the other hand, Doppler analysis of the tricuspid and mitral inflow pattern may be limited because of fusion of the early and late inflow waves during tachycardia, and a normal inflow pattern may be difficult to separate from pseudo-normalization.^{3,4} Furthermore these traditional parameters of systolic and diastolic myocardial function are influenced by several factors, including preload, afterload and heart rate.⁵⁻¹¹

Myocardial performance index (MPI) or Tei index,¹² a recently proposed Doppler-derived time interval index, combines both systolic and diastolic time intervals to generate a combined index of global ventricular func-

*Assistant professor of pediatric cardiology.

**Assistant professor of pediatric hematology and oncology and deputy for research affairs of the Kashan University of Medical Sciences, P.O.Box 18155/111, Qotb Ravandi Blvd. Kashan, (Zip Code: 87154) Telephone: 1) In Kashan: (0361) 552999, 2) In Tehran: 8268752 (Malakan), 2841523 (Momtazmanesh), 8274759 (Momtazmanesh), Fax: (0361) 552999. E-mail: malakan_rad@kaums.ac.ir, Momtazmanesh@kaums.ac.ir

tion, that is independent of geometric assumptions.

This study was designed to define normal values of RV MPI in healthy neonates.

PATIENTS AND METHODS

Fifty - one neonates including 26 female and 25 male infants were enrolled in the study. In 2001, during a three-month period, all neonates born at Beheshti's General Hospital were examined on Saturdays every week. Only those babies with a completely normal physical examination including a thorough cardiovascular evaluation were included. Those with any history of birth asphyxia, meconium aspiration pneumonia or any other problem that predisposed the infant to persistent pulmonary hypertension of the newborn (PPHN) were excluded. Because of ethical issues, no blood sampling was performed but those who were suspicious to have hypoglycemia or other electrolyte imbalances (such as premature or IUGR newborns, infants of diabetic mothers, macrosomic babies, ...) ¹³ were all excluded.

Data including prenatal history, mode of delivery (normal vaginal vs. cesarean section), gestational age, postnatal age (as hours), sex, weight and the state of infant alertness (asleep or awake but quiet) were recorded. The preterm neonates were excluded. Also, the crying infants were either soothed in some way or otherwise excluded from the study.

Then all the neonates who fulfilled the criteria underwent a complete two-dimensional and color-Doppler echocardiographic study within 48 to 72 hours after birth, using a Hewlett-Packard Sonos 1000 ultrasound system and a 5 MHZ transducer.

After routine echocardiographic examination, the tricuspid inflow waves were recorded from the apical four-chamber view with the pulsed-wave Doppler sample volume positioned at the tips of tricuspid leaflets in diastole; Doppler signals were displayed at a paper speed of 100mm/ second. Right ventricular (RV) ejection time was measured from the parasternal short-axis scan plane with a pulse-wave Doppler signal placed at the pulmonary valve annulus in the RV outflow tract. Calculation of isovolumic contraction time (ICT), isovolumic relaxation time (IRT) intervals and RV MPI are demonstrated in Fig. 1. To account for slight variations in R-R cycle




Fig. 1. Schema for measurement of Doppler intervals. The myocardial performance index is defined as $a-b/b$, where a is the interval between cessation and onset of the tricuspid inflow, and b is the ejection time (ET) of the right ventricular outflow tract. Isovolumetric relaxation time (IRT) is measured by subtracting the interval c between the R wave and the onset of tricuspid inflow. Isovolumetric contraction time (ICT) is obtained by subtracting IRT from $a-b$. ECG, Electrocardiogram.

length, each time interval was measured on >5 consecutive beats and then averaged.

Statistical analysis

Data are expressed as mean \pm SD or percentages where appropriate. A Mann-Whitney rank-sum test was used to compare age, weights and MPI in the two sexes; linear regression analysis was used to assess the relation of MPI to age, sex and weight.



Statistical significance was defined as P value < 0.05.

Table I. Normal values of RV MPI in neonates.

Variable	Female (n=26)	Male (n=25)	Total (n=51)
Age (hours)	34.2 \pm 14.5	25.0 \pm 11.6	29.74 \pm 13.8
Weight (Grams)	3028.1 \pm 527.2	3296.7 \pm 610.8	3158.7 \pm 575.9
MPI	0.21 \pm 0.118	0.27 \pm 0.15	0.23 \pm 0.14

Table II. Correlation coefficient between RV MPI and age and weight.

	Correlation coefficient	P value
MPI * Age	0.227	0.11
MPI * weight	0.021	0.45

**Fig. 2.** Four-chamber view, measurement of the interval between cessation and onset of the tricuspid inflow.**Fig. 3.** Parasternal short-axis view, measurement of the ejection time (ET) of the right ventricular outflow tract.

RESULTS

Fifty-one neonates were studied. Findings are shown as mean \pm SD in Table I. MPI values did not show a statistically significant difference between the two sexes (P value= 0.22). There was no correlation between MPI and variables of age and weight (Table II, Figures 2 and 3).

DISCUSSION

Our study demonstrated that the normal range of RV MPI in neonates is 0.23 ± 0.14 . To date, there have been few studies on normal values of RV MPI in neonates. As to our knowledge, on the basis of searching in the pubmed, the first study was by Tsutsumi¹⁴ and his colleagues in Japan and the second one by Eidem and his colleagues.¹⁶ It should be cited that there is slight discrepancy in the reported values of the above researches. Eidem and his colleagues reported a normal value of 0.35 ± 0.05 for RMPI in neonates and concluded that advancing gestational age causes no significant change in the fetal RV MPI. On the other hand Tsutsumi and his coworkers reported that Tei index undergoes gradual decrease with increasing gestational age. RV MPI has also been calculated in healthy children. Again, values of 0.24 ± 0.04 by Ishii¹⁵ and 0.32 ± 0.03 by Eidem,¹⁶ reflect the existing discrepancy in this regard. While our results show a lower RV MPI in neonates in comparison with those reported by Eidem for healthy children and neonates, they closely resemble the values obtained by Ishii in healthy children.

Looking back to the formula of RV MPI, we expect that our lower values be due to either a shorter interval between tricuspid inflow waves, i.e., an overall more rapid heart rate in neonates, or a more prolonged pulmonary ejection time. However, previous studies have shown that the correlation between MPI and heart rate is weak.¹⁷ The higher pulmonary arterial pressure within the initial 48 to 72 hours of life may be partly responsible for our findings. To date, the relationship between pulmonary arterial pressure and RV MPI has not been directly delineated. However, Caso and his colleagues studied the association between myocardial right ventricular relaxation time and pulmonary arterial pressure in chronic obstructive lung disease and concluded that myocardial relaxation time was positively related to pulmonary systolic pressure.¹⁸

Lack of correlation between RV MPI and age or weight of the neonates in our study is well explained by the homogenous nature of our studied group of neonates.

Overall, RV MPI is an easily measured index of ventricular myocardial performance, applicable in a wide variety of heart diseases either congenital or acquired.¹⁹⁻²⁴

REFERENCES

1. Snider AR: Methods for obtaining quantitative information from the echocardiographic examination. In: Snider AR, Serwer GA, Ritter SB (eds), *Echocardiography in Pediatric Heart Disease*. St. Louis: Mosby Year Book, pp. 133-236, 1997.
2. Colan SD: Principles of Echocardiography. In: Chang AC,

- Hanley FL, Wernovsky G, Wessel DL, (eds.), *Pediatric Cardiac Intensive Care*. Philadelphia: Williams and Wilkins, pp. 425-440, 1998.
3. Grodecki PV, Klein AL: Pitfalls in the echo-Doppler assessment of diastolic function. *Echocardiography* 10:213-34, 1993.
4. Appleton CP, Hatle LK: The natural history of left ventricular filling abnormalities: assessed by two-dimensional and Doppler echocardiography. *Echocardiography* 9: 437-457, 1992.
5. Nishimura RA, Abel MD, Housmans PR, Wames CA, Tajik AJ: Mitral flow velocity curves as a function of different loading conditions: evaluation by intraoperative transesophageal Doppler echocardiography. *J Am Soc Echocardiography* 2: 79-87, 1989.
6. Stoddard MF, Pearson AC, Kern MJ, Ractcliff J, Mrosek DG, Labovitz AJ: Influence of alteration of preload on the pattern of left ventricular diastolic filling as assessed by Doppler echocardiography in humans. *Circulation* 79: 1226-36, 1989.
7. Gardin JM, Rohan MK, Davidson DM, et al: Doppler transmitral flow velocity parameters: relationship between age, body surface area, blood pressure and gender in normal subjects. *Am J Noninvas Cardiol* 1: 3-10, 1987.
8. Demaria AN, Neumann A, Schubart PJ, Lee G, Mason DT: Systematic correlation of cardiac chamber size and ventricular performance determined with echocardiography and alterations in heart rate in normal persons. *Am J Cardiol* 43: 1-9, 1979.
9. Erbel R, Schwiezer P, Krebs W, Langen HJ, Meyer J, Effert S: Effects of heart rate on left ventricular volume and ejection fraction: a two-dimensional echocardiography study. *Am J Cardiol* 53: 590-7, 1984.
10. Harrison MR, Clifton GD, Pennell AT, DeMaria AN, Ca²ter A: Effect of heart rate on left ventricular diastolic transmitral flow velocity patterns assessed by Doppler echocardiography in normal subjects. *Am J Cardiol* 67: 622-7, 1991.
11. Galderisi M, Benjamine EJ, Evans JC, et al: Impact of heart rate and PR interval on Doppler indexes of left ventricular diastolic filling in an elderly cohort (the Framingham Heart Study). *Am J Cardiol* 72: 1183-7, 1993.
12. Tei C, Ling LH, Hodge DO, Bailey KR, Oh JK, Tajik JK, Seward JB: New index of combined systolic and diastolic myocardial performance: a simple and reproducible measure of cardiac function a study in normal and dilated cardiomyopathy. *J Cardiol* 26: 357-366, 1995.
13. Kalhan SC, Parimi PS: Disorders of carbohydrate metabolism. In: Fanaroff AA, Martin RJ, (eds), *Neonatal-Perinatal Medicine, Diseases of the Fetus and Infant*, 7th edition, St. Louis: Mosby, pp. 1351-1370, 2002.
14. Tsutsumi T, Ishii M, Eto G, Hota M, Kato H: Serial evaluation for myocardial performance in fetuses and neonates using a new Doppler index. *Pediatr Int* 41(6): 722-7, Dec 1999 .
15. Ishii M, Eto G, Tei C, Tsutsumi T, Hashino K, Sugahara Y, et al: Quantitation of the global right ventricular function in children with normal heart and congenital heart disease: a right ventricular myocardial performance index. *Pediatr Cardiol* Sep-21 (5): 416-421, Oct 2000.
16. Eidem BW, Edwards JM, Cetta F: Quantitative assessment of fetal ventricular function: establishing normal values of the myocardial performance index in the fetus. *Echocardiography* 18(1): 9-13, Jan 2001.
17. Poulsen SH, Nielsen JC, Andersen HR: The influence of heart rate on the Doppler-derived myocardial performance index. *J Am Soc Echocardiogram* 13: 379-84, 2000.
18. Caso P, Galderisi M, Cicala S, Cioppa C, et al: Association between myocardial right ventricular relaxation time and pulmonary arterial pressure in chronic obstructive lung disease. Analysis by pulsed Doppler tissue imaging. *J Am Soc Echocardiogr* 14 (10): 970-977, Oct 2001.
19. Poulsen SH, Jensen SE, Tei C, Seward JB: Egstrup K, et al: Value of the Doppler index of myocardial performance in the early phase of acute myocardial infarction, *J Am Soc Echocardiogr* 13:723-730, 2000.
20. Eidem BW, O'Leary P, Tei C, Seward JB: Usefulness of the myocardial performance index for assessing right ventricular function in congenital heart disease. *Am J Cardiol* 86:654-658, 2000.
21. Dujardin KS, Tei C, Yeo TC, Hodge DO, Rossi A, Seward JB: Prognostic value of a Doppler index combining systolic and diastolic performance in idiopathic-dilated cardiomyopathy. *Am J Cardiol* 82:1071-1076, 1998.
22. Khodes J: Another useful index of ventricular function in patients with single ventricle. *Am J Cardiol* 87 (9): 1137-1142, 2001.
23. Eidem BW, Sapp BG, Suarez CR, Cetta F: Usefulness of the myocardial performance index for early detection of anthracyclin-induced cardiotoxicity in children. *Am J Cardiol* 87 (9): 1120-22, 2001.
24. Ishii M, Tsutsumi T, Himeno W, Eto G, Furui J, Hashino K, Sugahara Y, et al: Sequential evaluation of left ventricular myocardial performance in children after anthracycline therapy. *Am J Cardiol* 86: 1279-1281, December 1, 2000.