





Med J Islam Repub Iran. 2023 (2 Aug);37.86. https://doi.org/10.47176/mjiri.37.86

Cost-Effectiveness of Health Literacy Enhancement Intervention on Patients with Type II Diabetes

Mahshid Lotfi¹, Farhad Lotfi^{2,3}, Shahram Sedghi¹*⁰, Sirous Panahi¹

Received: 15 Oct 2022 Published: 2 Aug 2023

Abstract

Background: Type II diabetes is considered a chronic disease that influences the affected person's quality of life and imposes a high economic burden on the patient and society. The enhancement of health literacy seems essential for self-management and disease control in patients with type II diabetes. Consequently, this study was performed to evaluate the cost-effectiveness of enhanced health literacy in type II diabetes patients.

Methods: This study was of quantitative and economic evaluation type. A population of 232 patients was selected among those referred to the Tafihan Shiraz clinic based on the quality of entry and exit. The health literacy educational intervention was carried out for three months. In order to collect information, researchers used the SF36 questionnaire and the checklist of costs. The Cost Effectiveness Ratio (CER) and Incremental Cost Effectiveness Ratio (ICER) were calculated. Moreover, the costs of each unit of increasing the quality of life before and after the intervention were calculated. A decision was made to determine the cost-effectiveness of the intervention.

Results: According to the results, 40% of the participants were females, and 34.5% were 30-40 years old. Examination of the quality of life in patients before the intervention indicated that the mean and standard deviation of the patients' quality of life before the intervention was 18.43 ± 44.99 , and the mean and standard deviation of the quality of life after the intervention was 49.57 ± 16.21 . Moreover, the patients' quality of life increased after the educational intervention. The total direct medical costs were \$717,484 and \$685,620 before and after the intervention, respectively. The total indirect costs were \$122,535 and \$122,119 before and after the intervention, respectively. The study was cost-effective with CER=0.9 and ICER= - 140,000 per increase in the quality of life.

Conclusion: Improving health literacy can have a range of benefits and improve the quality of life of patients with type II diabetes and reduce their treatment costs, and thereby, this may be seen as an effective step toward the recovery of patients with type II diabetes.

Keywords: Cost-Effectiveness, Quality of Life, Health Literacy

Conflicts of Interest: None declared

Funding: This study was supported as a doctoral dissertation by grant No. (IR.IUMS.REC.1399.4243) from the Research Deputy of the Iran University of Medical Sciences, Iran.

*This work has been published under CC BY-NC-SA 1.0 license. Copyright© <u>Iran University of Medical Sciences</u>

Cite this article as: Lotfi M, Lotfi F, Sedghi S, Panahi S. Cost-Effectiveness of Health Literacy Enhancement Intervention on Patients with Type II Diabetes. *Med J Islam Repub Iran.* 2023 (2 Aug);37:86. https://doi.org/10.47176/mjiri.37.86

Introduction

Health literacy is an acquired skill and is necessary for managing individual health. In addition, health literacy is related to literacy and requires people's knowledge, motivation, and competencies in accessing, understanding, evaluating, and using health information for decision-making. In order to remain compliant with their treatment and achieve desirable health outcomes, patients must have a high level of health literacy (1). People with inadequate

↑What is "already known" in this topic:

The improvement of health literacy improves the quality of life of type 2 diabetes patients. The improvement of health literacy reduces treatment costs for patients with type 2 diabetes.

\rightarrow *What this article adds:*

Considering the cost-effectiveness of the health literacy intervention, the improvement of the health literacy of type 2 diabetes patients in Iran has many limitations, most of which are due to the lack of expertise and specialist staff in treatment centers for type 2 diabetes patients.

Downloaded from mjiri.iums.ac.ir on 2025-07-01

Corresponding author: Dr Shahram Sedghi, sedghi.s@iums.ac.ir

¹ Librarianship and Medical Information Department, School of Health Management and Information Sciences, Iran University of Medical Sciences, Tehran, Iran

² National Center for Health Insurance Research, Tehran, Iran

^{3.} Health Human Resources Research Center, School of Management and Medical Information Sciences, Shiraz University of Medical Sciences, Shiraz, Iran

health literacy have worse health outcomes (1, 2). Approximately 25% of Americans have difficulty understanding health care information. In Iran, 50% of adults have limited health literacy (3). Additionally, racial and ethnic minorities with low health literacy rates among older adults are more susceptible to chronic illnesses (4).

Diabetes is a chronic disease. According to the International Diabetes Federation estimates, 463 million people around the world have diabetes, and this is expected to increase by up to 700 million patients. Most diabetic patients have type II diabetes, which accounts for 95% of the total population of diabetic patients. Currently, 8.5% of the American adult population is suffering from type II diabetes (5). Developing type II diabetes is caused by unhealthy lifestyle activities such as physical inactivity and poor diet (6). People with type II diabetes usually get aware of their disease late, and about 30%-80% of people with type II diabetes have not yet been diagnosed. The treatment of type II diabetes needs constant management of the lifestyle aspects such as diet, exercise, glucose monitoring, and especially some specific medications to achieve glycemic control goals. People with type II diabetes are expected to reduce or stop the progression of their disease by employing different intervention approaches (7).

Studies have demonstrated inadequate health literacy as one of the biggest challenges in the management and treatment of type 2 diabetes (5). Health literacy describes how much patients can acquire, process, and understand basic health information about managing and controlling type II diabetes. It is particularly important and crucial that the patients can understand the instructions related to the drugs prescribed for their treatment (8, 9). The low health literacy of type II diabetes patients about diet and adequate physical activity are known as the most important predisposing factors for developing complications, and the mortality of patients with type II diabetes increases since more than 95% of the care measures are accomplished by the patients (10, 11).

In Iran, 43.47% of people have inadequate and borderline health literacy, and 26.3% have adequate health literacy (12). A low level of health literacy is common in type II diabetic patients (13, 14). Coherent educational programs promote the awareness, attitude, and performance of diabetic patients and reduce treatment costs (15, 16).

The provision of proper conditions for training and increasing the knowledge of patients can decrease the costs of the health system. Some significant steps in reducing treatment costs include self-care programs and persuading type II diabetes patients to adhere to such programs (10). The cost-effectiveness method is an economic evaluation approach that focuses on the method to achieve objectives. This method helps decision-makers choose and implement different strategies. The examination of the cost-effectiveness of various types of therapeutic and non-therapeutic interventions plays a substantial role in the design and evaluation of health systems (17). Abdul Ahad et al. (2019) showed that examining the cost-effectiveness of health literacy provides ways to waste resources and costs. It also determines the effectiveness of health literacy intervention in treatment and has an effective role in the management of health care services in the field of diabetes (11-13, 18). The effect of enhancing health literacy in reducing direct medical and non-medical costs and increasing patients' quality of life has not been studied in Iran. Therefore, the present research was conducted to evaluate the cost-effectiveness of health literacy intervention in type II diabetic patients.

Methods

This research was of the type of economic evaluation. The sampling of patients was performed by a random sampling method, and samples were selected based on the inclusion and exclusion criteria. The sample size was obtained using the study of Pourreza et al. in 2013 (19). The mean and standard deviation in the same study was 71.97 ± 13 before the intervention and 76.06 ± 11.08 after the intervention. According to similar studies, the following formula was used to estimate the sample size as 210 subjects (19). According to the inclusion and exclusion criteria and the number of type II diabetes patients referred to Tafihan Shiraz clinic, and considering the probability of sample drop, a total of 232 patients were selected.

$$N = 2C \times \left[\frac{\sqrt{SD_1^2 + SD_2^2}}{m_1 - m_2} \right]^2$$

The inclusion criteria were as the following:

1. Patients aged 20-60 years old referred to the clinic

2. Diagnosed with type II diabetes by endocrinologists

3. Diagnosed with type II diabetes a year before the research

4. Ability to understand the Persian language

5. Willingness and informed consent to participate in the research

6. Referring to the clinic and having a medical record with the attending physician

The exclusion criteria were as the following:

1. Suffering from other chronic diseases except those related to the complications of diabetes

2. Suffering from mental retardation or the diagnosis of other psychological disorders, such as mood disorders and anxiety, before the diagnosis of diabetes

In this research, the cost-effectiveness of health literacy improvement in patients with type 2 diabetes in Shiraz City was investigated before and after the intervention. The demographic information questionnaire was utilized in this research to gather and determine the patients' age, gender, education level, economic status, marital status, and duration of the disease. Moreover, the 36-Item Short Form Survey (SF-36) questionnaire was employed to evaluate the quality of life of patients. This 36-item questionnaire is not limited to a certain age range, group, or disease. The goal of this questionnaire is to evaluate the state of health from physical and mental perspectives, which is obtained by combining the scores of the eight health domains. Eight areas of the SF-36 questionnaire are as follows:

1. General Health

- 2. Physical functioning
- 3. Role limitation due to physical reasons

- 4. Role limitation due to emotional reasons
- 5. Physical pain
- 6. Social functioning
- 7. Energy and vitality
- 8. Mental health

The reliability of the SF-36 questionnaire has been obtained to be 0.9 for diabetic patients (16). It should be noted that the benchmark in this study was to calculate the costs from the society's viewpoint, which encompassed direct treatment and non-treatment costs, and indirect costs. The information collection form included two parts. The first part involved the demographic information of the patient. The second part included information related to direct medical costs (e.g., the cost of physicians' visits, the cost of diabetes drugs, hospital costs, diagnostic and laboratory procedures), direct non-medical costs (e.g., transportation, lodging, and telephone costs, and the cost of informal care), and indirect costs (e.g., the lost productivity of the patient and lost productivity of the patient's companion) all of which were extracted through the form and calculated with the human capital approach method. Each dollar was considered to be equal to 41,790 Rials in this research, which is the official currency rate (17).

The researcher was present at Tafihan Health Center for three months (December, January, and February 2019). Accordingly, 24 face-to-face training sessions of 30-45 min were held in the form of small 13-20 people groups on Sundays and Wednesdays. The health literacy of the sample patients was improved with the help of the DELNET 2 educational model (18). This tool contains educational materials about blood sugar, nutrition, and various drugs in the area of diabetes, which is recognized as a standard tool by the American Diabetes Association and has been designed by nutritionists, physicians, and nurses. The educational materials of this tool were provided to all patients in the form of in-person training. Moreover, the researcher introduced and presented educational information sources and other useful treatment guides available in the field of type II diabetes (brochures, book introduction) to the patients. The intervention related to health literacy enhancement was carried out for three months, and the patients were followed up by the researcher for three months. The data obtained from the questionnaires were analyzed by the SPSS software (version 21) using the descriptive statistics method and the frequency distribution table before and after the health literacy intervention. The mean and standard deviation values were calculated before and after the intervention. In addition, the paired t-test was used to examine the changes before and after the intervention. Finally, the costeffectiveness of health literacy enhancement was calculated and described using the cost-effectiveness formula.

Data analysis

Cost-Effectiveness Ratio (CER)

The cost-effectiveness ratio in this research is the average cost of increasing the quality of life of each patient after the health literacy intervention compared to that before the intervention (17).

Costs Before Health Literacy Intervention

The costs of type 2 diabetes patients decreased before the intervention (xa).

The costs of type 2 diabetes patients did not decrease before the intervention (xb).

Costs After Health Literacy Intervention

The costs of type 2 diabetes patients have decreased after the intervention (yc).

The costs of type 2 diabetes patients have not decreased after the intervention (yd). CER = xa / yc

Where,

CER < 1 means that the intervention is cost-effective.

CER > 1 means that the intervention is not cost-effective

Incremental Cost-Effectiveness Ratio (ICER)

The incremental effectiveness cost ratio is defined as the difference in costs before and after the intervention to increase health literacy divided by the difference in the increase in quality of life before and after the intervention. The incremental cost-effectiveness ratio represents the incremental average associated with a unit of measurement (17).

ICER=xa - yc/ difference in a% - difference in c%

Decision rules

(i) If the incremental cost is negative and the incremental effect is positive (SE quadrant), the intervention is unequivocally cost-effective (it is dominant, achieving better outcomes at lower cost). (ii) If the incremental cost is positive and the incremental effect is negative (NW quadrant), the intervention is unequivocally not cost-effective (it is dominated, achieving poorer outcomes at a higher cost). (iii) If both the incremental cost and the incremental effect are negative (SW quadrant) or both the incremental cost and the incremental effect are positive (NE quadrant), no such unequivocal statements can be made. Determining whether the intervention is cost-effective depends on a threshold value (λ), defined as the maximum amount society is willing to pay for an incremental health gain or, equivalently, as the minimum amount society is willing to accept for foregoing an incremental health gain. The intervention will be regarded as cost-effective if its incremental cost-effectiveness ratio is lower than the threshold $(\Delta C/\Delta E < \lambda)$ for ICERs in the NE quadrant or higher than the threshold $(\Delta C/\Delta E > \lambda)$ for ICERs in the SW quadrant (17, 20) (Diagram 1).

The data obtained from the questionnaires were analyzed in the SPSS software (version 21) using the descriptive statistics method, the frequency distribution table before and after the health literacy intervention, and the mean and standard deviation before and after the intervention were expressed. The paired t-test was used to examine the changes before and after the intervention. Finally, using the cost-effectiveness calculation, the incremental cost-effectiveness was expressed as the cost-effectiveness of health literacy improvement.

Downloaded from mjiri.iums.ac.ir on 2025-07-01



Diagram 1. Cost-effectiveness plane for after and before Intervention

Results

The findings obtained from the demographic information are presented in Tables 1 and 2. The data were analyzed to investigate the relationship between demographic information and quality of life.

According to the findings, most of the study participants were women (39.1%). Most of the participants were aged 30-39 (34.5%) and had high school education (32.3%), and their income was between \$238-\$450 (37.9%). According to the results of Table 1 and Table 2, there was a significant relationship between demographic characteristics and quality of life before and after the intervention (P<0.001).

According to Table 3, the mean and standard deviation of the three dimensions of "role limitations due to physical health," "role limitations due to emotional problems," and "physical functioning" were obtained higher than other dimensions, while the "pain" dimension had the lowest values. Moreover, there was a significant relationship between quality of life and improvement of health literacy before and after the intervention (P<0.001).

According to Table 4, the total cost of the consuming items for all patients was obtained to be \$708.

According to Table 5, the cost of admission to the emergency unit and the costs of the physicians' visits and medications before the intervention were estimated to be more than others, and the cost of MRI was obtained lower than others. After performing the health literacy intervention, the cost of admission to the emergency unit and the costs of the physicians' visits and surgical services were calculated to be higher than other costs, and the cost of medications was estimated to be lower than others. According to the research hypothesis, the costs decreased after the educational intervention. According to Table 6, before the intervention, the expenses paid by the patient for diet, hiring a servant, and changes in the environment were greater than other costs. After the intervention, the total cost paid by the patients decreased, and the expenses paid by the patient for diet, changes in the environment, and auxiliary tools were more significant compared to other costs.

According to Table 7, the lost income of the patient and the patient's companion decreased by \$417 after the intervention. Moreover, the total lost income of the patient and the patient's companion decreased after the intervention with comparison to the state before the educational intervention.

The intention-to-treat analysis method was utilized to estimate the effect of the intervention accurately. The costs were calculated before and after the health literacy intervention. Moreover, the information related to the quality of life before and after the intervention was expressed as mean±standard deviation. The rate of changes in the patients' costs and quality of life before and after the intervention was calculated using the following formula, and the rate of cost-effectiveness was specified.

 $ICER = \Delta cost / \Delta effectiveness = -1773350$

The intention-to-treat analysis method was used to estimate the effect of the intervention appropriately. The costs were calculated before and after the health literacy intervention. The data related to the quality of life before and after the intervention was expressed as mean \pm standard deviation.

According to Table 8, the costs after the intervention decreased by \$141,858, the quality of life increased, and a significant difference was observed between the promotion of health literacy and demographic variables after the educational intervention; therefore, the intervention was effec-

Variable		Number	Mean ± Standard Deviation
		(%)	
Age	20-29	50 (21.6)	1.46±0.49
-	30-39	80 (34.5)	
	40-49	61 (26.3)	
	50-60	41 (17.7)	
Gender	Female	125 (39.1)	2.4±1.01
	Male	106 (33.1)	
Marital Status	Single	118 (50.9)	1.62 ± 0.75
	Married	85 (36.6)	
	Divorced	8 (19.8)	
Education Level	Illiterate	55 (23.7)	2.5±1.08
	Primary	49 (21.1)	
	High school	75 (32.3)	
	Academic	53 (22.8)	
Occupation	Jobless	60 (25.9)	2.97±1.54
*	Housewife	38 (16.4)	
	Employee	43 (18.9)	
	Retired	29 (12.5)	
	Self-Employed	62 (26.7)	
Income	Less than \$237	42 (18.1)	2.88±1.68
	\$238-\$450	88 (37.9)	
	\$451-\$687	31 (13.7)	
	\$688-\$924	31 (12.9)	
	\$925-\$1161	22 (9.5)	
	\$1162-\$1398	16 (6.9)	
	\$1399-\$1635	2 (0.9)	
	More than \$1635	0	

Table 1 Demographic characteristics of the participants in the study

Table 2. Demographic characteristics of the participants in the study

Variable		Number (%)	Mean ± Standard deviation
Disease Duration	Less than 1 year	54 (23.3)	2.43±1.21
	1-1.9 years	54 (23.3)	
	2-2.9 years	80 (34.5)	
	3-4 years	20 (8.6)	
	More than 4 years	16 (6.9)	
Blood Sugar	Less than 100	80 (53.3)	2.12±1.14
-	100-199	50 (21.6)	
	200-299	60 (25.9)	
	300-400	30 (12.9)	
	More than 400	10 (0.3)	

tive. Since any economic evaluation is associated with uncertainty, it was attempted in this study to test the generalizability of the results using the sensitivity analysis approach. Therefore, a one-way sensitivity analysis was performed, and the quality of life and costs increased by 20% before and after the intervention. Table 9 indicates the results of the sensitivity analysis before and after the intervention. Before the intervention, the new ICER equals 5,709,607 with a 20% increase in the costs, and with increasing the quality of life by 20%, the new ICER becomes 8,681,775. After the intervention, with a 20% increase in the costs, the new ICER equals 1,571,259, and with a 20% increase in the quality of life, the new ICER becomes 778,372.

Cost-Effectiveness Ratio (CER)

The CER was calculated as follows:

CER = 1301868 / 1443726 = 0.9

As a result, the health literacy intervention is cost-effective.

Incremental Cost-Effectiveness Ratio (ICER)

 $ICER = \Delta cost / \Delta effectiveness = -140000$

Considering that a positive CER and a negative ICER were obtained, there is an intervention in the area (SE quadrant); therefore, the intervention is definitely cost-effective.

Since every type of economic evaluation study is associated with uncertainty, it was attempted in the present study to test the generalizability of the results using sensitivity analysis. Therefore, a one-way sensitivity analysis was performed, and the quality and costs increased by 20% before and after the intervention, and the results were presented. Table 9 indicates the sensitivity analysis before and after the intervention. Diagram 2 demonstrates the fluctuations in price and quality of life before and after the intervention. Before the intervention, with a 20% increase in costs, the new ICER becomes 5,709,607, and with a 20% increase in the quality of life, the new ICER becomes 8,681,775. Moreover, after the intervention, with a 20% increase in costs, the new ICER becomes 1,571,259, and with a 20% increase in the quality of life, the new ICER becomes 778,372. As a result, the insensitivity of CER is variable.

The tornado diagram for sensitivity analysis is presented

	Before the Intervention	After the Intervention
Dimensions	Mean ± Standard Deviation	Mean ± Standard Deviation
Physical Functioning	49.50±14.35	50.28±16.41
Role Limitations due to Physical Health	66.95±18.88	69.76±24.34
Role Limitations due to Emotional Problems	58.62±19.18	70.83±30.48
Energy/Fatigue	44.87±20.26	42.37±17.44
Emotional Well-Being	44.91±19.37	48.56±19.68
Social Functioning	33.96±33.44	38.72±35.8
Pain	27.78±25.63	24.51±26.45
General Health	33.40±20.70	49.55±26.10
Total Mean	44 99+18 43	49 57+16 21

Table 4. Costs paid for the health literacy therapeutic intervention

Consuming Items	Costs (Dollars)
Cost of Printing the DELNET Booklet	498
Cost of the Instructor's Transportation	194
Total Cost	708

Table 5. Direct medical costs of type II diabetes before and after the health literacy educational intervention

Different Types of Direct Costs	Average Cost Per Pa-	Total Cost (Dollars)	Average Cost Per Patient	Total Cost (Dollars) After
	tient (Dollars) Before	Before the Interven-	(Dollars) After the Inter-	the Intervention
	the Intervention	tion	vention	
Physicians' visits	55	127,221	413	95,949
Admission to the Emergency	57	132,196	470	109,216
Unit				
Laboratory tests	427	99,029	297	6894
Medications	545	126,273	281	65,150
Surgical services	400	92,869	393	91,211
Physiotherapy	398	92,395	398	92,158
Radiology	250	58,043	409	94,764
MRI	244	56,622	295	68,230
Total Costs	2377	717,484	2956	685,620

Table 6. Direct non-medical costs of type II diabetes before and after health literacy educational intervention

Different Types of Direct Non-Medical Costs	Average Cost Per Patient (Dollars) Before the In-	Total Cost (Dollars) Be- fore the Intervention	Average Cost Per Patient (Dollars) Before the In-	Total Cost (Dollars) After the Intervention
	tervention		tervention	
Auxiliary Tools	444	102,819	333	77,233
Child Care	433	100 450	301	6941
Hiring Servants	610	141 436	323	75 101
Diet Costs	622	144 279	600	139 303
Changes in the Environ-	496	115 139	569	131,959
ment	.,,,	110,109	2007	101,909
Total Costs	2604	604,122	2127	493,011
Table 7 Indirect costs of type	II diabetes before and after t	he health literacy educational	intervention	
	Average Cost Per Pa-	Total Cost (Dollars) Be-	Average Cost Per Patient	Total Cost (Dollars) After
Indirect Costs	tient (Dollars) Before	fore the Intervention	(Dollars) After the Inter-	the Intervention
manoet costs	the Intervention	fore the intervention	vention	
Lost Income of Patient	501	116,560	53	12,248
Lost Income of the Patient's	118	5559	10	518
Companion				
Total Lost Income	619	122,119	63	122.535

Table 8. Changes in the cost and quality of life before and after the health literacy intervention

	Total Cost (Dollars)	Quality of Life	Differences Between the Costs (Dollars)	Quality of Life Difference
Before the Intervention	1,443,726	0.461094348659		
After the Intervention	1,301,868	0.54	141,868	0.08

as Diagram 3 for evaluating the strength of the results of the present investigation.

Discussion

This study was designed and conducted to evaluate the

cost-effectiveness of the health literacy intervention in patients with type II diabetes. According to the results, a significant difference was found between demographic variables and quality of life before and after the intervention. The mean and standard deviation for quality-of-life dimensions increased after the intervention. The total mean and

6 <u>http://mjiri.iums.ac.ir</u> Mad Lislam Rapub Iran 2

Med J Islam Repub Iran. 2023 (2 Aug); 37:86.

Table 9. One-way sensitivity analysis of costs and the quality of life before and after the intervention

One-Way Sensitivity Analysis	Result	One-Way Sensitivity Analysis	Result
20% Increase in Quality of Life Before Interven-	0.60	20% Increase in Quality of Life Before Intervention	0.60
tion			
Cost Difference	1,425,778	Cost Difference	1,425,778
Quality of Life Difference	0.02	Quality of Life Difference	0.18
New ICER	8,681,775	New ICER	778,372
20% Increase in Cost Before Intervention	17,411,330	20% Increase in Cost Before Intervention	15,700,396
Cost Difference	4,327,666	Cost Difference	1,190,955
Quality of Life Difference	0.08	Quality of Life Difference	0.08
New ICER	5,709,607	New ICER	1,571,259



Diagram 2. Incremental Cost-Effectiveness Ratio; after and before Intervention



Diagram 3. Tornado diagram for sensitivity analysis to evaluate the strength of the results

standard deviation of the dimensions of self-care was 44.99 ± 18.43 before the intervention, which became 49.57 ± 16.21 after the intervention (20). The research findings indicated that the quality of life has improved after the educational intervention. Chittleborough et al. (21) demonstrated that the quality of life of patients with diabe-

tes has a significant relationship with demographic characteristics, which is consistent with the results of the present investigation. Their results also indicated that the mean and standard deviation values of the quality-of-life dimensions, including mental health, physical functioning, and pain, are less than other dimensions. In the present study, the mean and standard deviation of social functioning, pain, and general health were higher than other dimensions before the intervention, while the mean and standard deviation of pain, social functioning, and energy/fatigue were lower than other dimensions after the intervention. Regarding the pain dimension, the results of this study were in line with Chittleborough et al.'s study.

Moreover, Cheng et al. (22) indicated that increasing knowledge improves the quality of life of patients with diabetes. In 2011, Vadstrup et al. (23) indicated that increasing the awareness and health literacy of patients with type II diabetes would enhance their quality of life, which is in line with the results of the present study. In recent years, several studies have been conducted regarding the quality of life of patients with diabetes, and the results have shown that the quality of life of these patients is lower than that of healthy people. Moreover, diabetes has a negative effect on a person's understanding and satisfaction with his/her life and causes a decrease in functional capacity, mental states, social functioning, and in general, the quality of life (24-26). Chen et al. (27) revealed that educational intervention in type II diabetes patients reduces blood sugar levels and improves the quality of life of patients, which is in line with the results of the present investigation. Moreover, Yuan et al. (28) indicated that health literacy enhancement intervention increases the quality of life of patients, which is also consistent with the results of this study.

In the present study, the direct and indirect costs of type II diabetes were defined and specified from the patients' perspective before and after the health literacy enhancement intervention. The total direct medical costs before and after the intervention were equal to \$717,484 and \$685,620, respectively. Moreover, the total indirect medical costs before and after the intervention were \$604,122 and \$493,011, respectively. In addition, the total indirect costs before and after the intervention were \$122,535 and \$122,119, respectively. The costs indicated a reduction after the intervention, and the intervention appeared influential. Kieu et al. (29) revealed that the total direct non-medical and indirect costs in 2017 accounted for 239 million dollars. The direct non-medical costs were estimated at 78 million dollars, while indirect costs accounted for 161 million dollars. The costs of being absent, attending sessions, and premature mortality corresponded to 17%, 73%, and 10% of the indirect costs. The average annual direct nonmedical costs for patients were equal to US\$56. The average annual absence and presence costs for working-age patients were equal to US\$61 and US\$267, respectively. Researchers have concluded that patients with type II diabetes have a high economic burden, which is in line with the results of this study. Sulistyaningrum et al. (30) evaluated the direct and indirect costs of diabetes in Indonesia. Accordingly, the average annual cost per patient accounted for 1207.8±673.4 US dollars. The direct cost accounted for the largest share of the total cost (87.95%), and the cost of medications was the significant share after the visiting fee of the general practitioners. Their findings seem consistent with the results of the present study. The average annual cost of using insulin with an oral hypoglycemic agent, the presence of hypertension and dyslipidemia, and poor levels were the highest frequent characteristics in each group. Saha et al. (31) indicated that the average total cost per year was \$600, covering the costs of medication, research, consultation, and transportation. Among them, the costs of medications were the highest (US\$350). Therefore, the cost of treating type II diabetes was proven to be high, which is consistent with the results of the present investigation.

In the present study, the cost-effectiveness of health literacy intervention was defined based on community costs. According to the results, the costs decreased, and the quality of life increased after the intervention. Therefore, the intervention has undoubtedly been predominantly effective. In a review of the studies on the cost-effectiveness of diabetes, it was found that the core emphasis of these studies has been on the economic evaluation of third-level preventions, i.e., various types of medication and behavioral interventions. In 2018, Saifulsyahira et al. (17) indicated that the health literacy improvement intervention reduces blood sugar and the treatment costs of patients with type 2 diabetes, which is in line with the present study.

Conclusion

It is concluded from the results of this research that an increase in health literacy is cost-effective and will increase the quality of life of patients and reduce their treatment costs. Therefore, it seems essential to increase the health literacy of all patients with type II diabetes. In order to improve the status of educating patients, it is necessary to align managers in this field and increase the recruitment of health literacy specialists, including librarians and medical informants. The present study demonstrated that inadequate health literacy is a common problem in patients with type II diabetes and imposes more costs on patients and society. Since the enhancement of the health literacy of patients with type II diabetes requires continuous efforts for several years, it is recommended to perform some relevant investigations on how to include related materials in textbooks as well as utilize them in social media and design mobile applications for patients to improve the health literacy level.

Limitations

One of the limitations of the present study was the noncooperation of the patients in the educational sessions. Moreover, the patients tried not to mention the treatment costs, which did not affect the participants' responses. In this research, the study samples were limited to Shiraz in Iran. As a result, a limited generalization is considered for obtained results.

Research Suggestions

The results indicated that the training of type 2 diabetes patients increases the quality of life and reduces the costs of the patients. As a result, it is suggested to plan for the training of type 2 diabetes patients in related clinics. Communication should be made to exchange knowledge between type 2 diabetic patients and clinics to increase health literacy. Moreover, future research should be conducted mainly with the aim of evaluating the effectiveness of evidence-based interventions to reduce costs. Furthermore, systematic reviews are required to better understand the effects of health literacy on the costs that type 2 diabetes patients should pay.

Ethical Approval

This study was conducted with the approval of the ethics committee of Iran University of Medical Sciences, Iran (IR.IUMS.REC.1399.424).

Consent for Participation and Publication

Prior to completing the Summary of Diabetes Self-Care Activities (SDSCA) questionnaire, some explanations were given to patients about the confidentiality and publication of research results, and they completed the questionnaire with satisfaction.

Acknowledgments

The authors appreciate the thoughtful comments and suggestions provided by the reviewers of this manuscript. Also, we thank all the participants for their time and consideration.

Conflict of Interests

The authors declare that they have no competing interests.

References

- Sørensen K, Van den Broucke S, Fullam J, Doyle G, Pelikan J, Slonska Z, et al. Health literacy and public health: a systematic review and integration of definitions and models. BMC Public Health. 2012;12(1):1-13.
- Berkman ND, Sheridan SL, Donahue KE, Halpern DJ, Crotty K. Low health literacy and health outcomes: an updated systematic review. Ann Intern Med. 2011;155(2):97-107.
- 3.Caballero J, Jacobs RJ, Ownby RL. Development of a computerized intervention to improve health literacy in older Hispanics with type 2 diabetes using a pharmacist supervised comprehensive medication management. Plos One. 2022;17(2):e0263264.
- 4.Caruso R, Magon A, Baroni I, Dellafiore F, Arrigoni C, Pittella F, et al. Health literacy in type 2 diabetes patients: a systematic review of systematic reviews. Acta Diabetol. 2018;55(1):1-12.
- 5.Abdullah A, Liew SM, Salim H, Ng CJ, Chinna K. Prevalence of limited health literacy among patients with type 2 diabetes mellitus: A systematic review. PLoS One. 2019;14(5):e0216402.
- 6.Marciano L, Camerini A-L, Schulz PJ. The role of health literacy in diabetes knowledge, self-care, and glycemic control: a meta-analysis. J Gen Intern Med. 2019;34(6):1007-17.
- 7.Hussain N, Said AS, Khan Z. Influence of health literacy on medication adherence among elderly females with type 2 diabetes in Pakistan. Int Q Community Health Educ. 2020;41(1):35-44.
- 8.Shan R, Sarkar S, Martin SS. Digital health technology and mobile devices for the management of diabetes mellitus: state of the art. Diabetologia. 2019;62(6):877-87.
- 9.Rahnavard S, Elahi N, Rokhafroz D, Hagighi MH, Zakerkish M. Comparison of the effect of group based and mobile based education on self-care behaviors in type ii diabetic patients. Iran J Diabetes Lipid Disord. 2019;18(2):55-63.
- Medina LA, Silva RA, de Sousa Lima MM, Barros LM, Lopes RO, Melo GA, Garcia Lira Neto JC, Caetano JÁ. Correlation Between Functional Health Literacy and Self-efficacy in People with Type 2 Diabetes Mellitus: Cross-sectional Study. Clin Nurs Res. 2022 Jan;31(1):20-8.
- 11. Pashaki MS, Eghbali T, Niksima SH, Albatineh AN, Gheshlagh RG. Health literacy among Iranian patients with type 2 diabetes: A systematic review and meta-analysis. Diabetes & Metabolic Syndrome: Clin Res Rev. 2019;13(2):1341-5.
- 12.Seyedoshohadaee M, Kaghanizade M, Nezami M, Hamedani B,

Barasteh S. The Relationship Between Health Literacy And General Health In Patients With Type 2 Diabetes. Iran J Diabetes Lipid Disord. 2016;15(5):312-9.

- Protheroe J, Rowlands G, Bartlam B, Levin-Zamir D. Health literacy, diabetes prevention, and self-management. J Diabetes Res. 2017;2017.
- 14.de Abreu IR, Baía C, Silva JM, Santos AM, Oliveira M, Castro F, Mozes M, Ferreira R, Alves LA. LitKDM2 study: the impact of health Literacy and knowledge about the disease on the metabolic control of type 2 diabetes mellitus. Acta Diabetol. 2022 Jun;59(6):819-25.
- 15.Akbari M, Ashrafi E, Rajab A. Effetiveness of instruct cognitive errors in the way of philisophy for children and adolscents in cognitive errors well being and blood sugar levels of children and adolscents with type 1 diabet. Iran J Diabetes Lipid Disord. 2019;18(4):199-206.
- 16. Woodcock AJ, Julious SA, Kinmonth AL, Campbell MJ. Problems with the performance of the SF-36 among people with type 2 diabetes in general practice. Qua Life Res. 2001 Sep;10(8):661-70.
- Saifulsyahira J, Salmiah MS, Juni MH. Cost effectiveness of a health literacy education module intervention on type 2 diabetic patients. Int J Public Health Clin Sci. 2018 Aug 16;5(4):82-94.
- Wolff K, Cavanaugh K, Malone R, Hawk V, Gregory BP, Davis D, et al. The diabetes literacy and numeracy education toolkit (DLNET). Diabetes Educ. 2009;35(2):233-45.
- 19.Jenkinson C, Coulter A, Wright L. Short form 36 (SF36) health survey questionnaire: normative data for adults of working age. Br Med J. 1993;306(6890):1437-40.
- Levin HM, McEwan PJ. Cost-effectiveness analysis: Methods and applications. Sage; 2000 Sep 27.
- 21. Chittleborough CR, Baldock KL, Taylor AW, Phillips PJ. Health status assessed by the SF-36 along the diabetes continuum in an Australian population. Qual Life Res. 2006;15(4):687-94.
- 22.Cheng L, Sit JW, Choi Kc, Chair Sy, Li X, He XI. Effectiveness of interactive self-management interventions in individuals with poorly controlled type 2 diabetes: A meta-analysis of randomized controlled trials. Worldviews Evid Based Nurs. 2017;14(1):65-73.
- 23.Vadstrup ES, Frølich A, Perrild H, Borg E, Røder M. Health-related quality of life and self-related health in patients with type 2 diabetes: effects of group-based rehabilitation versus individual counselling. Health Qual Life Outcomes. 2011;9(1):1-8.
- 24.Kalda R, Rätsep A, Lember M. Predictors of quality of life of patients with type 2 diabetes. Patient Preference Adherence. 2008;2:21.
- 25.Bourdel-Marchasson I, Druet C, Helmer C, Eschwege E, Lecomte P, Le-Goff M, et al. Correlates of health-related quality of life in French people with type 2 diabetes. Diabetes Res Clin Pract. 2013;101(2):226-35.
- 26.Sochocki MP, Verity S, Atherton PJ, Huntington JL, Sloan JA, Embil JM, et al. Health related quality of life in patients with Charcot arthropathy of the foot and ankle. Foot Ankle Surg. 2008;14(1):11-5.
- 27. Chen S, Qian D, Burström K, Burström B. Impact of an educational intervention in primary care on fasting blood glucose levels and diabetes knowledge among patients with type 2 diabetes mellitus in rural China. Patient Educ Counsel. 2020;103(9):1767-73.
- 28. Yuan Z, Jiao N, Liu X, Liu C. The effect of web-based educational intervention on psychological status and blood glucose in newly diagnosed patients with diabetes type 2 in rural China: A protocol for randomized trial. Medicine. 2021;100(8).
- 29.Kieu TTM, Trinh HN, Pham HTK, Nguyen TB, Ng JYS. Direct nonmedical and indirect costs of diabetes and its associated complications in Vietnam: an estimation using national health insurance claims from a cross-sectional survey. BMJ Open. 2020;10(3):e032303.
- 30.Sulistyaningrum IH, Kristina SA, Mukti AG. Analysis of Chronic Disease Direct Cost at Public Primary Health Cares in Indonesia. Bangladesh J Medical Sci. 2021 Jun 18;20(4):762-7.
- 31.Saha D, Jaeger J. America's new climate economy: a comprehensive guide to the economic benefits of climate policy in the United States. World Resources Institute. 2020 Jul 28:3.