



Med J Islam Repub Iran. 2024 (21 May);38.57. https://doi.org/10.47176/mjiri.38.57

Epidemiology of Proximal Femural Fractures among the Elderly People of Almaty City

Khadisha Kashikova¹*[®], Ergali Nabiyev², Ramazan Askerov³, Zhassulan Argynbayev³, Ussama AbuJazar², Arnat Baizakov³, Nurlan Turbekov²

Received: 18 Apr 2023 Published: 21 May 2024

Abstract

Background: Proximal femoral fractures are a global epidemiological concern due to their association with mortality and morbidity in the geriatric population.

Methods: We conducted an epidemiological study using hospital registry data to assess the incidence and associated factors of proximal femur fractures among individuals aged 60 years or older living in Almaty City. Student's t-test was used to assess for betweengroup differences.

Results: The data showed that the overall frequency of fractures among the population of Almaty City aged 60 years and older between 2014 and 2019 averaged 169.6 per 100,000, with a higher rate among women (190.3) compared to men (135.8). However, in age groups up to 70 years and over 85 years, the frequency of proximal femur fractures was higher among men. From 2014 to 2019, the incidence of proximal femur fractures increased by 1.6 times. An analysis of the distribution of fracture frequency by season revealed that winter was the most dangerous period.

Conclusion: Our research suggests a need for further epidemiological studies on the incidence of proximal femur fractures in various regions, identifying risk factors, and developing targeted regional prevention programs.

Keywords: Epidemiology, Fracture, Osteoporosis, Femur, Proximal Femur, Body Mass Index

Conflicts of Interest: None declared Funding: None

*This work has been published under CC BY-NC-SA 1.0 license. Copyright© <u>Iran University of Medical Sciences</u>

Cite this article as: Kashikova Kh, Nabiyev E, Askerov R, Argynbayev Zh, AbuJazar U, Baizakov A, Turbekov N. Epidemiology of Proximal Fenural Fractures among the Elderly People of Almaty City. *Med J Islam Repub Iran*. 2024 (21 May);38:57. https://doi.org/10.47176/mjiri.38.57

Introduction

Proximal femur fractures account for 15 to 55% of all fractures (1-3). Fractures of the femoral neck are observed in 50-55% of cases, and fractures of the trochanteric region occur in 30-40% (4-8).

Every year, the number of femoral fractures atincreases worldwide, and most of the affected individuals are the elderly, especially women (9-12). In 1990, there were 1,660,000 FRP fractures worldwide; their frequency is estimated to increase in 2050 to 6,260,000 per year (13).

- ^{2.} KazNMU named S.D. Asfendiyarov
- ^{3.} Kazakhstan's Medikal University

Almsot 90% of trochanteric fractures were registered in patients with varying degrees of osteoporosis (14-16). With a decrease in bone mineral density, femoral fractures may occur even with a minor low-energy injury (17-19). On the other hand, aging of the population, which is typical for all countries of the world, including the Republic of Kazakhstan increases the necessity of a sound estimation of the rate of femoral fractures and their associated features.

Therefore, we aimed to assess the incidence of fractures

†What is "already known" in this topic:

Proximal femur fracture is a global epidemiological concern, it commonly occurs in geriatric patients and leads to high morbidity and mortality.

\rightarrow *What this article adds:*

This research examined the occurrence of Proximal femoral fractures in individuals aged 60 years and above in Almaty City from 2014 to 2019. On average, there were 169.6 fractures per 100,000 cases. The majority of them are a result of domestic trauma, with women experiencing a higher incidence rate. Between 2014 and 2019, the frequency of fractures increased by 1.6 times. This study indicates the necessity of additional epidemiological research.

Corresponding author: Dr Khadisha Kashikova, hadisha.kash@gmail.com

^{1.} Caspian University, International School of Medicine, Almaty, Kazakhstan

of the proximal femur among individuals older than 60 years who live in Almaty, Kazakhstan, and to identify probable risk factors for fractures of proximal femur in this population.

Methods

We performed an epidemiological study using the data from the trauma departments of the City Clinical Hospital No. 4, City Clinical Hospital 7, Almaty, City Department of Statistics for Almaty. All patients older thahn 60 years who were hospitalized in City Clinical Hospital No. 4 trauma departments and City Clinical Hospital 7 of Almat in the period from January 1, 2014, to December 31, 2019, with proximal femur fracture were included.

The target population of the study were the residents of the city of Almaty aged 60 years and older, who accounted for 6.5% of the city's total population in 2019.

The research was carried out at the Department of Traumatology and Orthopaedics of KazNMU named after S.D.

When considering the causes of fractures, it is essential to consider the influence of various systemic diseases, which indirectly increase the risk of developing proximal femoral fractures. Thus, the data related to these underlying conditions as well as the immediate causes of fractures were collected. Seasonal fluctuations in the incidence of fractures at each fracture site were also recorded. These were used to identify a possible relationship between the incidence of proximal femoral fractures and the season of the year.

Descriptive statistical methods were used to present the data among different groups of patients. We used simple cross-tabulations and bar charts were used to present the data on the incidence of proximal femoral fractures in different demographic groups of the population. Two-tailed ttest with a 0.05 level of significance was employed to assess for betwee-group differences.

Results

A total of 54,252 people, including 20,625 men (38.0%) and 33,627 women (62.0%) of the city population was identified as the at-risk population(above 60 years old). During the study period, there were 297 cases of proximal fractures of the femur in persons 60 years of age and older, of which 102 fractures (34.3%) were in men and 195 (65.7%) in women in Almaty City.

The frequency of proximal fractures of the femur among men and period from 2014 to 2019 is shown in Figure 1. It was noted that the incidence of proximal fractures of the femur increased by 1.6 times over four years (1.5 times in men and 1.7 times in women). The average absolute number of fractures for one year in the city in people of this age was 74.2, 25.5 of which occurred in men and 48.7 in women. The distribution of patients with proximal fractures of the femur by sex and age from 2014 to 2019 is presented in Table 1 and Table 2.

In 80.0% of patients, domestic trauma was the cause of proximal fractures of the femur. Street injury was the culprit for 16.6% of cases, 3.4% of which were attributed to traffic injuries. The distribution of low-energy injuries by causes (in %) is shown in Figure 2. According to the figure, it can be noted that more than half (56%) of patients' household injuries were caused by a fall from the height of one's



Figure 1. Relative incidence of proximal fractures of the femur in men and women aged 60 years and older depending on gender and age (per 100,000 persons of the corresponding population)

Age group	Men		Women		Total	
	Ν	Incidence (%)	Ν	Incidence (%)	Ν	Incidence (%)
60-64	22	7.3	8	2.7	30	10.1
65-69	20	6.7	17	5.7	37	12.5
70-74	17	5.7	42	27.6	59	19.9
75-79	14	5.4	51	6.8	65	21.9
80-84	17	5.7	45	16.0	62	20.9
85+	12	4.0	32	10.6	44	14.8
Total	102	34.3	195	65.7	297	100.0

Table 1. Distribution of patients with proximal fractures of the femur by sex and age from 2014 to 2019

http://mjiri.iums.ac.ir 2

Med J Islam Repub Iran. 2024 (21 May); 38:57.

Age group	Both genders	Men	Women	The growth rate of each group compared to the previous one (both genders) (Percentage)
60-64	37.7±13.4	55.0±24.6	24.7±14.3	-
65-69	108.9±30.2	130.1±53.0	95.6±36.1	288.9
70-74	206.7±47.4	123.0±61.5	252.5±65.0	189.8
75-79	304.1±66.2	172.0±85.9	371.2±89.8	147.1
80-84	708.2±157.7	606.8±270.6	750.0±192.9	232.9
85+	511.2±153.7	760.5±236.6	430.5±162.3	72.2
Total	169.6±17.7	135.8±15.8	190.3±23.7	

Table 2. The frequency of proximal fractures of the femur in different age groups by sex (per 1,000,000 persons of the corresponding population) in 2014

height and while walking; in a third (32%) of patients, a fracture.

Regarding the seasonal variation in fracture rates, fractures were registered in 98 (32.9%) cases in Winter months, which is approximately two times the rate in the Summer months (15.3%) cases. In the spring period, 86 (28.9%) cases, and in autumn, 68 (22.9%) cases of fracture were observed (Figure 3). Thus, the vast majority (84.7%) of patients were injured during the cold seasons.

The average body mass index among the included patients was 21.6 kg/m². Fractures occurred in people with a BMI within the normal range.

Considering the underlying and comorbid conditions, 65% of patients had more than three concomitant diseases, 18% had two, and 17% had one disease each. The most common were diseases of the cardiovascular system in 263 (64.7%), respiratory in 49 (12.0%), and endocrine system in 17.0% of the patients. Of the diseases of the cardiovascular system, various forms of chronic coronary artery dis-



Figure 2. Distribution of low-energy injuries by causes, %



Figure 3. Distribution of injuries by seasons of the year depending on gender

ease were the most commonly detected (in 62 cases), hypertension occurred in 68 patients, and obliterating atherosclerosis in 13 patients. The consequences of acute stroke occurred in 13% of patients, and encephalopathy occurred in 12%. Metabolic disorders which consisted of diabetes mellitus and obesity occurred in 13% and 4% of the patients respectively. Diseases of the gastrointestinal system affected 4.6% and neoplastic diseases 1.2% of patients with proximal fractures of the femur.

Discussion

It is known that the vast majority of femoral fractures occur in the elderly. Hence, it is necessary to thoroughly assess the structure and frequency of these fractures and determine their dynamics in senior citizens. The current epidemiological study focused on the frequency of femoral fractures among the population of Almaty City, specifically in individuals aged 60 years and older from 2014 to 2019.

In terms of the dynamics of proximal fractures of the femur incidence, we observed a 1.6-fold increase in the frequency of PPBP from 2014 to 2019. The analysis of agespecific injury rates during the observation period revealed a significant growth in the frequency of proximal fractures of the femur up to the age of 80-84 years. The most significant growth was observed in the age group of 65-69 years, with a growth rate of 288.9%. A similar pattern was observed among the female population. In contrast, the frequency of proximal fractures of the femur among men steadily increased with age, except for the age group of 70-74 years, where the rate of proximal fractures of the femur was somewhat lower than the group of 65-69 year-olds. This finding is consistent with a similar study conducted in Ufa, Russian Federation, in 2000-2005, which also reported a higher incidence of proximal fractures of the femur among men in some age groups (20).

The overall fracture rate among the population aged 60 and over in 2019 was 169.6 per 100,000, with a similar rate observed in women (190.3) compared to men (135.8). However, we did not find a statistically significant difference in the frequency of proximal fractures of the femur by gender (P > 0.05). The coefficients we obtained partially coincide with the data from a similar study conducted in Ufa, Russian Federation, between 2000 and 2005. In our study, the ratio of injury frequency between men and women was 1:1.4, while the same analysis reported a ratio of 1:1.1. It is important to note that the incidence of injuries in our region was higher than in Ufa (129.8 prosantimilles), which may be attributed to the fact that the Russian survey group included persons over 50 years of age, while our

http://mjiri.iums.ac.ir

Med J Islam Repub Iran. 2024 (21 May); 38:57.

study focused on individuals over 60 years of age (20).

Interestingly, a higher incidence rate was recorded among men in their sixties. However, women tend to predominate among the cases with proximal fractures of the femur in their seventies. Among people in their eighties, the frequency of injuries among men prevails again. This pattern was also consistent with the Russian study findings (20). The reason for this pattern is unclear. However, it may be evaluated from physiological and social standpoints.

The ratio of the relative incidence of fractures of this localization in men and women, according to researchers, was 3:1 in England (21), 4.5:1 in Italy (22), and 3.8:1 in Argentina (23). Many researchers have proven that the risk of developing these fractures is lower in Asian women than in Caucasian women (24).

The causes of low-energy injuries also differed between men and women. In men, the most common cause of household injuries was a fall from their own height, accounting for 35.3% of cases, followed by falls while walking (20.6%) and falls from a height greater than their own (15.7%). In contrast, women were more likely to experience fractures from falls while walking (42.6%) and falls from their own height (32.3%). These differences in the causes of low-energy injuries between men and women may be attributed to variations in physical activity levels, balance, and muscle strength and are consistent with previous studies (21).

Our data also revealed a statistical correlation between the frequency of proximal fractures of the femur and the year's seasons, with the most hazardous period being winter, and spring and autumn periods being relatively less hazardous. This finding is consistent with the general epidemiological situation for proximal fractures of the femur in many countries with a continental climate (25). Some authors attribute the frequency of fractures in winter and spring to low vitamin D3 synthesis (25), while others attribute it to decreased neuromuscular coordination and vitamin D deficiency (26). Studies conducted in Sweden (27, 28), Great Britain (29, 30), Australia (31), Italy (32), and the United States have also confirmed the seasonality of proximal fractures of the femur fluctuations (33, 34), although data from other studies are contradictory (35-39).

Many researchers who assessed patients' physical activity concluded that low physical activity in this category of patients is a risk factor for fractures (40-49). Increased physical activity of the elderly(walking, climbing stairs, housework, and gardening) is considered a protective measure against fractures (48), as active movements increase the load on the bone, which in turn increases bone mineral density. An increase in muscle mass also serves as protection against local impact (47-49).

Conclusion

The high incidence and variable pattern of proximal femoral fractures found in this study highlight the need for further epidemiological research on the incidence of these fractures in different regions. This research should focus on identifying sex-specific and age-specific risk factors for the subsequent development of targeted regional programs to prevent femoral fractures.

4 <u>http://mjiri.iums.ac.ir</u>

Med J Islam Repub Iran. 2024 (21 May); 38:57.

Conflict of Interests

The authors declare that they have no competing interests.

References

- Mittal R, Banerjee S. Proximal femoral fractures: principles of management and review of literature. J Clin Orthop Trauma. 2012 Jun 1;3(1):15-23.
- Mironov SP. Standardized research in traumatology and orthopedics. M: Novosti; 2008:88.
- Sklyanchuk ED. Stimulation of osteogenesis in the complex treatment of post-traumatic disorders of bone regeneration: abstract. dis. ... Doctor of Medical Sciences: 14.00.22. M; 2009:35.
- Krivova AV. Dynamics of the frequency of fractures of the proximal part of the femur among the population of the city of Tver from 1994 to 2004. Osteoporoz Osteopatii. 2007;1:2-5.
- Dulaev AK, Tsed AN, Bobrin MI, Refitskiĭ Iu V, Khodin AI, Dzhusoev IG, et al. Application of intramedullary osteosynthesis with blocking pins in patients with peri- and intra-articular fractures. Vestn Khir Im I I Grek. 2012;171(4):49-54.
- Gordienko AI. Application of the PFN fixator in the treatment of fractures of the trochanteric region in elderly patients. Sb Tez Dokl 8 S"ezda Travmatol Ortop Rossii. amara; 2006;1:149.
- Skoroglyadov AV. Surgical treatment of subtrochanteric fractures of the femur. Kazan Med Zh. 2006;87(5):361-363.
- Basankin IV. On the issue of intracranial pressure and decompression of the proximal part of the femur in diseases of the hip joint. Sovremennye Tekhnol Travmatol Ortop. 2006;2:327.
- Atamansky AI. Our experience in endoprosthetics in severe bilateral pathology of the hip joint. Novye Tekhnol Lech Reabil Bolnykh Patol Sustavov. Kurgan; 2004:147-151.
- Frolov AV. Osteosynthesis of trochanteric and subtrochanteric fractures of the femur at the present stage. Vestn RUDN. Ser Med. 2008;2:98-100.
- Kostyukov VV. Lechenie perelomov sheyki bedra u lits pozhilogo i starcheskogo vozrasta: dis. ...kand. med. nauk: 14.00.22. Moscow, 2005; 232 p.
- 12. Zuckerman JD. Hip fracture. N Engl J Med. 1996;334(23):1519-25.
- Balaian VD, Tishkov NV, Barabash Iu A, Kauts OA. Khirurgicheskoe lechenie psevdoartrozov dlinnykh trubchatykh kostei s ispol'zovaniem dopolnitel'nykh ochagov kosteobrazovaniia [Surgical treatment of long tubular bone pseudoarthroses using additional osteogenesis foci]. Sibirskii meditsinskii zhurnal (Irkutsk). 2009;90(7):73-6.
- 14. Langlais F, Burdin P, Ropars M, Skalli W, Belot N, Lambotte JC. Osteosintheses mini vulnerants du femur proximal: quells enjeux pour les fractures du sujet age. Bull Acad Natl Med. 2005;189:1399-1412.
- Popsuishapka O, Litvishko V, Grigoryev V, Ashukina N. Treatment of bone fragments nonunion after dyaphiseal fracture. Orthopaedics Traumatology And Prosthetics. 2014(1):34-41.
- Malinin VL. Endoprotezirovanie tazobedrennogo sustava pri oskol'chatykh perelomakh proksimal'nogo otdela bedra u patsientov pozhilogo vozrasta. Osteoporoz i osteoartroz – problema XXI: Materialy konferentsii. Moscow; 2009:111-113.
- Høgh J, Lund B, Lucht U. Trochanteric and subtrochanteric fractures. The operative results in a prospective and comparative study of Ender nailing and McLaughli Acta Orthop Scand. 1981 Dec;52(6):639-43.
- Girshin SG. Klinicheskie lektsii po neotlozhnoy travmatologii. Moscow: Izdatel'skiy dom "Azbuka", 2004; 544 p.
- 19 Lomtatidze ES, Lomtatidze VE, Potseluyko SV, Toropov EA. Analysis of the functional results of internal osteosynthesis in fractures of the proximal humerus. NN Priorov Journal of Traumatology and Orthopedics. 2003 Sep 15;10(3):62-6.
- Nurlygayanov RZ, Khafizov NK, Fayzullin AA. Chastota perelomov proksimal'nogo otdela bedrennoy kosti sredi zhiteley goroda Ufy (retrospektivnoe epidemiologicheskoe issledovanie). Osteoporoz I Osteopatii. 2009;1:7-9.
- Almigdad A, Mustafa A, Alazaydeh S, Alshawish M, Bani Mustafa M, Alfukaha H. Bone Fracture Patterns and Distributions according to Trauma Energy. Adv Orthop. 2022 Sep 9;2022:8695916.
- Mazzuoli GF, Gennari C, Passeri M, Celi FS, Acca M, Camporeale A, et al. Incidence of hip fracture: an Italian survey. Osteoporos Int. 1993;3(Suppl. 1):8-9.
- Bagur A, Mautalen C, Rubin Z. Epidemiology of hip fractures in an urban population of central Argentina. Osteoporos Int. 1994;4:332-5.

- 24. Melton LJ 3rd, Cooper C. Magnitude and Impact of Osteoporosis and Fractures. In: Marcus R, Feldman D, Kelsey J, editors. Osteoporosis. 2nd ed. San Diego: Academic Press; 2001. p. 557-67.
- Komissarov AN, Palshin GA, Rodionova SS. Frequency of proximal femur fractures among residents of Yakutsk. Osteoporoz I Osteopatiya. 2004;1:2-3.
- Wickham CA, Walsh K, Cooper C, Barker DJ, Margetts BM, Morris J, et al. Dietary calcium, physical activity, and risk of hip fracture: a prospective study. BMJ. 1989;299:889-92.
- 27. Holmberg S, Thorngren KG. Statistical analysis of femoral neck fractures based on 3053 cases. Clin Orthop. 1987;218:32-41.
- Zetterberg C, Elmerson S, Anderson G. Epidemiology of hip fractures in Göteborg, Sweden, 1940-1983. Clin Orthop. 1984;191:43-52.
- Stewart IM. Fractures of the neck of femur: incidence and implications. BMJ. 1955;1:698-701.
- Bastow MD, Rawlings J, Allison SP. Undernutrition, hypothermia, and injury in elderly women with fractured femur: an injury response to altered metabolism? Lancet. 1983;1:143-6.
- Lau EMC, Woo JW, Kwok TCY, Ho SC. The seasonality of hip fracture and its relationship with weather conditions in New South Wales. Aust J Public Health. 1995;19:76-80.
- 32. Canniggia M, Morreale P. Epidemiology of hip fractures in Siena, Italy, 1975-1985. Clin Orthop Relat Rec. 1989;238:131-8.
- 33. Jacobsen SJ, Goldberg J, Miles TP, Brody JA, Stiers W, Rimm AA. Seasonal variation in the incidence of hip fracture among white persons aged 65 years and older in the United States, 1984–1987. Am J Epidemiol. 1991;133:996-1004.
- 34. Jacobsen SJ, Sargent DJ, Atkinson EJ, O'Fallon WM, Melton LJ. Population-based study of the contribution of weather to hip fracture seasonality. Am J Epidemiol. 1995;141:79-83.
- 35. Aharonoff GB, Koval KJ, Skovron ML, Zuckerman JD. Circumstances of falls causing hip fractures in the elderly. Clin Orthop Relat Res. 1998;348:10-14.
- Gallagher JC, Melton LJ, Riggs BL, Bergstrath E. Epidemiology of fractures of the proximal femur in Rochester, Minnesota. Clin Orthop Relat Res. 1980;150:163-71.
- 37. Parker MJ, Martin S. Falls, hip fractures and the weather. Eur J Epidemiol. 1994;10(4):441-2.
- Pedrazzoni M, Girasole G, Bertoldo F, Bianchi G, Minisola S, Rossini M, et al. Seasonal variation in the incidence of hip fractures in Emilia-Romagna and Parma. Bone. 1993;14(Suppl 1):S57-63.
- Aharonoff GB, Koval KJ, Skovron ML, Zuckerman JD. Circumstances of falls causing hip fractures in the elderly. Clin Orthop Relat Res. 1998;348:10-14.
- 40. "BMI Classification". Global Database on Body Mass Index. World Health Organization. 2006.
- 41. Meier DE, Luckey MM, Wallenstein S, Lapinski RH, Catherwood B, Goldsmith SJ. Calcium, vitamin D, and parathyroid hormone status in young white and black women: association with racial differences in bone mass. J Clin Endocrinol Metab. 1991;72(3):703-10.
- 42. Brandro CM, Vieira JG. Fatoresenvolvidos no pico de massayssea. Arq Bras Endocrinol Metab. 1999;43(6):401-8.
- 43. Parker MJ, Martin S. Falls, hip fractures and the weather. Eur J Epidemiol. 1994;10(4):441-25.
- 44. Benevolenskaya LI. Rukovodstvo po osteoporozy [Guide to Osteoporosis]. Moscow: BINOM, Laboratoriya Znaniy; 2003. 524 p.
- 45. Kreiger N, Tenenhouse A, Joseph L, Mackenzie T, Poliquin S, Brown JP, et al. An epidemiological study of hip fracture in postmenopausal women. Am J Epidemiol. 1982;116:141-8.
- 46. Farmer ME, White LR, Brody JA, Bailey KR. Anthropometric indicators and hip fracture. The NHANES I epidemiologic follow-up study. J Am Geriatr Soc. 1989;37(1):9-16.
- Dunitz M. Osteoporosis: diagnosis and management. London: Martin Dunitz, 1998. p. 1-16.
- Cooper C, Barker DJ, Wickham C. Physical activity, muscle strength, and calcium intake in fracture of the proximal femur in Britain. BMJ. 1988;297:1443-6.
- Cummings SR, Nevitt MC, Browner WS, Stone K, Fox KM, Ensrud KE, et al. Risk factors for hip fracture in white women. Study of Osteoporotic Fractures Research Group. N Engl J Med. 1995;332(12):767-73.

Downloaded from mjiri.iums.ac.ir on 2024-07-18