

*Systematic Review Article* <u>http://mjiri.iums.ac.ir</u> Medical Journal of the Islamic Republic of Iran (MJIRI)

Med J Islam Repub Iran. 2025 (16 Jun);39.81. https://doi.org/10.47176/mjiri.39.81



# AI-Powered Clinical Decision Support Systems in Disease Diagnosis, Treatment Planning, and Prognosis: A Systematic Review

Marzieh Nojomi<sup>1</sup>, Ebrahim Babaee<sup>1\*®</sup>, Zahra Rampisheh<sup>1</sup>, Mahshid Roohravan Benis<sup>1</sup>, Mahdi Soheyli<sup>1</sup>, Nasibeh Rady Raz<sup>2\*</sup>®

Received: 12 Feb 2025 Published: 16 Jun 2025

# Abstract

**Background**: Artificial intelligence (AI) is transforming healthcare with applications that can surpass human performance in prevention, detection, and treatment. This systematic review aimed to collect and assess the impact and success of AI technologies across various healthcare domains.

Methods: A systematic search of major databases (including PubMed, Scopus, and ISI) was conducted for articles published up to 2023. Keywords related to AI-driven disease detection, classification, and prognosis were used. Non-English articles or those with inaccessible full texts were excluded. Data was extracted by two researchers, and the quality of selected articles was evaluated based on the strengths and limitations stated by the authors.

**Results**: In total, 123 articles were included. AI contributions were categorized into three areas. For disease detection (n=75), Coronavirus disease 2019 (COVID-19) was the most frequent topic (n=18), followed by oncology. Chest X-rays were the most common input (n=15). In disease classification (n=23), oncology (especially breast cancer) was the most researched field (n=7), primarily using breast imaging. For prediction and prevention (n=25), oncology was again the most studied category, with clinical and laboratory parameters being the most utilized input (n=12).

**Conclusion**: AI-driven clinical decision support systems (CDSS) exhibit strong diagnostic and prognostic accuracy in imaging and laboratory settings. However, many models function as "black boxes," which limits interpretability and clinician trust. Data bias and challenges in integrating AI tools into practice also persist. The findings suggest that future work should focus on explainable AI and rigorous real-world validation to safely implement these tools in healthcare.

Keywords: Artificial Intelligence, Clinical Decision Support Systems, Diagnosis, Treatment, Prognosis, Disease, Prediction

Conflicts of Interest: None declared Funding: None

\*This work has been published under CC BY-NC-SA 4.0 license. Copyright© Iran University of Medical Sciences

*Cite this article as*: xxxxx. AI-Powered Clinical Decision Support Systems in Disease Diagnosis, Treatment Planning, and Prognosis: A Systematic Review. *Med J Islam Repub Iran.* 2025 (16 Jun);39:81. https://doi.org/10.47176/mjiri.39.81

## Introduction

Artificial Intelligence (AI) is a multidisciplinary study that aims to create a machine capable of perceiving data, inferring information, reaching intelligence, wisdom, cognition, and ultimately making decisions. AI aims to assist human beings in their decision-making by preparing a framework for processing all data at the same time, and presenting logical thinking and problem-solving (1). These machines are designed to handle vast, complex tasks without limitations in time or accuracy (2).

Corresponding authors: Dr Ebrahim Babaee, babaee.e@iums.ac.ir Dr Nasibeh Rady Raz, radyraz.n@iums.ac.ir

<sup>1</sup> Preventive Medicine and Public Health Research Center, Psychosocial Health Research Institute, Department of Community and Family Medicine, School of Medicine, Iran University of Medical Sciences

<sup>2</sup> Department of Artificial Intelligence in Medicine, Faculty of Advanced Technologies in Medicine, Iran University of Medical Sciences, Tehran, Iran

According to the literature, AI is applied in a broad range of applications, technologies, and facilities, from software programs to robots. Like the other services and jobs, using AI in medical care is inevitable today (3). AI could compensate for some care delivery deficiencies, such as a lack of manpower and time-consuming tasks (2). Generally, AI is used in prevention, early detection of disease, and personalized and targeted therapy. Using a variety of computational tools, AI can process different types of data, such

↑ *What is "already known" in this topic:* Artificial intelligence (AI)-powered clinical decision support systems (CDSS) demonstrate high diagnostic and prognostic accuracy in imaging and laboratory settings. However, challenges such as the "black-box" nature of models, dataset biases, and

such as the "black-box" nature of models, dataset biases, and integration barriers into clinical workflows limit their widespread adoption and trust among healthcare providers.

### $\rightarrow$ *What this article adds:*

This systematic review provides a focused analysis of AI-powered CDSS across diagnosis, treatment, and prognosis, addressing architectural suitability (eg, CNNs for imaging, RNNs for temporal data). It critically examines limitations, proposes strategies for explainability and bias mitigation, and highlights ethical and workflow considerations for real-world deployment.

as laboratory, clinical, images, signals, et cetera, to find irregular patterns of disease, perform estimation, and prediction. The ability to work with multimodal data creates a holistic view for physicians to make quick and precise decisions.

Studies show that AI may enable better disease prevention, diagnosis, and treatment. Among the main fields of disease that use AI tools, we can mention basic interventions in the fields of cancer, neurology, cardiology, and diabetes (4-6). For instance, many studies in radiologic diagnosis of different types of lung disease find the remarkable diagnostic value of various types of AI methods (7-9). However, there is a need for research to comprehensive study and address the effect of AI applications on healthcare. Therefore, here, we systematically reviewed the evidence to find the effect of different AI methods usage on the medical interventions classified as prevention, diagnosis, and treatment.

# Methods

# Study Design

Search Strategy and Study Screening Process: This systematic review was conducted in 2024. Iran University of Medical Sciences approved the study (IR.IUMS.REC.1401.711). To select appropriate and relevant studies, an extensive electronic search was conducted. For this purpose, PubMed, ISI, Cochrane, Scopus, Embase, Science Direct, and Elsevier databases were searched. Articles published until 2023 were searched. To select studies, by applying Mesh term strategy we used keywords such as "AI powered," "AI powered," "AI- powered," "AI assisted," "AI assisted," "AI based," "AI based," "AI enabled," "AI enabled," "AI aided," "AI aided," "Machine learning powered," "Machine learning assisted," "Machine learning based," "Machine learning enabled," "Machine learning aided," "Deep learning powered," "Deep learning assisted," "Deep learning based," "Deep learning enabled," "Deep learning aided," "Neural Network Computer," "Computer based," "Computer assisted," "Computer enabled," "Computer aided," "Computer powered," "System based," "System assisted," "System enabled," "System aided," "System powered," "AI," "Deep learning," "Machine learning," "Clinical Decision Support Systems," and "Clinical Decision Support". Based on the main purpose of the study, the keywords of prevention, classification, and diagnosis (detection) were also used to search for studies. The types of included studies were intervention clinical trials, randomized controlled trials, case-control, prospective and retrospective cohort, and cross-sectional. Also, references to the selected articles were searched manually. For an extensive search, 3 researchers conducted the resource search process separately and eventually coordinated the selected studies.



Identification of new studies via databases and registers

Figure 1. Finding and screening flowchart

2 <u>http://mjiri.iums.ac.ir</u> Mod Llalam Bonub Iran 202

Med J Islam Repub Iran. 2025 (16 Jun); 39:81.

In the first search phase, 2024 articles were selected. Duplicated articles were detected by 1 researcher and supervised by a subsequent researcher using EndNote (X17) software, and 1250 articles were removed. The criteria used for duplication detection were similar in the titles, first author name, and the year of publication. Here, we focus on the open-access journals and those publicly available for the possibility of further investigation. A total of 126 articles were excluded due to full-text unavailability. The number of remaining articles after this process reached 648. Subsequently, the titles and abstracts of articles were evaluated based on inclusion criteria, and 243 articles met the inclusion criteria. Using a full-text review, 120 articles were excluded due to inappropriate content. Finally, 123 eligible studies were reviewed. The finding and screening flowchart was plotted using the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) flow diagram tool (10) and reported in Figure 1.

#### Inclusion and Exclusion Criteria

We considered all studies with any study design as eligible for inclusion if they examined PICO as a tool (Table 1) for developing a search strategy for identifying potentially relevant studies. We applied other restrictions in this review, such as studies related to the English language. Also, for quality assessment, we used evaluation criteria that are usual in artificial intelligence in medicine journals, including accuracy, precision, et cetera. We evaluate AI-based systems rather than the process of diagnosis. The articles whose full texts were not accessible were excluded.

### **Data Extraction**

A standard checklist was developed for data collection and extraction. The design of the checklist was done under the supervision of the AI expert of the project. The designed checklist included the information of the extracted articles, such as the name of authors, year of publication, the country where the study took place, the type of AI model, the type of disease, the type of data processed, the AI performance measures and the limitations that mentioned in the manuscripts. To evaluate the quality of the selected articles, the strengths and limitations expressed in the articles by the authors were used as a proxy for the quality evaluation tool of the selected studies. Data extraction was done by 2 researchers of the study under the supervision of an AI expert. Any ambiguities and disagreements were listed and discussed in a session by these 2 researchers. If any problem remained after the discussion, it was investigated and resolved by the third person in the study. In the data-extracting process, the effort was to ensure that there was no missing data.

The methodological quality and potential for bias of the selected studies, particularly those evaluating diagnostic accuracy, were assessed using the Quality Assessment of Diagnostic Accuracy Studies 2 (QUADAS-2) tool. This tool evaluates studies across 4 key domains: patient selection, index test, reference standard, and flow and timing. Each domain was assessed for risk of bias, and the first 3 domains were also evaluated for concerns regarding applicability.

# **Results**

This review included studies published until 2023. Through a systematic search of electronic databases and manual screening of references, 123 articles were identified. The findings of this review indicated that AI has made significant contributions to the field of medicine in various areas, including diagnosis, prognosis, and classification. Based on these findings, our included articles were classified into these three parts, presented in Tables 2 to 4.

# Application of Artificial Intelligence in the Detection of Diseases

Our review identified a total of 75 studies that employed AI algorithms for the early detection and diagnosis of medical conditions. The details of each study, including the AI model used, performance measurement, data types, and study limitations (if available), are shown in Table 2. These studies have covered a wide range of conditions, with respiratory infections, including COVID-19 disease, being the most commonly studied topic (in 18 out of the 75 studies).

The second most researched field in this category was oncology, with a focus on gastrointestinal and skin cancers.

AI was used in various applications for detection, ranging from the interpretation of medical images to the analysis of clinical and laboratory data. In this category, chest X-ray images were the most frequently used inputs (in 15 of the 75 studies), followed by CT scan images (in 9 of the 75 studies). It is worth noting that AI-based systems demonstrated high accuracy and efficiency in detecting abnormalities.

In this category, out of 75 studies reviewed, 37 studies (49.3%) have employed "deep learning methods," while 22 studies (29.3%) utilized "machine learning methods." This demonstrates a significant reliance on deep learning algorithms for the accurate detection and diagnosis of medical conditions.

#### Classification

In the domain of classification, our review identified 23 studies that employed AI techniques to categorize medical data into distinct classes.

Population or Problem	Intervention	Comparison	Outcome
Studies of any study design that	Any type of applied AI models	Investigating the efficacy	sensitivity, specificity, accuracy, etc.
have investigated the use of AI in		of AI models compared to	
the diagnosis, classification, and		existing standard medical	
prediction of diseases.		methods	

[ Downloaded from mjiri.iums.ac.ir on 2025-07-27

Study	AI Model	AI Mode details	Disease	Sample	Data Type	AI Performance, Measurement	Country	limitation
Abbasi, 2022 (11)	machine-learning	XGBoost random forest	COVID-19	6710	chest X-ray	F1 score: 0.90 ROC curve: 0.96	Pakistan	N/A
Abdelhamid, 2022 (12)	deep-learning	SVM transfer-learning TensorFlow` optimization algorithm: RMSprop	COVID-19	7395	chest X-ray	accuracy: 99.3 sensitivity: 99 specificity: 99 F1-Score: 99.3	online dataset	N/A
Akgün, 2021 (13)	Deep Learning	VGG19, ResNet50V2, Dense- Net121, and MobileNet	COVID-19	460	cough sound	accuracy: 86.42%	Cambridge data	N/A
Ali, 2022 (14)	convolutional neural networks		COVID-19 (Omicron virus)	915	chest X-ray	ROC: 0.9888 Sensitivity: 96.2 Accuracy: 98 Precision: 100	Egypt	N/A
Alotaibi, 2022 (15)	Deep Learning	Long-term and short-term memory CNN-bidirectional LSTM CNN-LSTM	Ischemic Stroke	48	MRI	F1 score: 75.3 Accuracy: 70.2 Precision: 68.1 F1 score: 94.2 Accuracy: 94.2 Precision: 94.5 F1 score: 93.8 Accuracy: 93.8 Precision: 96.5	online dataset	N/A
Aponte-Hao, 2021 (16)	machine learning	XGBoost model	frailty	5,466	Clinical Frailty Scale (CFS)	sensitivity: 78.14 specificity: 74.41	Canada	N/A
Babukarthik, 2022 (17)	Deep Learning	Genetic Deep Learning Con- volutional Neural Net- work (GDCNN) Particle Swarm Optimization	COVID-19	5071	chest X-ray	accuracy: 97.23 sensitivity: 98.62 specificity: 97.0 precision: 93.0	publicly availa- ble datasets	N/A
Balgetir, 2021 (18)	deep learning	VGG16, VGG19, ResNet, DenseNet, MobileNet, NasNetMobile, and NasNet[arge	multiple sclerosis	105	images showing plantar pressure distribution	accuracy: 89.23 sensitivity: 89.65 specificity: 88.88		N/A
Baz, 2022 (19)	deep learning	deep learning model dubbed Parallel Convolution Neuron Net- works (PCN2)	COVID-19	328	chest X-ray	accuracy: 99.9 F1-score: 0.99	online dataset	N/A

Study	AI Model	AI Mode details	Disease	Sample size	Data Type	AI Performance, Measure- ment	Country	limitation
Bendifallah,2022 (20)	Machine learn- ing	ML models such as Logistic Regression (LR), Random Forest (RF), Deci- sion Tree (DT), eXtreme Gradient Boosting (XGB), and hard/soft Voting Classifier are considered ensemble learning tech- niques	endometriosis	8000	features about diagno- sis, symptoms, imaging, medical treatment, fertility and surgical treat- ments, and follow-up.	sensitivity: 1 specificity: 80 F1-score: 88	online dataset	N/A
Bhargava, 2022 (21)	machine learn- ing	K-NN SRC	COVID-19	31,454	chest X-ray / CT im- ages	Accuracy: 91.70 Sensitivity: 90.69 Specificity: 88.70 Accuracy: 94.40 Sensitivity: 72.00 Sensitivity: 86.00	nine distinct da- tasets	N/A
		ANN SVM				Accuracy: 96.16 Sensitivity: 91.20 Specificity: 97.40 Accuracy: 99.14		
						Specificity: 92.86		
Bozkurt, 2020 (22)	machine learn- ing	Decision Tree, Support Vec- tor Machines, k-Nearest Neighborhood Algorithm and Ensemble classifiers	Obstructive Sleep Apnea	10	ECG	Accuracy: 85.12 Sensitivity: 85 Specificity: 86	Turkey	N/A
Chen, 2019 (23)	deep learning	receptive field block, dense up sampling convolu- tion	Prostate cancer	50	MRI	Recall: 90.82 Precision: 85.53 F1-score: 88.10	dataset from MICCAI Grand Challenge	small sample size
Chen, 2022 (24)	multiple learn- ing	deep attention-based MIL	Lung cancer	1018	Chest CT images	Recall: 87 accuracy: 80 PPV: 92 NPV: 59 AUC: 84	Lung Image Da- tabase Consortium (LIDC-IDRI)	being depended on preexisting or human expert seg- mentation
Choi, 2021 (25)	Deep learning	Deep learning-based CAD algorithm (DCAD)	thoracic abnor- malities	244	chest X-ray	AUC: 0.9112	Korea	the algorithm cov- ered only 3 major thoracic le- sions

Study	AI Model	AI Mode details	Disease	Sample size	Data Type	AI Performance, Measurement	Country	limitation
Deperlioglu, 2020 (26)	Deep learning	Autoencoder Neural Networks (AEN)	Cardiovascular dis- eases	449	heart sounds	Accuracy: 100 Sensitivity: 100 Specificity: 100	PASCAL dataset	Potential interrup- tions in wireless
				409		Accuracy: 99.8 Sensitivity: 99.65 Specificity: 99.13	PhysioNet da- taset	negative user expe- riences
Ding, 2021 (27)	convolutional neural network	algorithm named 'HRNet'	oral and maxillofacial dis- ease	912	facial images		China	Lower accuracy for abnormal group compared to normal group
Asmare, 2021 (28)	machine learning	support vector machine classi- fier	Rheumatic Heart Disease	170: 124 cases 46 con- trols	heart sound	F1-score: $96.0 \pm 0.9$ recall: $95.8 \pm 1.5$ precision: $96.2 \pm 0.6$ specificity: $96.0 \pm 0.6$	in-house col- lected data with data from a freely available public database	N/A
Padmavathi Kora, 2017 (29)	Neural Network	Particle Swarm Optimization	Myocardial Infarc- tion	1806	ECG Signals	99.3% accuracy, sensitivity of 99.97%, and specificity of 98.7%		N/A
Yuan Liu, 2020 (30)	deep learning	deep convolutional neural net- work	skin diseases	16114	Skin Image	accuracy: 0.66	USA	N/A
Zhiyong Liu, 2021 (31)	deep learning	ResNet-101 and RPN net- works	prostate cancer		ultrasound image		China	N/A
Erito Marques de Souza Filho, 2022 (32)	Machine Learn- ing	Logistic Regression (LR), KNearest- Neighbors (KNN), Classifica- tion and Regression Tree (CART), AdaBoost (AB), Gradient Boosting (GB), Extreme Gradient Boosting (XGB), Random Forests (RF) and Support Vector Ma- chine (SVM)	Depression	100	clinical-laboratory and sociodemographic data	accuracy> .85	Brazil	Low socio-eco- nomic information and data
Muhammed, M, 2021 (33)	deep learning	Alex Net is the CNN model	COVID-19		chest X-ray	accuracy of 97.97%	Nigeria	N/A

6

Study	AI Model	AI Mode details	Disease	Sample size	Data Type	AI Performance, Measurement	Country	limitation
Ju Gang Nam, 2019 (34)	deep learning	deep learning-based auto- matic detection algorithm	Malignant Pulmo- nary Nodules	600	chest X-ray	specificity of 100%	South Korean	only included ma- lignant nodules - small nodules
Alvaro D. Orjuela- Cañón, 2022 (35)	Machine Learn- ing	algorithm-in-the-loop	tuberculosis	233	smear sample	accuracy 78%, sensitivity 90% , specificity 33%	Colombia	high incidence of TB in the data set, selection bias
ChunSu Park, 2022 (36)	deep learning	N/A	bone marrow edema	73	MRI	sensitivity (79%) specificity (90%)	Korea	n/A
Hatice Catal Reis, 2022 (37)	Deep Learning	InceptionResNetV2, Incep- tionV3, Mobile Net, ResNet- 101, DenseNet-169, NASNetMobile, Efficient- NetB0 algorithms	Covid-19	2400	CT and Chest X-Ray images	96.58 % sensitivity	Turkey	a small dataset - jpeg format
Prottoy Saha, 2021 (38)	Deep Neural Network	convolutional neural net- work(, EMCNet)	COVID-19	4600	chest X-ray	98.91% accuracy	Bangladesh	misclassify some COVID-19-posi- tive cases as nega- tive
Arkaprabha Sau, 2019 (39)	Machine Learn- ing	CatBoost, Logistic Regres- sion, Naïve Bayes, Random Forest	anxiety and de- pression	470	online public data	accuracy 82.6 % and precision 84.1%	India	N/A
Vijendra Singh, 2022 (40)	Deep Neural Network	Support Vector Machine (SVM), K-Nearest Neighbor (KNN), Logistic regression, Random Forest, and Naïve Bayes classifier	Chronic Kidney Disease	N/A	lab features	accuracy 100%	usa	small data sets
Fu M, 2018 (41)	Deep Learning	Multi-layer up-sampling structure	pancreas cancer	236	CT images	Precision: 77.36 ± 17.96, Recall:79.12 ± 16.27, DSC: 76.36 ± 14.34, JAC- CARD: 63.72 ± 17.05	China	N/A
Ghazal TM, 2022 (42)	Deep Learning	AlexNet , MATLAB 2020a	Skin cancer	2400	image	Accuracy: 87.1% Sensitivity: 89.0% Specificity: 94.2% PPV: 93.2% NPV:82.5%	Malaysia UAE Pakistan	N/A

Table 2. AI in the detection and diagnosis of medical condition
---

Study	AI Model	AI Mode details	Disease	Sample	Data Type	AI Performance, Measure-	Country	limitation
				size		ment		
Gopalakrishnan T, 2022 (43)	Deep Learning	Synergic Deep Learning (SDL) method with extreme Gradi- ent Boosting (XGBoost), called SDL-XGBoost	Muscular dystro- phy		MRI images	Accuracy: 96.18 Precision:87.60 Sensitivity: 97.92% Specificity: 95.64% F-Score: 92.44 Kappa:0.1800	India KSA Saudia Arabia Yemen	N/A
Hemanth DJ, 2020 (44)	Deep Learning	CNN model with eight lay- ers: image input layer, con- volutional layer, ReLU layer, cross-channel normal- ization layer, max pooling layer, fully connected layer, softmax layer, and classifica tion layer.	diabetic retinopa- thy	400	retinal fundus images	Accuracy: 97% Precision:94% Sensitivity: 94% Specificity: 98% F-Score: 94% Gmean: 96% Recall: 94%	India Turkey	N/A
Hsu FR, 2022 (45)	Deep Learning	ShuffleNet	Biliary atresia (BA)	1976	ultrasound images	Accuracy: 90.57% Precision:85.08% Sensitivity: 67.83% Specificity: 96.76% F-Score: 75.48% AUC: 92.62%	Taiwan	the size of the US image database, the doubt of over fitting, and more requirements of test sets for verifi- cation. In addition, the overdiagnosis of BA by ShuffleNet was an issue
Hu S, 2020 (46)	Deep Learning	Portable Handheld Slit- Lamp Based on a Smartphone Camera	cataract	N/A	N/A	N/A	China	N/A
Huang C, 2022 (47)	Deep Learning	Deep Transferred Efficient- Net with SVM (DTE-SVM)	Tuberculosis	N/A	CT images	Accuracy: 94.62±1.00 Precision: 95.30±1.24 Sensitivity: 93.89±1.96 Specificity: 95.35±1.31 F-Score: 94.62±1.00	China	N/A
Hussain L, 2020 (48)	Machine learn- ing	Decision Tree (DT), Naïve Bayes (NB), SVM Gaussian, SVM RBF and SVM Polynomial	congestive heart failure	72	RR time series inter- val data	Sensitivity (93.06%), Specificity (81.82%), Accuracy (88.79%), AUC (0.95)	Saudi Arabia Pakistan	N/A

Study	AI Model	AI Mode details	Disease	Sample size	Data Type	AI Performance, Measurement	Country	limitation
Hussain W, 2019 (49)	Machine learning	a new entropy index Permuta- tion Fuzzy Entropy (PFEN)	Epileptic Seizure	21	EEG	accuracy of 98.72%, sensitivity of 98.82%, specificity of 98.63%	China Japan	N/A
Hwang EJ, 2019 (50)	Deep Learning	deep learning–based auto- matic detection algorithms (DLADs)	Active Pulmonary Tuberculosis	65548	chest X-ray	classification performance of 0.977–1.000, localization performance of 0.973–1.000, Sensitivities and specificities for classification were 94.3%– 100% and 91.1%–100%	Korea	N/A
Jia X, 2022 (51)	Machine learning	WOA-SVM model	Breast Cancer	683	clinical data of breast cancer	Accuracy: 99%	China	N/A
Jia Yj, 2022 (52)	convolutional neural network	convolutional neural network VGG16 and gradient en- hanced tree model	Lung Disease		Chest sonography	Accuracy: 0.415 F-value: 0.452 Recall:0.496	China	N/A
Jo Y,2017 (53)	Deep Learning	HoloConvNet	optical screening of anthrax spores		holographic imaging	Accuracy:96.3%	Korea	N/A
Kaiume M, 2021 (54)	Deep Learning	a software based on a deep convolutional neural network (DCNN)	Rib fracture	256	CT images	Sensitivity: 0.645 (0.586– 0.703) , PPV:0.793 (0.738– 0.848)	Japan	N/A
Khan T,2019 (55)	Deep Learning	a deep learning classifier with 8 layers	Snoring	1000	sound	Accuracy: 0.96	USA	N/A
Khurana Y, 2022 (56)	Deep Learning	ResNet-50, EfficientNetB0, VGG-16 and a custom convo- lutional neural network (CNN)	COVID-19	8000	chest CT & X-ray	Precision:99.8% Specificity: 99.8% F1-Score: 98.7% Recall: 97.6%	India	N/A
N. Toda, 2023 (57)	Machine Learn- ing	N/A	detecting pulmo- nary nodules, masses	453	chest X-ray	mean wAFROC FOM score: 93	Japan	small and designed datasets, CT not used for ground truth labeling, US FDA guideline in- stead of Japanese guidelines are used.
S. Toften, 2021 (58)	neural network	N/A	sleep apnea	40	Somnofy and pulse ox- imetry signal	Cohen's kappa: 0.81	Norway	small data

Table 2. AI in the detection and diagnosis of medical conditions

http://mjiri.iums.ac.ir Med J Islam Repub Iran. 2025 (16 Jun); 39:81.

9

Study	AI Model	AI Mode details	Disease	Sample size	Data Type	AI Performance, Measurement	Country	limitation
S. Toraman, 2020 (59)	convolutional neural network	N/A	COVID-19		chest X-ray	accuracy:97.24	Turkey	small data
S. Trajanovski, 2021 (60)	deep learning	N/A	tongue squamous cell carcinoma	N/A	hyperspectral imaging (HSI)	average dice coeffi- cient=0.891±0.053  area under the ROC-curve=0.924±0.036	China	N/A
K. E. Trinkley, 2021 (61)	Machine Learn- ing	N/A	drug-induced long QT syndrome	7510	Electronic health rec- ord	N/A	United States	more analytical ap- proaches, more types of data
L. M. Tseng, 2020 (62)	deep learning	N/A	Ventricular fibril- lation	N/A	ECG	recall: 99 accuracy: 97	Taiwan	N/A
A. S. Vatian, 2022 (63)	Deep Learning	N/A	acute myocardial infarction		ECG	accuracy:85 F-scores : 74	Russian	N/A
J. Verdu-Diaz, 2020 (64)	Machine Learn- ing	N/A	Genetic diagnosis of muscular dys- trophies	976	MRIs	accuracy: 95.7 sensitivity : 92.1 specificity: 99.4	Russian	N/A
R. Verma, 2023 (65)	Machine Learn- ing	N/A	glaucoma		fundus images	Precision: 97.2 Recall: 97.3 accuracy: 97.1	India	N/A
M. Viscaino, 2021 (66)	Deep Neural Network	based on convolutional and recurrent neural networks for video otoscopy analysis.Long Short-term Memory	Ear disorders	875	video otoscopy	accuracy: 98.15 precision : 91.94 sensitivity : 91.67 specificity : 98.96 F1-score : 91.51	Chile	N/A
M. Viscaino,, 2022 (67)	convolutional neural network	N/A	otologic diagnosis		images	accuracy: 92 sensitivity: 85 specificity: 95 precision: 86 F1-score : 85	Chile	N/A
H. Wang, 2022 (68)	Deep Learning	N/A	Brown adipose tis- sue	368	PET/CT images	average DICE coefficient (DSC): 0.9057 average Hausdorff distance: 7.2810	United States	small data
J. Wang, 2021 (69)	convolutional neural network	N/A	Congenital heart disease	1308	five-view echocardio- grams video records	accuracy: 93.9	china	small data
X. Y. Wu, 2022 (70)	Machine Learn- ing	N/A	Coronavirus dis- ease	21	CT imaging	accuracy: 59 Sensitivity: 91.2 Specificity: 18.5 false-positive rate: 81.5	China	singgle center, data type, and AI model

[ Downloaded from mjiri.iums.ac.ir on 2025-07-27 ]

# 10

Study	AI Model	AI Mode details	Disease	Sample size	Data Type	AI Performance, Measurement	Country	limitation
Y. Xiang, 2021 (71)	Deep Learning	(DenseUnet composed of en- coder module, context ex- traction module and decoder module was established)auto- matic segmentation model based on the magnetic reso- nance image	N/A	19	MRI images	accuracy: 81	China	small data
C. N. Xu, 2022 (72)	convolutional neural network	-	glioma	470	MR images	Dice coefficient: 0.81 intraclass correlation coeffi- cients: 0.75 area under the curve: 0.958		computational re- sources
L. Yan,, 2021 (73)	convolutional neural network	feature selection module based on attention mecha- nisms	cervical lesions	1400	colposcopy image	sensitivity: 74.6 accuracy: 85.5 specificity: 95.7 AUC:0.909	China	N/A
Z. H. Yao, 2021 (74) S. L. Yi, 2022 (75)	deep learning Machine Learn- ing	(ResLab ) N/A	bacterial vaginosis rectal cancer	1078<	image ultrasound images	accuracy: 82.19 accuracy: 94.66 precision: 94.7 recall: 94.65 F1 values : 94.67	china china	N/A N/A
T. Yin, 2022 (76)	Machine Learn- ing	upport vector machine (SVM)	functional dyspep- sia	745	FD patients were col- lected from two clini- cal trials	accuracy: 0.773	china	N/A
W. Zeng, 2022 (77)	convolutional neural network	N/A	Fetal head circum- ference	999	ultrasound images	mean absolute difference: 1.97( ± 1.89) Dice similarity coefficient: 97.61	china	N/A
Y. Zeng, 2020 (78)	Deep learning	N/A	colorectal cancer	26000	optical coherence to- mography (OCT)	sensitivity: 100 specificity: 99.7 AUC: 0.998	United States	small traing data
Q. Zhang, 2019 (79)	Deep learning	N/A	hyperlipidemia	446	physiological infor- mation and doctors' diagnosis results	accuracy: 91.49 sensitivity: 87.50 specificity: 93.33 precision: 87.50	china	all factors not con- sider
X. N. Zhang, 2022 (80)	Clinical decision support systems	N/A	Type 2 diabetes		State-University Part- nership Learning Net- work	N/A	United States	N/A

Table 2. AI in the detection and diagnosis of medical conditions

Study	AI Model	AI Mode details	Disease	Sample size	Data Type	AI Performance, Measurement	Country	limitation
Z. Zhang, 2022 (81)	Deep Learning	N/A	meibomian gland dysfunction	2420	Image	sensitivity: 88 specificity 81	China	N/A
C. Zhao, 2021 (82)	convolutional neural network	N/A	Melanoma	2420	mei-bography images	Jaccard index: 86.84	china	N/A
A. E. Zhdanov, 2023 (83)	machine learning	N/A	retinal dystrophy		signal	N/A	Romania	N/A
J. Zhou, 2022 (84)	N/D	N/A	espiratory function evaluation	220	spirometers signal	accuracy: 68	China	N/A
X. J. Zhou, 2022 (85)	convolutional neural network	N/A	dental	210 pa- tients with one or more car- ies and 94 patients without caries	panoramic radiographs	accuracy: 0.8272 precision: 0.8538 recall: 0.8770 F1 score: 0.8652 AUC: 0.9005	china	N/A

Study	AI Model	AI Mode details	Disease	Sample size	Data Type	AI Perfformance, Measurment	Country	limitation
Abdalla, 2020 (86)	conventional neural network	N/A	Arrhytmia		ECG		Dataset from Physionet, MIT	N/A
Alzubaidi, 2021 (87)	Deep Learning	N/A	breast cancer	400	H&E-stained breast biopsy images	Accuracy: 93.2	BACH 2018 Grand Challenge dataset	N/A
			diabetic foot ulcers	754	foot skin images	precision: 95.1 recall: 94.5 F1 score: 94.8	DFU 2 dataset	N/A
			wound types	783	wound images	precision: 88.1 recall: 84.8 F1 score: 86.4 accuracy: 87.94	a combination of Google search images and DFU 2 da- taset	N/A
Ara, 2022 (88)	Deep Learning	VGG network CNN models	Diabetic Macular Edema (DME), Choroidal Neovascu- larization (CNV), Drusen	84,000	Optical Coherence Tomography Images	Recall: 0.9990 Precision: 0.9990 F1-Score: 0.9990 Accuracy: 0.9990	publicly available da- taset	N/A
Badawy, 2021 (89)	fuzzy logic and deep learning	Eight CNN based SS models have been utilized: FCN-AlexNet, UNet, Se- gNet-VGG16, SegNet- VGG19, and DeepLabV3+(ResNet18, ResNet50, MobileNet-V2, and Xception.	breast cancer	1200	breast ultrasound im- ages	accuracy: 95.45 intersection over union: 78.70 F1- score: 68.08	MT_small_dataset.	N/A
Bajaj, 2017 (90)	NNLS classifier	CoHOG and Eig(Hess)-Co- HOG	Alcoholism	120	alcoholic EEG signals	Accuracy: 95.83 Sensitivity: 100 Specificity: 91.67	Online dataset	N/A
Basha, 2021 (91)	quasi-reflection- based learning procedure	swarm intelligence-driven convolutional neural network	brain tumor	16000	Brain MRI	accuracy: 95	Online dataset	N/A
Burlina , 2017 (92)	deep learning	transfer learning and univer- sal features derived from deep convolutional neu- ral networks (DCNN).	age-related macular degeneration	5664	fundus images	accuracy: 79.4, 81.5, 93.4 for different classifications	NIH AREDS dataset	N/A

Table 3. AI in classification of medical conditions

# Table 3. AI in classification of medical conditions

Study	AI Model	AI Mode details	Disease	Sample size	Data Type	AI Perfformance, Measurment	Country	limitation
Hua Li – 2019 (93)	deep learning	DenseNet-II neural network m	Breast cancer	2042	mammogram images	accuracy of 94.55%	China	
Eduardo Ramirez, 2019 (94)	neural networks and fuzzy sys- tems	type-1 and type-2	2-lead cardiac arrhyth- mias	N/A	electrode signals or leads	92.90% and 92.70% of classification	Mexico	N/A
Xinbo Ren,2023 (95)	deep learning	lock-SegNet (DBSegNet)	cardiovascular	22210	optical coherence to- mography (OCT) im- ages	91.81%	China	N/A
Dr. K. SRI- DHAR, 2022 (96)	Deep Neural Network	ResNet and DenseNet,CNN	Leukemic Cells	N/A	blood cell counts	accuracy of 95.59%	India	N/A
Gholami E, 2021 (97)	convolutional neural network	deep non-parametric transfer (DNPT), DeepRF	Gastric cancer	970	tongue images	Accuracy: 73.78	Iran	the size of the input network
Gite S, 2023 (98)	deep learning	U-Net ++	TB or other pulmonary lung diseases	662	X-ray images	Dice: 0.9796 Specificity: 0.9932 Mean-iou: 0.9598 Sensitivity: 0.9753 Recall: 0.9838 Precision: 0.9685 Accuracy: 0.9874	India	N/A
Y. Tashtoush (99)	convolutional neural networks	CNN that is augmented with convolutional block attention modules (CBAM)	Lung cancer	N/A	CT lung	acuracy: 83.49	United States	small data
C. M. Vasile, 2021 (100)	deep learning ensemble method	that fused two deep learning models, one based on convo- lutional neural network and the other based on transfer learning (5-CNN, VGG-19)	thyroid disorders	N/A	ultrasound images	accuracy: 97.35 specificity: 98.43 Sensitivity: 95.75 positive predictive value: 95.41 negative predictive value: 98.05		need representa- tive images, small data
W. Y. Wang, 2023 (101)	Deep Learning	PointNet++	knee arthroplasty	N/A	Image	N/A	China	N/A
T. Wongsiricho, 2018 (102)	hybrid ASSC technique, Multi-Layer Hy- brid Machine Learning Model	Decision Tree (DT) and Support Vector Machine	sleep disorder	100	polysomnographic data signal	accuracy: 0.694±0.22 in subject-specific classification and 0.942±0.02 in subject- independent classifica- tion.	Thailand	N/A

# 14

Table 3. AI in classification of medical conditions

Study	AI Model	AI Mode details	Disease	Sample size	Data Type	AI Perfformance, Measurment	Country	limitation
L. Yi, L. Zhang, 2022- (103)	multi-label soft- max loss (MLSL)	N/A	lung nodules	N/A	image	AUC: 82.78	china	N/A
J. B. Zang, 2022 (104)	deep residual net- work model	N/A	Cardiovascular disease	152	ECG	accuracy : 97.89	china	N/A
V. Zarikas, 2015 (105)	Bayesian net- works (BNs)	N/A	pulmonary infections	N/A	lab data	sensitivity: 90	Austria	N/A
J. Zech, 2018 (106)	Natural Language Modeling	bag-of-words (BOW), word embedding, and Latent Di- richlet allocation-based ap- proaches		96 303	CT reports text	AUC: 0.966 Sensitivity: 92.59 specificity : 89.67	United States	N/A
X. Zhang, 2021 (107)	decision support systems with team-based care	decision support systems with team-based care	type 2 diabetes		clinical data	online survey agree: 80	United States	N/A
Q. Zhao, 2023 (108)	deep learning	N/A			ECG	accuracy: highly con- sistent	china	N/A

Study	AI Model	AI Mode details	Disease	Sample size	Data Type	AI Perfformance, Measurment	Country	limitation
Amyar,2022 (109)	multi-task multi- scale learning framework	multilayer perceptron and convolutional neural network (CNN)	lung cancers esophageal cancers	195	PET images PET images	area under the ROC curve:77 area under the ROC curve:71	France	N/A
Burdick, 2020 (110)	machine learning	XGBoost Classifier	COVID-19	197	clinical param- eters	AUC: 0.86 Sensitivity: 0.90 Specificity: 0.58 LR+: 2.15 LR-: 0.17 DOR: 12.57	US	N/A
chicco, 2020 (111)	Machine learning	Random forests Gradient boosting SVM radial	heart failure	299	clinical param- eters	F1 score: 0.754 Accuracy: 0.585 F1 score: 0.750 Accuracy: 0.585 F1 score: 0.720 Accuracy: 0.543	Pakistan	small size of the dataset
Bailoor , 2021 (112)	computational he- modynamic mod- els	linear discriminant classifier	transcatheter aortic valve replacement	29	heart sound	accuracy: 90	United States	N/A
Shaline Jia Thean Koh1- 2022 (113)	deep learning	Inception-ResNet-v2 (Gradient Class Acti- vation Map)	Covid-19	795	PCR	Accuracy:98.13%, sensitivity:97.7% speci- ficity :99.1%.	Public open da- taset	N/A
Renu Narain, 2016 (114)	Neural Network	quantum neural net- work	Cardiovascular	689	symptoms	98.57% accuracy	India	N/A
Roberto Ne- gro,2020 (115) Nathan Orlando,	machine learning deep learning	N/A N/A	benign Thyroid Nodule prostate segmentation	402	needle aspira- tions ultrasound im-	accuracy:85% - sensitivity: 0.70; specific- ity: 0.99	Italian	criteria used for nodule classifi- cation - number of noduels
2020 (118) Uvais Qidwai, 2022 (117)	Machine Learn- ing-Adaptive Neuro-Fuzzy In- ference System	N/A	Age-related macular-de- generation	58	ages clinical exami- nation data	accuracy (>92%)	united kingdom	low sample size
Xavier Rafael- Palou, 2022 (118)	Neural Network	hierarchical generative and probabilistic net- work	Lung Nodule		Lung Nodule image	accuracy of 84%	Spain	low sample size, segmentations were generated semi-automati- cally,our method relied on a single axial slice of the tumour

# Table 4. AI in determining prognosis of medical conditions

Study	AI Model	AI Mode details	Disease	Sample size	Data Type	AI Perfformance, Measurment	Country	limitation
Göltepe Y, 2021 (119)	Machine learn- ing	RF(Random forest); k-NN(k-nearest neighbors); NB(Naïve Bayes); LR(Logistic regres- sion); DT(Decision tree); SVM(Support vec- tor machine)	lung cancer	56	Images	Z-score Accuracy :0.83	Turkey	N/A
Han X, 2021 (120)	Deep Learning	LDDMM-based Registration Net- work	pancreatic cancer	40	CT and CBCT im- ages	improved segmentation accuracy	USA	N/A
Hasimbegovic E, 2021 (121)	Machine learn- ing	ML-based ap- proaches for under- standing complex clinical decision- making processes	Severe Symptomatic Aortic Stenosis	692	The cohort registry data	Area under the receiver operator char acteristics curve: 0.91 Accuracy:92% Sensitivity: 92% Specificity: 90%	Austria	N/A
Hossain MM, 2022 (122)	Fuzzy convolu- tional neural network (fuzzy CNN)	QUANTITATIVE FEATURE EX- TRACTION MA- CHINE PSO BASED FINE-TUNED FUZZY CNN	Ultrasound Image Quality Identification	2600	Ultrasound Image	Accuracy (99.62%), Precision (99.62%), Recall (99.61%), F1-score (99.61%)	Bangladesh Saudi Arabia Australia	N/A
Jung JW, 2022 (123)	Machine learn- ing	Extreme Gradient Boosting algorithm (XGBoost) model as a machinelearning classiier	postoperative delirium following knee arthro- plasty	3980	Clinical data	AUC score was 0.82 (95% CI: 0.80 - 0.83) and the sensitivity, speciicity was 0.72 and 0.73 re- spectively.	Korea	N/A
Işik AH, 2013 (124)	Neural Network	BP-ANN-based Mobile Information Device Applet application is developed with the Java 2 Micro Edi- tion environment.	Chronic Pulmonary Disease	486	Spirometry Data	Accuracy: 98.7% , Specificity: 97.83% , Sensitivity: 97.63% , Correlation values: 0.946	Turkey	N/A

## Table 4. AI in determining prognosis of medical conditions

http://mjiri.iums.ac.ir Med J Islam Repub Iran. 2025 (16 Jun); 39:81.

17

## Table 4. AI in determining prognosis of medical conditions

Study	AI Model	AI Mode details	Disease	Sample size	Data Type	AI Perfformance, Measurment	Country	limitation
G. Sumana, 2021 (125)	Neural Networks	N/A	renal syndromes		laboratory data	expert evaluation assessment	India	N/A
J. Tang, 2023 (126)	machine learning	including Backpropa- gation artificial neural network (BP-ANN), random forest (RF), support vector ma- chine (SVM), and na- ive Bayes classifier (NBC)	Adrenocortical carcinoma	825	clinical data	5-year AUROCs=0.890, 0.847, and 0.854	China	small data
A. Vodenčarevic, 2018 (127)	machine learning	N/A	Rheumatoid Arthritis (RA), a chronic inflamma- tory disease		laboratory and pharmacy data	AUC value: 80	Netherlands	N/A
Y. J. Wang, 2022 G. T. Werneburg, 2022 (128)	Deep Learning machine learning	N/A kernel techniques	Alzheimer's Disease overactive bladder treat- ments Onabotulinumtox- inA (OBTX-A) injection and sacral neuromodula- tion (SNM).	127	dCDT images clinical data	accuracy: 85 accuracy: 95	China United States	small data N/A
C. H. Wu, 2015 (129)	neural network models fuzzy rule- based expert sys- tem	N/A	chronic kidney disease		clinical and lab data	accuracy: 88.40		N/A
I. Y. Zhang, 2023 (130)	machine learning	N/A	malignant tumor		clinical value	(for pediatric, adolescent, and young adult) average C-index = 86.8%, 85.2%, and 88.6% [average time-de- pendent AUC = 76.5%, 88.1%, and 90.0%		not available data,
C. J. Zimmermann, 2021 (131)	feedback and adapted the tool	N/A	ill older adult trauma	48	trauma clini- cians in Wis- consin, Texas, and Oregon	qualitative content analysis	United States	N/A
A. Vallée, 2022 (132)	N/A	N/A	human immunodeficiency virus (HIV)	8180	electronic med- ical record (EMR)	N/A	France	A patient with a previous HIV diagnosis was included, did not compare rates of HIV testing during the study

# 18

Table 3 presents the details of each study. These studies covered a wide range of medical conditions, with oncology (specifically breast cancer) being the most studied topic in 7 of 23 studies. Another widely researched area was cardiovascular diseases, which were addressed in 4 of 23 studies. AI algorithms demonstrated remarkable capabilities in accurately classifying patients based on various data, with breast images (ultrasound or mammograms) being the most used items.

Out of 23 articles reviewed in this category, 11 studies (47.8%) have utilized "deep learning methods." In addition, 4 studies (17.4%) have utilized "neural networks," further highlighting the versatility and effectiveness of AI in classifying medical conditions.

## **Prognosis and Prevention**

In the domain of prognosis and prevention, our review identified 25 studies that leveraged AI methodologies to predict the progression and outcomes of medical conditions, as shown in Table 4. These studies covered a wide spectrum of diseases, ranging from heart failure to neurological disorders. Oncology again was the most studied field in this category. AI-based prognostic models exhibited impressive predictive performance, enabling clinicians to anticipate patient outcomes with greater accuracy and foresight. The category's most frequently used items were clinical and laboratory parameters, including PCR, used in 12 out of 25 articles. Furthermore, AI algorithms demonstrated the ability to identify prognostic factors that might otherwise go unnoticed, thereby facilitating early intervention and risk mitigation strategies.



# **Comparing the 3 Classes**

Furthermore, we evaluated research in the 3 classes based on sample size, AI models, and data type.

Figure 2 shows the comparison from the aspect of sample size. As the result shows for the classification, the AI-based system required a larger sample size. Also, detection and diagnosis need a larger sample size than prognosis. Considering the AI models, we first investigated the AI-based model as presented in Figure 3. The most used model is DL. Then, we analyzed them in each of the 3 classes. The most used model in detection-diagnosis and classification is the deep learning model, with about 49.3% and 47.8%, respectively, and prognosis uses ML, with 56%. These results indicate that for detection and classification, deep learning is the most used model, while in prognosis, in which there is a need to find patterns among data, ML is the most used model that uses a smaller sample size than the deep learning model.

We further evaluated the classes from the aspect of data type in AI-based systems. As the results in Tables 2 to 4 indicate, in detection-diagnosis and classification, different modalities of images and signals are the 2 most used data types. More than 75% and 60% are images, and more than 15% and 26% are signals in those 2 classes. Also, the most used data type in prognosis is laboratory data, with more than 53% usage. These results show that for detection and classification, images and signals carry proper information,





Downloaded from mjiri.iums.ac.ir on 2025-07-27

Figure 2. (A) Detection and Diagnosis, (B) classification, and (C) prognosis



Figure 3. AI-based models in studies. Neural networks (NN), convolutional neural networks (CNN), and deep learning (DL) stand for 3 types of network-based ML algorithms, and ML class stands for the rest of the ML algorithms.

#### Table 5. Methodology checklist: the QUADAS-2 tool components for risk of bias assessment the selected studies (questions)

- 1.Was a consecutive or random sample of patients enrolled? (Yes / No / Unclear)
- 2.Was a case-control design avoided? (Yes / No / Unclear)
- 3.Did the study avoid inappropriate exclusions? (Yes / No / Unclear)
- 4. Could the selection of patients have introduced bias? (Risk: Low / High / Unclear)
- 5.Is there concern that the included patients do not match the review question? (Concern: Low / High / Unclear)
- 6.Were the index test results interpreted without knowledge of the results of the reference standard? (Yes / No / Unclear)
- 7.If a threshold was used, was it pre-specified? (Yes / No / Unclear)
- 8.Could the conduct or interpretation of the index test have introduced bias? (Risk: Low / High / Unclear)
- 9.Is there concern that the index test, its conduct, or interpretation differ from the review question? (Concern: Low / High / Unclear)
- 10.Is the reference standard likely to correctly classify the target condition? (Yes / No / Unclear)
- 11.Were the reference standard results interpreted without knowledge of the results of the index test? (Yes / No / Unclear)
- 12.Could the reference standard, its conduct, or its interpretation have introduced bias? (Risk: Low / High / Unclear)
- 13.Is there concern that the target condition as defined by the reference standard does not match the review question? (Concern: Low / High / Unclear)
  - 14.Was there an appropriate interval between index test(s) and reference standard? (Yes / No / Unclear)
  - 15.Did all patients receive a reference standard? (Yes / No / Unclear)
  - 16.Did patients receive the same reference standard? (Yes / No / Unclear)
  - 17.Were all patients included in the analysis? (Yes / No / Unclear)
- 18.Could the patient flow have introduced bias? (Risk: Low / High / Unclear)

while in prognosis, laboratory data analysis leads to extracting early signs of disease.

The specific signaling questions used for the methodological quality and potential for bias of the selected studiesassessment, adapted from the standard QUADAS-2 checklist, are provided in Table 5. The summary of the risk of bias and applicability judgments for the diagnostic accuracy studies included in this review (as listed in Table 2) is presented in Figure 4.

## Discussion

The information obtained from studies was analyzed from several aspects, including the applied AI models, the investigated disease, sample size, data type, and measurement criteria. Most studies used structures based on DL in the model section. The most used methods were deep learning, ML, CNN, hybrid fuzzy-based learning systems, classical neural networks (NN), decision support systems (DSS), Bayesian network (BN), and particle swarm optimization (PSO). ML algorithms consist of two classes of nonnetwork-based ML algorithms, such as random forest, and network-based ML algorithms, such as NN. Here, for precise classification, we separate each type of network-based algorithm of NN, CNN, and DL, which are mostly used in papers, and consider the rest of the ML algorithms as ML class.

From the point of view of diseases, the two top investigated diseases were COVID-19 and cancer. Also, in terms of data type, various medical data types such as images, signals, clinical data, and geographical data were used in studies. However, medical images were used as the data investigated in most of the studies.

In the sample size section, the largest data volume included 96,303 CT image samples, and the smallest included 19 MRI image samples. It shows that the proper sample size

	Risk of bias domains							Risk of bias domains					
		D1	D2	D3	D4	Overall		01 1 10	D1	D2	D3	D4	Overall
	Study 1	<u> </u>	× ×		$\overline{}$	×		Study 42			<b>.</b>		
	Study 2	<u>×</u>			<u> </u>	×		Study 43	+	+	<b>H</b>	+	+
	Study 3	<u> </u>		<u> </u>	<u> </u>	×		Study 44		•	-	+	+
	Study 4	<u> </u>		<u> </u>	<u> </u>	×		Study 45		-	•	•	
	Study 5	<b>X</b>		+	-	×		Study 46	<u>×</u>	×	<u> </u>	<u> </u>	
	Study 6	+	+	+	+	+		Study 47		•	+	<b>+</b>	<b>+</b>
	Study 7	×		-	-	×		Study 48		•	•	+	+
	Study 8	×	+	+	+			Study 49			×	×	
	Study 9	-	•	-	-	<u> </u>		Study 50		+	+	+	+
	Study 10	+	-	+	+	+		Study 51	+	+	+	+	+
	Study 11	×	-	-	-	×		Study 52	×	+	+	+	
	Study 12	×	×	+	+	×		Study 53		-	+	+	
	Study 13	+	+	+	+	+		Study 54	+	+	+	+	+
	Study 14	×	-	-	-	×		Study 55			+	-	×
	Study 15	×	+	+	+	×		Study 56	+	+	+	+	+
	Study 16	×	+	+	+	X		Study 57	×	-	+	-	×
	Study 17	×	+	+	+	X		Study 58	×	+	+	+	
	Study 18	-	-	+	-	-		Study 59	+	+	+	+	+
	Study 19	×	+	+	+	×		Study 60	×	-	+	-	×
	Study 20	+	+	+	+	+		Study 61	×	+	+	+	X
Study	Study 21	×	+	+	+	×	Study	Study 62	+	+	+	+	+
0,	Study 22	+	+	+	+	+		Study 63	+	+	+	+	+
	Study 23	×	+	-	+	×		Study 64	×	×	+	-	×
	Study 24	×	+	+	+	×		Study 65	×	×	X	+	×
	Study 25	+	-	X	+	X		Study 66	+	×	×	+	×
	Study 26		<b>(</b>	X	Ť	X		Study 67	×	×	+	+	X
	Study 27	X	+	+	+	X		Study 68	×	×	+	+	×
	Study 28	×	+	-	+	X		Study 69	×	×	+	+	×
	Study 29	+	-	+	+	<u> </u>		Study 70	×	×	+	X	×
	Study 30	x	+	<b>(</b>	X	X		Study 71	×	×	+	+	×
	Study 31	×	+	+	+	X		Study 72	×	×	+	+	×
	Study 32	×	+	+	$\overline{+}$	X		Study 73	×	×	+	×	×
	Study 33	-	×	×	+	X		Study 74	+	×	X	+	X
	Study 34	×	×	+	-	X		Study 75	×	×	+	+	X
	Study 35	+	+	+	+	+		Study 76	×	×	+	+	×
	Study 36	X	-	+	-	X		Study 77	×	×	X	X	X
	Study 37	×	×	-	-	X		Study 78	Ň	X	×	Ŧ	X
	Study 38	x	Ŧ	( <del>•</del>	Ŧ	X		Study 79	X	X	+	+	×
	Study 39	×	+	+	+	×		Study 80	Ň	X	+	+	×
	Study 40	Ŧ	•	<b>+</b>	÷	+		Study 81	X	X	×	<b>(</b>	×
	Study 41	×	X	<b>H</b>	-	×		Study 82	Ň	X	+	<b>(+</b> )	X
	Dormains: D1: Patient selection. D2: Index test. D3: Reference standard. D4: Flow & timing.				Ju	dgement High Some concerns	I		Dornains: D1: Patient se D2: Index test D3: Reference D4: Flow & tim	lection. standard. ning.		Ju G	dgement High Some concerns

M. Nojomi, et al.

Figure 4. Methodology checklist: the QUADAS-2 tool for studies of diagnostic test accuracy (listed according to table 2)

depends on the applied model. Besides, it should be noted that this sample size variation might affect results. Moreover, on the measurement criteria, accuracy, specificity, F1score, and area under the curve were among the most used criteria in studies, and the accuracy of the model was at the

top. The highest accuracy value of the models is equal to 99.9% in the diagnosis of COVID-19 using deep learning and 328 data points related to chest radiography, and also in the classification of diabetic macular edema using deep learning and 84000 tomographic images.

21

			Ris	k of bias dom	ains	
		D1	D2	D3	D4	Overall
	Study 83	<b>N</b>	<b>N</b>	<b>X</b>	+	<b>N</b>
	Study 84	×	×	×	<b>(+)</b>	×
	Study 85	×	×	•	<b>(+)</b>	×
	Study 86	×	×	+	+	
	Study 87	×	×	+	+	×
	Study 88	×	×	+	+	×
	Study 89	×	×	+	+	×
	Study 90	×	×	+	+	
	Study 91	×	×	+	+	
	Study 92	×	×	+	+	
	Study 93	×	×	+	+	×
	Study 94	×	×	+	+	×
	Study 95	×	×	×	×	×
	Study 96	+	×	+	+	×
	Study 97	×	×	+	×	×
	Study 98	×	×	+	+	X
	Study 99	×	×	×	×	X
	Study 100	+	×	+	+	X
	Study 101	×	×	+	+	X
dy	Study 102	×	×	+	+	X
Stu	Study 103	×	×	×	×	X
	Study 104	×	×	+	+	X
	Study 105	×	X	+	+	X
	Study 106	×	X	+	+	X
	Study 107	×	X	+	+	X
	Study 108	×	X	+	+	X
	Study 109	×	X	+	+	X
	Study 110	×	X	+	+	X
	Study 111	X	X	+	+	X
	Study 112	X	Ň	+	(+)	X
	Study 113	<del>•</del>	Ň	<del>•</del>	(+)	X
	Study 114	X	Ň	<del>•</del>	X	Ň
	Study 115	Ň	Ň	<del>•</del>	(+)	Ň
	Study 116	Ň	x	<del>•</del>	x	Ň
	Study 117	Ň	Ň	<b>H</b>	<b>H</b>	Ň
	Study 118	Ř	Ň	<b>A</b>	<b>A</b>	×
	Study 119	Ň	Ň	<b>A</b>	<b>H</b>	Ň
	Study 120	Ā	Ň	<b>A</b>	<b>H</b>	×
	Study 121	×	Ň		×	×
	Study 122	Ā	×	<b>—</b>	<b>A</b>	
		Domains:	otion	<b>—</b>		Judgement
		D2: Index test. D3: Reference s	tandard.			😣 High 🗭 Low
		D4: Flow & timir	) <b>0</b> .			

Figure 4. Methodology checklist: the QUADAS-2 tool for studies of diagnostic test accuracy (listed according to table 2)

# Architectural Suitability and Domain Characteristics

CNNs are particularly well-suited to high-dimensional imaging tasks because their convolutional filters and pooling layers efficiently learn and aggregate spatial features, such as edges, textures, and anatomical structures, across multiple scales. In contrast, RNNs (and LSTM variants) excel at modeling temporal dependencies in sequential data, such as time-series laboratory values or vital signs, by maintaining an internal state that captures information



Figure 4. Methodology checklist: the QUADAS-2 tool for studies of diagnostic test accuracy (listed according to table 2)

across prior time steps. The nature of the input modality further dictates model choice and hyperparameter tuning: structured clinical data (eg, tabular lab results, demographics) often employ fully connected networks or tree-based models with feature engineering and regularization parameters (eg, learning rate, tree depth) optimized for tabular distributions, whereas unstructured imaging data require convolutional architectures with appropriately chosen filter sizes, depths, and spatial dropout rates to balance representational power and overfitting risk. Finally, hybrid CNN-RNN pipelines enable multimodal integration-first extracting spatial embeddings from images via CNNs, then modeling their temporal evolution or combining them with sequential clinical measurements in RNN layers-thereby capturing both spatial and temporal patterns for more comprehensive clinical predictions.

# Limitations

A key strength of this study is its comprehensive and systematic approach, as it analyzed a wide range of studies from multiple reputable databases, ensuring a thorough review of AI applications in medical diagnosis, classification, and prognosis. Additionally, the study highlighted AI's significant contributions across different medical fields, particularly in disease detection and prediction, providing valuable insights for future research. However, a notable weakness is the lack of emphasis on the speed and efficiency of AI models, as the study primarily focuses on accuracy without discussing the practical implications of AI adoption in clinical settings. Another limitation is its exclusion of non-English studies and studies with inaccessible full texts, which may introduce selection bias and limit the generalizability of its findings. Furthermore, while the paper categorizes AI applications effectively, it does not critically assess the challenges of AI implementation, such as ethical concerns, data biases, and real-world integration hurdles. Finally, considering the interpretability and data requirements, we should explain that DL models, compared to ML models, are less interpretable and need more data. However, using explainable artificial intelligence (XAI) techniques, this problem can be solved. Also, the mentioned limitations in the studies include lack of data, having small datasets that can be solved using data augmentation

or federated learning, asymmetry of data, dependence on the data labeling process, incomplete or inaccurate data that lead to data bias, and the need for an expert's opinion in the data labeling process.

Moreover, many high-performance DL models remain "black boxes," limiting clinician trust; integrating XAI methods such as saliency mapping or SHAP can partially mitigate this but adds complexity. The reliance on Englishlanguage, publicly accessible datasets may also introduce geographic and demographic biases, and small sample sizes exacerbate overfitting; strategies like data augmentation and federated learning could improve generalizability. Practical integration into clinical workflows remains challenging due to workflow disruption, interoperability issues with electronic health records, and regulatory uncertainties, underscoring the need for clinician-AI-AI codesign. Finally, ethical and regulatory concerns, including patient privacy, accountability for AI-driven decisions, and the lack of standardized approval pathways-must be addressed to ensure safe and compliant deployment.

## **Conclusion and Future Work**

AI-based methods with a variety of approaches can be used in different areas of medicine, such as automated triage systems or real-time imaging analysis, based on the type of data and the amount of available data. DL, as the most popular strategy, has multiple processing layers in the network. They are used in many medical applications with high accuracy. In medical diagnoses, classification, and prognoses, the focus has been on the use of AI methods, which increase the accuracy, and criteria for selecting methods that increase the speed of diagnoses, classification, and prognoses were not provided. This may be a suggestion for future investigation. Also, the efficiency of AI-based methods compared with manual ones is discussed in most studies, which indicates the willingness to use AI in the medical industry. Also, with the human in the loop in medicine and medical staff duty, all AI-based applications are assistants, and the final choice is made by the human expert who uses such AI-based aid.

As a result, no work is focused on replacing clinical professionals with AI. This demonstrates the complementarity and aid of artificial intelligence in medical services. It is

23

indicated that in the future, it will be explored in which medical services AI may be fully implemented, as well as how human presence will be involved. Robotic surgery is one relevant example in this subject. Robotic surgery is one relevant example in this subject.

## **Authors' Contributions**

M.N. and E.B. conceptualized the study. N.R.R., Z.R., and M.R.B. conducted the literature search and data extraction. N.R.R. and M.S. contributed to data analysis and manuscript revision. All authors reviewed and approved the final manuscript.

## **Ethical Considerations**

This study was approved by the Ethics Committee of Iran University of Medical Sciences (IR.IUMS.REC.1401.711).

### Acknowledgment

The authors would like to thank the Iran University of Medical Sciences for their support.

### **Conflict of Interests**

The authors declare that they have no competing interests.

## References

- Xu Y, Liu X, Cao X, Huang C, Liu E, Qian S, et al. Artificial intelligence: A powerful paradigm for scientific research. Innovation. 2021;2(4):100179.
- 2. Davenport T, Kalakota R. The potential for artificial intelligence in healthcare. Future Healthc J. 2019;6(2):94-8.
- Alowais SA, Alghamdi SS, Alsuhebany N, Alqahtani T, Alshaya AI, Almohareb SN, et al. Revolutionizing healthcare: the role of artificial intelligence in clinical practice. BMC Med Educ. 2023;23(1):689.
- 4. Elter M, Schulz-Wendtland R, Wittenberg T. The prediction of breast cancer biopsy outcomes using two CAD approaches that both emphasize an intelligible decision process. Med Phys. 2007;34(11):4164-72.
- Gholami E TS, Kheirabadi M. Diagnosis of Gastric Cancer via Classification of the Tongue Images using Deep Convolutional Networks. J Inf Syst Telecommun (JIST). 2021;35(9):191-6.
- 6. Gu ZC, Huang SR, Dong L, Zhou Q, Wang J, Fu B, et al. An Adapted Neural-Fuzzy Inference System Model Using Preprocessed Balance Data to Improve the Predictive Accuracy of Warfarin Maintenance Dosing in Patients After Heart Valve Replacement. Cardiovasc Drugs Ther. 2022;36(5):879-89.
- Gite S, Mishra A, Kotecha K. Enhanced lung image segmentation using deep learning. Neural Comput Appl. 2023;35(31):22839-53.
- Y. Application of Recurrent Neural Network Algorithm in Intelligent Detection of Clinical Ultrasound Images of Human Lungs. Comput Intell Neurosci. 2022;2022:9602740.
- Bajwa J, Munir U, Nori A, Williams B. Artificial intelligence in healthcare: transforming the practice of medicine. Future Healthc J. 2021;8(2):e188-e94.
- 10. PRISMA Flow Diagram Generator 2019 [Available from: http://prisma.thetacollaborative.ca/.
- Abbasi WA, Abbas SA, Andleeb S, Bibi M, Majeed F, Jaleel A, et al. COVIDX: Computer-aided diagnosis of COVID-19 and its severity prediction with raw digital chest X-ray scans. Quant Biol. 2022;10(2):208-20.
- Abdelhamid AA, Abdelhalim E, Mohamed MA, Khalifa F. Multi-Classification of Chest X-rays for COVID-19 Diagnosis Using Deep Learning Algorithms. Appl Sci (Switzerland). 2022;12(4).
- 13. Akgün D, Kabakuş AT, Karapinar Şentürk Z, Şentürk A, Küçükkülahli E. A transfer learning-based deep learning approach for automated COVID-19 diagnosis with audio data. Turk J Electr Eng Comput Sci. 2021;29:2807-23.
- 14. Ali EM, Elnady MA, Abdella AK, Elstohy RA. Detection of omicron
- 24 <u>http://mjiri.iums.ac.ir</u> Mod Llalam Popub lug
- Med J Islam Repub Iran. 2025 (16 Jun); 39:81.

virus based on convolutional neural network. J Theor Appl Inf Technol. 2022;100(9):2871-84.

- Alotaibi NS, Alotaibi AS, Eliazer M, Srinivasulu A. Detection of Ischemic Stroke Tissue Fate from the MRI Images Using a Deep Learning Approach. Mob Inf Syst. 2022;2022.
- 16. Aponte-Hao S, Wong ST, Thandi M, Ronksley P, McBrien K, Lee J, et al. Machine learning for identification of frailty in Canadian primary care practices. Int J Popul Data Sci. 2021;6(1).
- 17. Babukarthik RG, Chandramohan D, Tripathi D, Kumar M, Sambasivam G. COVID-19 identification in chest X-ray images using intelligent multi-level classification scenario. Comput Electr Eng. 2022;104.
- Balgetir F, Bilek F, Kakakus S, Arslan-Tuncer S, Demir CF. Detection of ataxia in low disability MS patients by hybrid convolutional neural networks based on images of plantar pressure distribution. Mult Scler Relat Disord. 2021;56.
- Baz A, Baz M. Pcn2: Parallel cnn to diagnose covid-19 from radiographs and metadata. Intell Autom Soft Comput. 2022;31(2):1051-69.
- 20. Bendifallah S, Puchar A, Suisse S, Delbos L, Poilblanc M, Descamps P, et al. Machine learning algorithms as new screening approach for patients with endometriosis. Sci Rep. 2022;12(1).
- Bhargava A, Bansal A, Goyal V. Machine learning-based automatic detection of novel coronavirus (COVID-19) disease. Multimed Tools Appl. 2022;81(10):13731-50.
- 22. Bozkurt F, Ucar MK, Bozkurt MR, Bilgin C. Detection of Abnormal Respiratory Events with Single Channel ECG and Hybrid Machine Learning Model in Patients with Obstructive Sleep Apnea. IRBM. 2020;41(5):241-51.
- Chen A, Zhu L, Zang H, Ding Z, Zhan S. Computer-aided diagnosis and decision-making system for medical data analysis: A case study on prostate MR images. J Manag Sci Eng. 2019;4(4):266-78.
- Chen JH, Zeng HY, Zhang C, Shi ZW, Dekker A, Wee L, et al. Lung cancer diagnosis using deep attention-based multiple instance learning and radiomics. Med Phys. 2022;49(5):3134-43.
- Choi SY, Park S, Kim M, Park J, Choi YR, Jin KN. Evaluation of a deep learning-based computer-aided detection algorithm on chest radiographs: Case-control study. Medicine. 2021;100(16):e25663.
- 26. Deperlioglu O, Kose U, Gupta D, Khanna A, Sangaiah AK. Diagnosis of heart diseases by a secure Internet of Health Things system based on Autoencoder Deep Neural Network. Comput Commun. 2020;162:31-50.
- Ding M, Kang Y, Yuan Z, Shan X, Cai Z. Detection of facial landmarks by a convolutional neural network in patients with oral and maxillofacial disease. Int J Oral Maxillofac Surg. 2021;50(11):1443-9.
- Asmare MH, Filtjens B, Woldehanna F, Janssens L, Vanrumste B. Rheumatic heart disease screening based on phonocardiogram. Sensors. 2021;21(19).
- Kora P. ECG based Myocardial Infarction detection using Hybrid Firefly Algorithm. Comput Methods Programs Biomed. 2017;152:141-8.
- Liu Y, Jain A, Eng C, Way DH, Lee K, Bui P, et al. A deep learning system for differential diagnosis of skin diseases. Nat Med. 2020;26(6):900-8.
- 31. Liu Z, Yang C, Huang J, Liu S, Zhuo Y, Lu X. Deep learning framework based on integration of S-Mask R-CNN and Inception-v3 for ultrasound image-aided diagnosis of prostate cancer. Future Gener Comput Syst. 2021;114:358-67.
- 32. de Souza Filho EM, Veiga Rey HC, Frajtag RM, Arrowsmith Cook DM, Dalbonio de Carvalho LN, Pinho Ribeiro AL, et al. Corrigendum to "Can machine learning be useful as a screening tool for depression in primary care?" J Psychiatr Res. 2022 Apr:148:61-62.
- Muhammed M, Boukar MM, Aldullahi SE, Dane S. The Application of Artificial Intelligence Technique (CNN-Alexnet) in Diagnosing COVID-19 Using Chest X-ray Images. J Res Med Dent Sci. 2021;9(5):21-6.
- 34. Nam JG, Park S, Hwang EJ, Lee JH, Jin KN, Lim KY, et al. Development and validation of deep learning-based automatic detection algorithm for malignant pulmonary nodules on chest radiographs. Radiology. 2019;290(1):218-28.
- 35. Orjuela-Canon AD, Jutinico AL, Awad C, Vergara E, Palencia A. Machine learning in the loop for tuberculosis diagnosis support. Front Public Health. 2022;10.
- 36. Park C, Kim M, Park C, Son W, Lee SM, Seok Jeong H, et al. Diagnostic performance for detecting bone marrow edema of the hip on

dual-energy CT: Deep learning model vs. musculoskeletal physicians and radiologists. Eur J Radiol. 2022;152.

- 37. Reis HC, Turk V. COVID-DSNet: A novel deep convolutional neural network for detection of coronavirus (SARS-CoV-2) cases from CT and Chest X-Ray images. Artif Intell Med. 2022;134.
- Saha P, Sadi MS, Islam MM. EMCNet: Automated COVID-19 diagnosis from X-ray images using convolutional neural network and ensemble of machine learning classifiers. Informatics in Medicine Unlocked. 2021;22.
- Sau A, Bhakta I. Erratum: Screening of anxiety and depression among seafarers using machine learning technology (Informatics in Medicine Unlocked (2019) 16, (S235291481830193X), (10.1016/j.imu.2018.12.004)). Inform Med Unlocked. 2019;16.
- Singh V, Asari VK, Rajasekaran R. A Deep Neural Network for Early Detection and Prediction of Chronic Kidney Disease. Diagnostics. 2022;12(1).
- 41. Fu M, Wu W, Hong X, Liu Q, Jiang J, Ou Y, et al. Hierarchical combinatorial deep learning architecture for pancreas segmentation of medical computed tomography cancer images. BMC Syst Biol. 2018;12.
- 42. Ghazal TM, Hussain S, Khan MF, Khan MA, Said RAT, Ahmad M. Detection of Benign and Malignant Tumors in Skin Empowered with Transfer Learning. Computational Intelligence and Neuroscience. 2022;2022.
- 43. Gopalakrishnan T, Sudhakaran P, Ramya KC, Sathesh Kumar K, Al-Wesabi FN, Alohali MA, et al. An automated deep learning based muscular dystrophy detection and classificationmodel. Comput Mater Continua. 2022;71(1):305-20.
- Hemanth DJ, Deperlioglu O, Kose U. An enhanced diabetic retinopathy detection and classification approach using deep convolutional neural network. Neural Computing and Applications. 2020;32(3):707-21.
- 45. Hsu FR, Dai ST, Chou CM, Huang SY. The application of artificial intelligence to support biliary atresia screening by ultrasound images: A study based on deep learning models. PLoS One. 2022;17(10 October).
- 46. Hu S, Wu H, Luan X, Wang Z, Adu M, Wang X, et al. Portable Handheld Slit-Lamp Based on a Smartphone Camera for Cataract Screening. J Ophthalmol. 2020;2020.
- 47. Huang C, Wang W, Zhang X, Wang SH, Zhang YD. Tuberculosis Diagnosis using Deep Transferred EfficientNet. IEEE/ACM Trans Comput Biol Bioinform. 2022;Pp.
- Hussain L, Aziz W, Khan IR, Alkinani MH, Alowibdi JS. Machine learning based congestive heart failure detection using feature importance ranking of multimodal features. Math Biosci Eng. 2020;18(1):69-91.
- 49. Hussain W, Wang B, Niu Y, Gao Y, Wang X, Sun J, et al. Epileptic Seizure Detection With Permutation Fuzzy Entropy Using Robust Machine Learning Techniques. IEEE ACCESS. 2019;7:182238-58.
- Hwang EJ, Park S, Jin KN, Kim JI, Choi SY, Lee JH, et al. Development and Validation of a Deep Learning-based Automatic Detection Algorithm for Active Pulmonary Tuberculosis on Chest Radiographs. Clin Infect Dis. 2019;69(5):739-47.
- 51. Jia X, Sun XL, Zhang XA. Breast Cancer Identification Using Machine Learning. Math Probl Eng. 2022;2022.
- 52. Jia YJ. Application of Recurrent Neural Network Algorithm in Intelligent Detection of Clinical Ultrasound Images of Human Lungs. Comput Intellig Neurosci. 2022;2022.
- 53. Jo Y, Park S, Jung J, Yoon J, Joo H, Kim MH, et al. Holographic deep learning for rapid optical screening of anthrax spores. Sci Adv. 2017;3(8).
- 54. Kaiume M, Suzuki S, Yasaka K, Sugawara H, Shen Y, Katada Y, et al. Rib fracture detection in computed tomography images using deep convolutional neural networks. Medicine (Baltimore). 2021;100(20):E26024.
- 55. Khan T. A deep learning model for snoring detection and vibration notification using a smartwearable gadget. Electronics (Switzerland). 2019;8(9).
- 56. Khurana Y, Soni U. Leveraging deep learning for COVID-19 diagnosis through chest imaging. Neural Computing and Applications. 2022;34(16):14003-12.
- 57. Toda N, Hashimoto M, Iwabuchi Y, Nagasaka M, Takeshita R, Yamada M, et al. Validation of deep learning-based computer-aided detection software use for interpretation of pulmonary abnormalities on chest radiographs and examination of factors that influence readers' performance and final diagnosis. Jpn J Radiol. 2023;41(1):38-44.

- Toften S, Kjellstadli JT, Tyvold SS, Moxness MHS. A Pilot Study of Detecting Individual Sleep Apnea Events Using Noncontact Radar Technology, Pulse Oximetry, and Machine Learning. J Sens. 2021;2021.
- 59. Toraman S, Alakus TB, Turkoglu I. Convolutional capsnet: A novel artificial neural network approach to detect COVID-19 disease from Xray images using capsule networks. Chaos Solitons Fractals. 2020;140.
- 60. Trajanovski S, Shan C, Weijtmans PJC, De Koning SGB, Ruers TJM. Tongue Tumor Detection in Hyperspectral Images Using Deep Learning Semantic Segmentation. IEEE Trans Biomed Eng. 2021;68(4):1330-40.
- 61. Trinkley KE, Pell JM, Martinez DD, Maude NR, Hale G, Rosenberg MA. Assessing Prescriber Behavior with a Clinical Decision Support Tool to Prevent Drug-Induced Long QT Syndrome. Appl Clin Inform. 2021;12(1):190-7.
- 62. Tseng LM, Tseng VS. Predicting Ventricular Fibrillation through Deep Learning. IEEE Access. 2020;8:221886-96.
- 63. Vatian AS, Gusarova NF, Dobrenko NV, Zmievsky DA, Kabyshev MV, Polevaya TA, et al. Automated evaluation of ECG parameters during the COVID-19 pandemic. Sci Tech Inf Technol.Mech Opt. 2022;22(6):1166-77.
- 64. Verdu-Diaz J, Alonso-Perez J, Nunez-Peralta C, Tasca G, Vissing J, Straub V, et al. Accuracy of a machine learning muscle MRI-based tool for the diagnosis of muscular dystrophies. Neurology. 2020;94(10):E1094-E102.
- Verma R, Shrinivasan L, Hiremath B. Machine learning classifiers for detection of glaucoma. Int J Artif Intell (IAES). 2023;12(2):806-14.
- 66. Viscaino M, Maass JC, Delano PH, Cheein FA. Computer-Aided Ear Diagnosis System Based on CNN-LSTM Hybrid Learning Framework for Video Otoscopy Examination. IEEE Access. 2021;9:161292-304.
- 67. Viscaino M, Talamilla M, Maass JC, Henríquez P, Délano PH, Cheein CA, et al. Color Dependence Analysis in a CNN-Based Computer-Aided Diagnosis System for Middle and External Ear Diseases. Diagnostics. 2022;12(4).
- Wang H, Wang Z, Wang J, Li K, Geng G, Kang F, et al. ICA-Unet: An improved U-net network for brown adipose tissue segmentation. J Innov Opt Health Sci. 2022;15(3).
- 69. Wang J, Liu X, Wang F, Zheng L, Gao F, Zhang H, et al. Automated interpretation of congenital heart disease from multi-view echocardiograms. Med Image Anal. 2021;69.
- Wu XY, Ding F, Li K, Huang WC, Zhang Y, Zhu J. Analysis of the Causes of Solitary Pulmonary Nodule Misdiagnosed as Lung Cancer by Using Artificial Intelligence: A Retrospective Study at a Single Center. Diagnostics. 2022;12(9).
- 71. Xiang Y, Wu Y, Zhang X, Hu X, Liu J, Lei L, et al. Automatic segmentation of levator ani muscle in MRI images based on DenseUnet model. Journal of Third Military Medical University. 2021;43(18):1720-8.
- 72. Xu CN, Peng YY, Zhu WF, Chen ZY, Li JR, Tan WH, et al. An automated approach for predicting glioma grade and survival of LGG patients using CNN and radiomics. Front Oncol. 2022;12.
- 73. Yan L, Li S, Guo Y, Ren P, Song H, Yang J, et al. Multi-state colposcopy image fusion for cervical precancerous lesion diagnosis using BF-CNN. Biomed Signal Process Control. 2021;68.
- 74. Yao ZH, Chen W, Li C, Yang HY, He YL, Tan YS, et al. Automatic Diagnosis of Vaginal Microecological Pathological Images Based on Deep Learning. Prog Biochem Biophys. 2021;48(11):1348-57.
- 75. Yi SL, Wei YR, Luo XM, Chen D. Diagnosis of rectal cancer based on the Xception-MS network. Phys Med Biol. 2022;67(19).
- 76. Yin T, Zheng H, Ma T, Tian X, Xu J, Li Y, et al. Predicting acupuncture efficacy for functional dyspepsia based on routine clinical features: a machine learning study in the framework of predictive, preventive, and personalized medicine. EPMA J. 2022;13(1):137-47.
- Zeng W, Luo J, Cheng JR, Lu YL. Efficient fetal ultrasound image segmentation for automatic head circumference measurement using a lightweight deep convolutional neural network. MEDICAL PHYSICS. 2022;49(8):5081-92.
- Zeng Y, Xu S, Chapman WC, Li S, Alipour Z, Abdelal H, et al. Realtime colorectal cancer diagnosis using PR-OCT with deep learning. Theranostics. 2020;10(6):2587-96.
- Zhang Q, Liu YL, Liu GH, Zhao G, Qu ZG, Yang WM. An automatic diagnostic system based on deep learning, to diagnose hyperlipidemia. Diabetes Metab Syndr. 2019;12:637-45.
- 80. Zhang XN, Svec M, Tracy R, Ozanich G. Clinical decision support systems with team-based care on type 2 diabetes improvement for

http://mjiri.iums.ac.ir

Med J Islam Repub Iran. 2025 (16 Jun); 39:81.

Medicaid patients: A quality improvement project. Int J Med Inform. 2022:158

- 81. Zhang Z, Lin X, Yu X, Fu Y, Chen X, Yang W, et al. Meibomian Gland Density: An Effective Evaluation Index of Meibomian Gland Dysfunction Based on Deep Learning and Transfer Learning J Clin Med. 2022;11(9).
- 82. Zhao C, Shuai R, Ma L, Liu WJ, Wu ML. Segmentation of dermoscopy images based on deformable 3D convolution and ResU-NeXt plus. Med Biol Eng Comput. 2021;59(9):1815-32.
- 83. Zhdanov AE, Dolganov AY, Zanca D, Borisov VI, Luchian E, Dorosinsky LG. Evaluation of the effectiveness of the decision support algorithm for physicians in retinal dystrophy using machine learning methods. Comput Opt. 2023;47(2):272-7.
- 84. Zhou J, Wang P, Guo L, Cao J, Zhou M, Dai R. Automated interpretation of the pulmonary function test by a portable spirometer in Chinese adults. Clin Respir J. 2022;16(8):555-61.
- 85. Zhou XJ, Yu GX, Yin QY, Liu Y, Zhang ZL, Sun J. Context Aware Convolutional Neural Network for Children Caries Diagnosis on Dental Panoramic Radiographs. Comput Math Methods Med. 2022;2022.
- 86. Abdalla FYO, Wu L, Ullah H, Ren G, Noor A, Mkindu H, et al. Deep convolutional neural network application to classify the ECG arrhythmia. Signal Image Video Process. 2020;14(7):1431-9.
- 87. Alzubaidi L, Fadhel MA, Al-Shamma O, Zhang JL, Santamaria J, Duan Y. Robust application of new deep learning tools: an experimental study in medical imaging. Multimed Tools Appl. 2022;81(10):13289-317
- 88. Ara RK, Matiolański A, Dziech A, Baran R, Domin P, Wieczorkiewicz A. Fast and Efficient Method for Optical Coherence Tomography Images Classification Using Deep Learning Approach. Sensors. 2022:22(13).
- 89. Badawy SM, Mohamed AEA, Hefnawy AA, Zidan HE, GadAllah MT, El-Banby GM. Automatic segmentation of breast tumors in ultrasound images based on combining fuzzy logic and deep learning-A feasibility study. PloS One. 2021;16(5).
- 90. Bajaj V, Guo YH, Sengur A, Siuly S, Alcin OF. A hybrid method based on time-frequency images for classification of alcohol and control EEG signals. Neural Comput. Appl. 2017;28(12):3717-23.
- 91. Basha J, Bacanin N, Vukobrat N, Zivkovic M, Venkatachalam K, Hubalovsky S, et al. Chaotic Harris Hawks Optimization with Quasi-Reflection-Based Learning: An Application to Enhance CNN Design. SENSORS, 2021;21(19).
- 92. Burlina P, Pacheco KD, Joshi N, Freund DE, Bressler NM. Comparing humans and deep learning performance for grading AMD: A study in using universal deep features and transfer learning for automated AMD analysis. Comput Biol Med. 2017;82:80-6.
- 93. Li H, Zhuang S, Li DA, Zhao J, Ma Y. Benign and malignant classification of mammogram images based on deep learning. Biomedical Signal Processing and Control. 2019;51:347-54.
- 94. Ramirez E, Melin P, Prado-Arechiga G. Hybrid model based on neural networks, type-1 and type-2 fuzzy systems for 2-lead cardiac arrhythmia classification. Expert Syst Appl. 2019;126:295-307.
- 95. Ren X, Zhao Y, Fan J, Wu H, Chen Q, Kubo T. Semantic segmentation of superficial layer in intracoronary optical coherence tomography based on cropping-merging and deep learning. Infrared Phys Technol. 2023:129
- 96. Sridhar K, Yeruva AR, Renjith PN, Dixit A, Jamshed A, Rastogi R. Enhanced Machine learning algorithms Lightweight Ensemble Classification of Normal versus Leukemic Cells. J Pharm Negat Results. 2022:13:496-505
- 97. Gholami E, Tabbakh SRK, Kheirabadi M. Diagnosis of Gastric Cancer via Classification of the Tongue Images using Deep Convolutional Networks. J Inf Syst Telecommun. 2021;9(35):191-6.
- 98. Gite S, Mishra A, Kotecha K. Enhanced lung image segmentation using deep learning. Neural Comput. 2022.35:22839-22853.
- 99. Tashtoush Y, Obeidat R, Al-Shorman A, Darwish O, Al-Ramahi M, Darweesh D. Enhanced convolutional neural network for non-small cell lung cancer classification. Int J Electr Comput Eng. 2023;13(1):1024-38.
- 100. Vasile CM, Udristoiu AL, Ghenea AE, Popescu M, Gheonea C, Niculescu CE, et al. Intelligent Diagnosis of Thyroid Ultrasound Imaging Using an Ensemble of Deep Learning Methods. Medicina-Lithuania, 2021:57(4).
- 101. Wang WY, Zhou HF, Yan YX, Cheng X, Yang P, Gan LZ, et al. An automatic extraction method on medical feature points based on PointNet plus plus for robot-assisted knee arthroplasty. Int J Med Robot

http://mjiri.iums.ac.ir

26 Med J Islam Repub Iran. 2025 (16 Jun); 39:81. Comput Assist Surg. 2023;19(1).

- 102. Wongsirichot T, Hanskunatai A. A multi-layer hybrid machine learning model for automatic sleep stage classification. Biomed Eng Appl Basis Commun. 2018;30(6).
- 103. Yi L, Zhang L, Xu X, Guo J. Multi-label softmax networks for pulmonary nodule classification using unbalanced and dependent categories. IEEE Trans Med Imaging. 2022; Pp.
- 104. Zang JB, Wang JL, Zhang ZD, Zheng YQ, Xue CY. Improving ECG Classification Performance by Using an Optimized One-Dimensional Residual Network Model. Appl Sci. 2022;12(24).
- 105. Zarikas V, Papageorgiou E, Regner P. Bayesian network construction using a fuzzy rule based approach for medical decision support. Expert Syst. 2015;32(3):344-69.
- 106. Zech J, Pain M, Titano J, Badgeley M, Schefflein J, Su A, et al. Natural language-based machine learning models for the annotation of clinical radiology reports. Radiology. 2018;287(2):570-80.
- 107. Zhang X, Svec M, Tracy R, Ozanich G. Clinical decision support systems with team-based care on type 2 diabetes improvement for Medicaid patients: A quality improvement project. Int J Med Inform. 2021;158:104626.
- 108. Zhao Q, Che Z, editors. Interpretation research of deep learning ECG classification results based on classification contribution value. Proc SPIE; 2023.
- 109. Amyar A, Modzelewski R, Vera P, Morard V, Ruan S. Multi-task multi-scale learning for outcome prediction in 3D PET images. Comput Biol Med. 2022:151.
- 110. Burdick H, Lam C, Mataraso S, Siefkas A, Braden G, Dellinger RP, et al. Prediction of respiratory decompensation in Covid-19 patients using machine learning: The READY trial. Comput Biol Med. 2020;124.
- 111. Chicco D, Jurman G. Machine learning can predict survival of patients with heart failure from serum creatinine and ejection fraction alone. BMC Med Inform Decis Mak. 2020;20(1).
- 112. Bailoor S, Seo JH, Schena S, Mittal R. Detecting Aortic Valve Anomaly From Induced Murmurs: Insights From Computational Hemodynamic Models. Front Physiol. 2021;12.
- 113. Jia SQ, Wang Y, Wang WZ, Zhang Q, Zhang X. Value of medical imaging artificial intelligence in the diagnosis and treatment of new coronavirus pneumonia. Expert Syst. 2022;39(3).
- 114. Narain R, Saxena S, Goyal AK. Cardiovascular risk prediction: A comparative study of framingham and quantum neural network based approach. Patient Prefer Adherence. 2016;10:1259-70.
- 115. Negro R, Rucco M, Creanza A, Mormile A, Limone PP, Garberoglio R, et al. Machine Learning Prediction of Radiofrequency Thermal Ablation Efficacy: A New Option to Optimize Thyroid Nodule Selection. Eur Thyroid J. 2020;9(4):205-12.
- 116. Orlando N, Gillies DJ, Gyacskov I, Romagnoli C, D'Souza D, Fenster A. Automatic prostate segmentation using deep learning on clinically diverse 3D transrectal ultrasound images. Med Phys. 2020;47(6):2413-26.
- 117. Qidwai U, Qidwai U, Raja M, Burton B. Smart AMD prognosis through cellphone: an innovative localized AI-based prediction system for anti-VEGF treatment prognosis in nonagenarians and centenarians. Int. Ophthalmol. 2022;42(6):1749-62.
- 118. Rafael-Palou X, Aubanell A, Ceresa M, Ribas V, Piella G, Ballester MAG. Prediction of Lung Nodule Progression with an Uncertainty-Aware Hierarchical Probabilistic Network. Diagnostics. 2022;12(11).
- 119. Göltepe Y. Performance of lung cancer prediction methods using different classification algorithms. Comput Mater Continua. 2021;67(2):2015-28.
- 120. Han X, Hong J, Reyngold M, Crane C, Cuaron J, Hajj C, et al. Deeplearning-based image registration and automatic segmentation of organs-at-risk in cone-beam CT scans from high-dose radiation treatment of pancreatic cancer. Med Phys. 2021;48(6):3084-95.
- 121. Hasimbegovic E, Papp L, Grahovac M, Krajnc D, Poschner T, Hasan W, et al. A Sneak-Peek into the Physician's Brain: A Retrospective Machine Learning-Driven Investigation of Decision-Making in TAVR versus SAVR for Young High-Risk Patients with Severe Symptomatic Aortic Stenosis, J Pers Med. 2021:11(11).
- 122. Hossain MM, Hasan MM, Rahim MA, Rahman MM, Yousuf MA, Al-Ashhab S, et al. Particle Swarm Optimized Fuzzy CNN With Quantitative Feature Fusion for Ultrasound Image Quality Identification. IEEE J Transl Eng Health Med. 2022;10:1800712
- 123. Jung JW, Hwang S, Ko S, Jo C, Park HY, Han HS, et al. A machinelearning model to predict postoperative delirium following knee

arthroplasty using electronic health records. BMC Psychiatry. 2022;22(1).

- 124. Işik AH, Güler I, Şener MU. A low-cost mobile adaptive tracking system for chronic pulmonary patients in home environment. Telemed J E Health. 2013;19(1):24-30.
- 125. Sumana G, Kalaiselvi K, Vijayalakshmi J, Shanthi S, Aparna G, Joseph MK. A design and development of support system for prediction of various renal syndromes using artificial neural networks. Int J Syst Assur Eng Manag.
- 126. Tang J, Fang Y, Xu Z. Establishment of prognostic models of adrenocortical carcinoma using machine learning and big data. Front Surg. 2023;9.
- 127. Vodenčarevic A, Van Der Goes MC, Medina OAG, De Groot MCH, Haitjema S, Van Solinge WW, et al., editors. Predicting flare probability in rheumatoid arthritis using machine learning methods. DATA 2018 -Proceedings of the 7th International Conference on Data Science, Technology and Applications; 2018.
- 128. Werneburg GT, Werneburg EA, Goldman HB, Mullhaupt AP, Vasavada SP. Machine learning provides an accurate prognostication model for refractory overactive bladder treatment response and is noninferior to human experts. Neurourol Urodyn. 2022;41(3):813-9.
- 129. Wu CH, Chiu RK, Wang SA. A cloud-based fuzzy expert system for the risk assessment of chronic kidney diseases. Int J Bus Syst Res. 2015;9(4):315-33.
- 130. Zhang IY, Hart GR, Qin B, Deng J. Long-term survival and second malignant tumor prediction in pediatric, adolescent, and young adult cancer survivors using Random Survival Forests: a SEER analysis. Sci Rep. 2023;13(1).
- 131. Zimmermann CJ, Zelenski AB, Buffington A, Baggett ND, Tucholka JL, Weis HB, et al. Best case/worst case for the trauma ICU: Development and pilot testing of a communication tool for older adults with traumatic injury. J Trauma Acute Care Surg. 2021;91(3):542-51.
- 132. Vallée A, Sveltlane D, Trichereau J, Neveu S, Fourn E, Majerholc C, et al. Electronic medical record alert increases HIV screening rates: the Foch hospital pilot POP-up project. BMC Health Serv Res. 2022;22(1).