

THE PATTERN OF MALIGNANT TUMORS IN KERMAN PROVINCE

H. TABRIZCHEE, M.D., M. MASOOMIAN, M.D., F. AHANI, M.D.,
AND M.S. ZARE, M.D.

*From the Department of Pathology, Kerman University of Medical Sciences, Kerman,
Islamic Republic of Iran.*

ABSTRACT

This investigation was done in Kerman from March 1989 to March 1993 to study the epidemiology of cancers in this province and to compare the results with a previous study performed in this area and some available statistics.

A total number of 2881 new cases of malignancy were registered and the crude relative frequency of cancers was studied. Gastric cancer with a frequency of 9.7%, malignant lymphoma with 8.8% and leukemia with 6.8% were the three most common malignancies in this province, excluding skin malignancy.

Keywords: Cancer epidemiology, Gastric carcinoma, Malignant lymphoma, Breast carcinoma.

MJIRI, Vol. 12, No. 1, 19-23, 1998.

INTRODUCTION

Epidemiologic studies have shown that the frequency and distribution of cancer varies enormously between population subgroups around the world which are defined by such variables as race, age, sex, occupation, geographic factors and life style. This epidemiologic study includes all malignant tumors of Kerman province registered during a four year period from March 1989 to March 1993.

A previous epidemiologic study of malignant tumors was also done in this region during a nine year period from March 1980 to March 1989. Some available statistics from other areas of Iran and other countries are compared and discussed in this article.

MATERIAL AND METHODS

The data used in this retrospective epidemiologic study

was collected from pathology records registered in pathology laboratories in Kerman province from March 1989 to March 1993. The laboratories included two pathology centers of Kerman Medical School, one private pathology laboratory in Kerman and one university pathology center in Rafsanjan. No other pathology center was functional during the period of this study in this province. There is a possibility that a few cases (<1%) from the neighboring provinces (Sistan, Hormozgan, Fars and Yazd) may have been included in our samples.

During this four year period, a total number of 37,366 pathologic samples were referred to these diagnostic centers in this region, of which 2881 new cases of malignancy were diagnosed. The samples were examined histologically or by the study of blood and bone marrow smears. All cases were studied and diagnosed by conventional morphologic methods. Clinical information such as sex, age and site of sampling were recorded. Cases with repeated biopsy and those which were diagnosed by cytology alone were excluded. All hematologic samples without bone marrow confirmation were also excluded.

Address for correspondence:

H. Tabrizchee, M.D., Esteghlal Street, Tabrizchee Alley, No. 44, Kerman,
I.R. Iran.

Malignant Tumors in Kerman Province

Table I: Crude relative frequency and rank order of most common cancers in Kerman province.

RO*	Both Sexes Description	Both Sexes CRF%	Males Description	Males CRF%	Females Description	Females CRF%
1	Skin	19.10	Skin	20.71	Skin	17.08
2	Stomach	9.70	Stomach	12.20	Breast	12.70
3	Malignant Lymphoma	8.80	Malignant Lymphoma	9.70	Malignant Lymphoma	7.50
4	Leukemia	6.80	Leukemia	7.00	Leukemia	6.70
5	Breast	5.70	Bladder	5.50	Stomach	6.30
6	Colorectal	4.70	Colorectal	4.30	Colorectal	5.30
7	Bladder	4.20	Lung	4.20	Cervix	4.00
8	CNS	3.70	CNS	3.90	Endometrium	3.70
9	Liver	3.20	Prostate	3.80	CNS	3.40
10	Lung	2.90	Larynx	3.70	Thyroid	3.00
11	Esophagus	2.60	Liver	3.50	Liver	2.80
12	Larynx	2.40	Bone	2.90	Bladder	2.50
13	Bone	2.30	Esophagus	2.70	Esophagus	2.40
14	Prostate	2.10	Testis	1.40	Gallbladder	2.30
15	Cervix	1.70	Oral cavity	1.10	Ovary	2.20
16	Endometrium	1.60	Small Intestine	1.00	Soft tissue	2.10
17	Thyroid	1.50	Soft tissue	1.00	Bone	1.60
18	Soft tissue	1.50	Kidney	0.90	Oral cavity	1.20
19	Gallbladder	1.30	Eye	0.80	Eye	1.20
20	Oral cavity	1.20	Gallbladder	0.60	Lung	1.10

*RO: Rank Order. CRF: Crude Relative Frequency.

RESULTS

Examination of 37,366 pathological samples resulted in the diagnosis of 2881 new cases of malignancy in Kerman province during a four year study (March 1989-March 1993). Therefore, 7.71% of the reported cases were positive for malignancy. The male:female ratio for malignant tumors was 1.35:1. Significantly high male: female ratios in non-sex organs were found in cancers of salivary glands (11: 1), larynx (5.5: 1), lung (4.9: 1), adenocarcinoma of the esophagus (4.2: 1), carcinoma of the urinary bladder (2.93: 1), carcinoma of the pancreas (2.66) and the stomach (2.57: 1).

Cancer of the thyroid (male: female ratio, 0.21: 1) and gallbladder (0.37: 1) exhibited female preponderance.

Maximum age frequency of cancer for males and females were in the seventh and sixth decades of life, respectively (Fig. 1).

Table II: Crude relative frequency and rank order of most common cancers in the pediatric age group (below 5 years old) in Kerman province.

RO*	Description	CRF%
1	Leukemia	41.40
2	Malignant Lymphoma	18.50
3	CNS Malignancy	14.30
4	Wilm's tumor	5.70
5	Testis	5.70
6	Soft Tissue	5.70
7	Eye	2.80
8	Other	5.70

*RO: Rank Order. CRF: Crude Relative Frequency.




Fig. 1. Age and sex distribution of cancer in Kerman province.

Table III: Crude relative frequency and rank order of most common cancers in the pediatric age group (below 15 years old) in Kerman province.

RO*	Description	CRF%
1	Leukemia	41.50
2	Malignant Lymphoma	21.00
3	CNS Malignancy	14.50
4	Bone	5.00
5	Soft Tissue	4.00
6	Wilm's tumor	2.50
7	Testis	2.50
8	Skin	2.50
9	Other	13.00

*RO: Rank Order. CRF: Crude Relative Frequency.

Overall, carcinoma was about five times more common than sarcoma in both sexes. The most common type of carcinoma was gastric adenocarcinoma (11.6%) and the most common type of sarcoma was malignant lymphoma (8.8%). The crude relative frequency (CRF) of the twenty most common cancers are given in Table I.

In males, the first two outstanding cancers were of the stomach (12.2%) and malignant lymphoma (9.7%) if skin cancer (20.7%) was excluded, but in females, breast cancer (12.7%) and malignant lymphoma (7.5%) ranked as the first and second most common types of cancer after skin cancer (17.08%).

In skin cancer, which ranked first in both sexes, basal cell carcinoma (67.5%) was the most common type, followed by squamous cell carcinoma (23.9%) and malignant melanoma (3.9%).

In this study, a total number of 200 cases of malignancy (6.9%) belonged to the pediatric age group (below 15 years old). The three most common malignant tumors in this group were leukemia (41.4%), malignant lymphoma (18.5%) and central nervous system tumors (14.3%) (Tables II and III).

DISCUSSION

The main aim of this work was to compare the results of this epidemiologic study with a previous one in Kerman province and some available cancer statistics from other areas. Many types of cancer showed almost the same crude relative frequency (CRF) in this study and the previous one, though other forms showed a higher or lower frequency. The most significant increase in frequency was related to CNS malignancy, liver, endometrium and lung cancers. An obvious decrease in frequency of malignancy is seen in cancers of the esophagus, urinary bladder, stomach and uterine cervix. Table IV and V compare the differences between the results of recent and previous studies.

For some differences we have a logical explanation

Table IV: The most common cancers with a higher frequency in our recent study in Kerman province.

Description	Previous Study CRF%	Recent Study CRF%	P-Value
CNS Malignancy	<1	3.70	<0.00001
Liver	1	3.20	<0.00001
Endometrium	2.65	3.70	<0.039
Lung	2.08	2.90	<0.006

CRF: Crude Relative Frequency
P-Value < 0.05.

Malignant Tumors in Kerman Province

Table V: The most common cancers with a lower frequency in our recent study in Kerman province.

Description	Previous Study CRF%	Recent Study CRF%	P-Value
Cervix	7.74	4	<0.00003
Esophagus	6.21	2.60	<0.00001
Urinary Bladder	7.01	4.20	<0.00001
Stomach	11.36	9.70	<0.022

CRF: Crude Relative Frequency
P-Value < 0.05.

while for the others we have no justifications yet. As shown in Table IV, CNS malignancy shows a marked difference of frequency in the two studies. The most probable reason for this difference could be the establishment of a neurosurgery department in this area in 1988, which has provided samples to the pathology laboratories since that time, whereas neurosurgical samples were rare in the previous study.

Advances in diagnostic methods to detect liver malignancy in this region can explain the higher frequency of liver cancer in this study. Such diagnostic methods include ultrasonography, CT scanning, MRI and aspiration needle biopsy of the liver which were not regularly available in the previous study. For endometrial adenocarcinoma which shows a 1% increase in frequency as compared with the previous study, we have no definite explanation.

The increase in frequency of lung cancer may also be due to the advances in diagnostic methods, mainly endoscopic and radiologic facilities. No reasonable explanation for the lower frequency of urinary bladder, esophageal and gastric cancer in this study can be provided at the present time.

The decrease in frequency of uterine cervix carcinoma can be due to the extensive practice of vaginal smears as a routine test for many adult females. Some cancers (salivary gland, larynx, lung and urinary bladder) show a significant male:female preponderance. Whether that is due to a higher rate of smoking habits in males or some other occupational and habitual factor remains to be answered in further studies.

When the statistics of common cancers in this region are compared with the available data from other countries (mainly from the American Cancer Society [A.C.S.] reports)², some major differences could be found. Gastric carcinoma which is the most common cancer in this region with a CRF of 9.7% shows a high frequency in China, Japan and Russia, but is not a common type of cancer in the U.S.A.² A survey among Japanese emigrants to the U.S.A. shows a major role for environmental and cultural influences.³ Whether such factors are applicable in this

region or not needs further investigation.

The frequency of malignant lymphoma (8.8%; confidence interval [C.I.] 95% = 7.8-9.8%) and leukemia (6.8%; C.I. 95% = 5.9-7%) in this study seems fairly high compared to the previous study (7% and 6.2%, respectively).

According to other studies the CRF for malignant lymphoma is 7.6³ to 8.4%⁴ in Iran, 11.7% in Lebanon,⁶ 13.5% in Iraq,⁷ 10.3% in Saudi Arabia⁸ and 4.7% in the U.S.A.²

Breast carcinoma with a CRF of 12.7% (CI[95%]=2.9-4.7%) is the most common malignancy in females in this region, excluding skin cancer. The frequency of breast cancer in American women is about 31%², while in China this is not a common malignancy.³ Most probably some cultural differences such as age of marriage, number of pregnancies and habit of breast feeding can account for such differences, although other risk factors such as hereditary factors may also influence the frequency of breast cancer. The other major difference in frequency belongs to prostatic carcinoma which shows almost a low frequency in this area and the Middle East with a CRF of 3.8% (CI [95%]= 10.9-14.3%) and 3.3%⁶, respectively, compared to the high frequency in American males (36%).² Possible explanations for such differences are as follows:

a- Prostatic carcinoma has a geographic distribution throughout the world. It is a common malignancy in the western countries while it has a lower frequency in eastern countries such as Japan.³

b- Prostate cancer is a malignancy of older men, usually over fifty years of age, which are a relatively minor age group in the general population of this country.

c- Prostatic carcinoma in our study was diagnosed on prostatectomy specimens of patients with a clinical diagnosis of benign prostatic hyperplasia and suspected cases of prostatic carcinoma according to digital rectal examinations. Such methods of detection cannot detect those cases of prostatic carcinoma which are detected in developed countries by studying serum prostatic-specific antigen (PSA) and transrectal sonography and needle biopsy.

The other major difference between our results and

American statistics pertains to lung cancer. The CRF of lung cancer in this study was 4.2% (CI [95%]= 2.9-4.7%) in males and 1.1% in females, whereas in American studies this was 14% and 13%, respectively.² Except for some differences in diagnostic methods, we have no other explanation for such a difference.

Colorectal carcinoma in Americans also shows a high frequency with a CRF of 10% and 12% in males and females, respectively.² In our study the CRF for colorectal carcinoma was 4.3% (CI [95%]= 3.5-5.3%) and 5.3% (CI [95%]= 4.1-6.3%), while in the Middle East these figures were 5.6% in males and 4.3% in females.⁶ Such a difference can be explained by environmental or mainly dietary differences.³

ACKNOWLEDGEMENT

We would like to thank the research deputy of the Kerman University of Medical Sciences for his support in this project.

REFERENCES

1. Gupta RK, Tabrizchee H: Pattern of malignant tumors in Kerman province. *Med J Islam Rep Iran* 5: 91, 1991.
2. American Cancer Society: Cancer statistics, Vol. 45, No. 1, 1995.
3. Cotran R, Kumar V, Robbins S: *Pathologic Basis of Disease*. 5 th edition, Philadelphia: W.B. Saunders Co, pp. 241-303, 1994.
4. Haghighi P, Nabizadeh I, Asvadi S, Mahallateh EA: Cancer in southern Iran. *Cancer* 27: 965-977, 1971.
5. Habibi A: Cancer in Iran, a survey of most common cases. *J National Cancer Inst* 34: 533-569, 1965.
6. Azar HA: Cancer in Lebanon and the Near East. *Cancer* 15: 66-78, 1962.
7. Majeed AMA: Cancer in Nineva province. *Ann Coll Med Mosul (Iraq)* 13: 9-19, 1982.
8. El Akkad SM, et al: Pattern of cancer in Saudi Arabia, referred to King Faisal Specialist Hospital. *Cancer* 58: 1172-8, 1986.

