

# COMPARATIVE STUDY BETWEEN EFFICACY OF METRONIDAZOLE AND FURAZOLIDONE IN 7 TO 11 YEAR OLD CHILDREN WITH GIARDIASIS

IRAJ SHARIFI, M.D.

*From the Department of Parasitology and Microbiology, Medical School, Kerman University of Medical Sciences,  
Kerman, Islamic Republic of Iran.*

## ABSTRACT

Metronidazole and furazolidone were compared for efficacy in 268 seven to 11 year old children infected with giardiasis. With metronidazole the cure rate was 82.3% and 80.0% in boys and girls respectively, while furazolidone showed an efficacy of 74.4% and 71.7% in the corresponding groups. The overall efficacy was 81.1% with metronidazole and 73.3% with furazolidone. There were no significant differences between metronidazole and furazolidone efficacies in regard to the age groups or sexes.

*MJIRI, Vol. 6, No. 2, 105-107, 1992*

## INTRODUCTION

*Giardia lamblia* is the most common intestinal parasite, distributed worldwide, particularly where sanitation is poor. It is more prevalent in children than in adults.<sup>1</sup>

Four drugs are effective in the treatment of giardiasis; quinacrine, metronidazole, furazolidone, and tinidazole.<sup>2</sup> Furazolidone and metronidazole have been routinely used in clinics and each have their advocates.<sup>3-5</sup> Comparative studies have varied considerably in their patient population, drugs tested, dose and duration of treatment, method of patient allocation, ascertainment of compliance, and criteria for cure.<sup>2,3,6,7,8</sup>

Metronidazole is used in children and adults, but in children it should be given in smaller doses.<sup>3</sup> However, some authorities have not considered furazolidone in the list of drug priority for treating giardiasis<sup>3</sup>, while others prefer furazolidone.<sup>4,5</sup>

The present study is an attempt to compare the efficacy of five-day metronidazole with 10-day furazolidone in treatment of 268 seven to 11 year old children.

## MATERIAL AND METHODS

Children aged 7 to 11 years old with laboratory-

confirmed giardiasis were enrolled after obtaining informed, oral consent from their parents. They were randomly divided into two groups. Infected children were weighed and assigned to one of two regimens. For one group (122 children), metronidazole was given orally in a dosage of 25 mg/kg/day divided into three doses for a period of 5 days.<sup>9</sup> The second group (146 children) received furazolidone 8 mg/kg/day orally divided into three or four doses for a period of 10 days.<sup>4,10</sup> Both drugs were used in tablet form with meals.

Post-treatment stool specimens were obtained at four weeks. Fresh stool specimens were concentrated by the formalin-ether solutions.<sup>11</sup> Cure was defined as failure to identify giardia in post-treatment stool examination. Treatment failures were classified as cysts found in any post-treatment specimens. Children unable to tolerate furazolidone or metronidazole were retreated with the other drug and excluded from the study groups.

Statistical analysis was by  $\chi^2$ -test. P value of <0.05 was considered significant.<sup>12</sup>

## RESULTS

The efficacy of metronidazole and furazolidone in boys, girls and both sexes are shown in Tables I, II and

## Metronidazole and Furazolidone in Giardiasis

**Table I.** The efficacy of metronidazole and furazolidone in 7 to 11 year old boys with Giardiasis

Age (years)	Metronidazole		Furazolidone	
	no.	percent	no.	percent
7-8	24	83.3	31	71.0
9-10	19	84.2	30	76.7
11	19	78.9	25	76.0
Total	62	82.3	86	74.4

III, respectively. The efficacy of metronidazole was better in both boys and girls than the corresponding sexes or age groups treated with furazolidone (Table I and II). The overall efficacy with the course of therapy was 81.1% with metronidazole and 73.3% with furazolidone (Table III). The difference between efficacies of the two drugs was not statistically significant in regards to the age groups or sexes.

### DISCUSSION

The results obtained from this study indicate that in routine medical practice, one should expect cure rates of 81.1% and 73.3% for metronidazole and furazolidone, respectively. Thus still quite a few number of patients remain untreated. Therefore, stool examination one month post-treatment should be done to make sure whether treatment was efficient.

Although the difference is not statistically significant, furazolidone, in the commonly recommended dose as used in this study, appears to be less effective than metronidazole. This finding is in accord with those found by others.<sup>8</sup> However, an exact comparison is very difficult, because each worker has used different therapeutic dosages in different age groups with completely different courses of therapy.

Furazolidone given in larger than therapeutic doses over long periods of time has been shown to be associated with an increased incidence of mammary tumors in rats.<sup>13</sup> With metronidazole administration, mammary and lung tumors occur with doses similar to those used therapeutically.<sup>14,15</sup> It is also mutagenic in bacteria<sup>16</sup> and its use in children has generally been reserved for infections that fail to respond to other drugs. Metronidazole is a more effective drug; however, it is not currently licensed for giardiasis, and there is concern over its mutagenicity.<sup>4</sup>

At this time, none of the drugs currently available for the treatment of giardiasis in children are ideal. The point of prime importance is that usage of these drugs

**Table II.** The efficacy of metronidazole and furazolidone in 7 to 11 year old girls with Giardiasis

Age (years)	Metronidazole		Furazolidone	
	no.	percent	no.	percent
7-8	19	68.4	23	69.6
9-10	20	85.0	16	75.0
11	21	85.7	21	71.4
Total	60	80.0	60	71.7

**Table III.** The efficacy of metronidazole and furazolidone in 7 to 11 year old boys and girls with Giardiasis

Age (years)	Metronidazole		Furazolidone	
	no.	percent	no.	percent
7-8	43	76.7	54	70.4
9-10	39	84.6	46	76.1
11	40	82.5	46	73.9
Total	122	81.1	146	73.3

must be carefully prescribed. Because of their carcinogenic effects, they should never be given repeatedly by physicians. A drug that is effective, nontoxic, and free of carcinogenic potential would be a welcome development in the treatment of children with giardiasis.

### ACKNOWLEDGMENT

The author gratefully acknowledges the valuable assistance of Dr. Khirolah Ghorbani, M.D. at the Department of Pediatrics and also kind cooperations of Dr. Mohsen Janghorbani, PhD., at the Department of Community Medicine, Kerman Medical School, and Dr. Iraj Norozian at the College of Veterinary Medicine, University of Tehran.

The metronidazole and furazolidone used in this study were supplied by Pharmamed Ltd. and Tehran Darou Laboratories Co. Ltd., respectively.

### REFERENCES

1. Yardley JH, Bayless TM: Giardiasis. *Gastroenterology* 52: 301-304, 1967.
2. Stevens DP: Selective primary health care; strategies for control of disease in the developing world. XIX. *Giardiasis* 7: 530-535, 1985.

## I. Sharifi, M.D.

3. WHO: The Fourth Report of the WHO Expert Committee, Containing the Sixth Model List of Essential Drugs, 6-7, 1990.
4. Plorde JJ: Giardiasis. In: Braunwald E, et al (eds), Harrison's Principles of Internal Medicine. New York, McGraw-Hill, Inc. 1: 802-805, 1991.
5. Forfar JO, Arneil GC: Textbook of Paediatrics. London, Churchill Livingstone Co, 1493, 1990.
6. Craft JC, Murphy TV, Nelson JD: Furazolidone and quinacrine: Comparativestudyof therapy for giardiasis inchildren. Am J Dis Child 135: 164-166, 1981.
7. Murphy TV, Nelson JD: Five and ten day therapy with furazolidone for giardiasis. Am J Dis Child 137: 267-270, 1983.
8. Lerman SJ, Walker RA: Treatment of giardiasis. Clin Ped 21: 409-414, 1982.
9. Mahmoud AF: Intestinal protozoa. In: Behrman RE, Vaughan VC (eds), Nelson Textbook of Pediatrics. 12th edition, Philadelphia, W.B. Saunders Co, 837, 1983.
10. Mahmoud AF, Kaplan BS: In: Behrman RE, Vaughan VC (eds), Nelson Textbook of Paediatrics. Philadelphia, W.B. Saunders Co, 725, 1987.
11. Ash LR, Orihel TC: A guide to laboratory procedures and identification. Raven Press, 12-13, 1987.
12. Armitage P: Statistical methods in medical research. Oxford, Blackwell Scientific Publications, 131, 1977.
13. Anonymous: Treatment of giardiasis (Med lett) Drug Ther 18: 39, 1976.
14. Legator MS, Connor TH, Stoeckel M: Detection of mutagenic activity of metronidazole and niridazole in body fluids of humans and mice. Science, 188: 1118, 1975.
15. Rustia M, Shubik P: Induction of lung tumors and malignant lymphomas in mice by metronidazole. JNCI 48: 721-729, 1972.
16. Voogd CE, Van der Stel JJ, Jacobs JJ: The mutagenic action of nitroimidazole: 1. Metronidazole, nimorazole, dimetridazole and ronidazole. Muta Res 26: 483-490, 1974.