

“In situ fusion for spondylolysis” is regaining its lost popularity

Hamid Behtash, MD.¹, Ebrahim Ameri, MD.², Ganjavian Mohamadsaleh, MD.³,
Farzad Omidi-Kashani, MD.⁴, Marzieh Nojomi, MD., MPH.⁵

Departments of Orthopedic Surgery, Rasul-e-Akram and Mehrad Hospitals, Tehran, Iran.

Abstract

Background: Spondylolysis and spondylolisthesis can be associated with significant low back pain, especially in physically active adolescents. Non-operative management is usually successful in improving symptoms, but surgical intervention is occasionally required. The aim of this study was to determine the effect of in situ posterolateral fusion in the treatment of refractory cases with spondylolysis.

Methods: In this prospective before and after study, we described our experience in 13 patients managed by in situ fusion after failing multimodality non-operative treatment. All surgical procedures were performed by the senior author and by a similar technique. The spondylolytic vertebra and the one below were fused, in situ. Finally, clinical outcome and recovery rates of clinical symptoms were evaluated by Henderson's functional capacity and Oswestry Disability Index version 2.1, respectively.

Results: The mean duration of non-operative management was 36 (12-72) months. There were 8 males and 5 females. Average pre- and postoperative Oswestry Disability Indices were $28.4\% \pm 13.7\%$ and 4.9 ± 7.8 respectively ($P=0.001$, significant). All patients had follow-up contact on an average of 42.3 months (range 30 - 62 months). Based on Henderson's clinical outcome functional capacity at the final follow-up stage clinical outcomes were excellent in 10, good in 2 and poor in 1 patient. The case with poor result had a pseudoarthrosis and was re-operated. Finally he had an excellent outcome.

Conclusion: We accept that the number of our cases is not high significantly but it can be claimed that in situ fusion is a safe and effective modality to treat symptomatic patients with spondylolysis and low-grade spondylolisthesis. A study with much more cases is strongly recommended.

Keywords: spondylolysis, intertransverse fusion, posterolateral approach, in situ fusion.

Introduction

Spondylolysis is characterized by a defect in the pars interarticularis of the vertebral arch. This defect is estimated to occur in 3-7% of adolescents and young adults and is often asymptomatic[1]. However, in some cases, it

can be associated with significant low back pain (LBP). This is especially true in the young athletic population which nearly 50% of cases of low back pain can be attributed to spondylolysis or spondylolisthesis. This is in contrast to patients older than 25 years which only 5% of cases of LBP are caused by spondylolysis and

1. Assistant Professor of Orthopedic and Spine Surgery, Iran University of Medical Sciences, Tehran, Iran.

2&3. Assistant Professor of Orthopedic and Spine Surgery, Iran University of Medical Sciences, Tehran, Iran.

4. **Corresponding author**, Assistant Professor of Orthopedic, Spinal fellowship, Mashhad University of Medical Sciences, Mashhad, Iran. Address: No 226, South Bozorgmehr 20, Sajad Street, Mashhad, Iran. Tel: +98915 5149248 – +98511 7646500, email: omidif@mums.ac.ir

5. Associated Professor of Community Medicine, Iran University of Medical Sciences, Tehran, Iran.

spondylolisthesis [2,3].

Anatomically the fifth lumbar vertebra is affected most often, followed by L4 and then L3. Although the exact mechanism is still unknown, pars injuries are mainly believed to be the result of repetitive microtrauma [3]. Symptomatic patients often describe LBP that often radiates into the buttocks or proximal thigh. True symptoms and signs of radiculopathy are uncommon and necessitate evaluation for disc herniation, foraminal stenosis or instability. When radicular symptoms are present, radiographic evaluation should include dynamic (flexion and extension) radiographs, and MRI should be considered [4,5].

Non-operative management should be attempted prior to operative intervention. Restriction in activity, followed by rehabilitation and abdominal strengthening exercises, are often successful in improving symptoms [6]. These treatments rarely result in bony fusion across the defect in the pars interarticularis. Recurrence of symptoms can occur with resumption of competitive activities and results in dropout of athletes as physical demands increase on the body. This dropout probably accounts for a lower incidence of spondylolysis and spondylolisthesis in professional football player when compared with high-school or college athletes [7].

The young patients (<25years) with spondylolysis or grade I spondylolisthesis (Meyerding grading system [8]) who fail non-operative management are offered either direct repair of the pars defects [9,10] or in situ posterolateral intertransverse fusion [11,12].

In this study, we have explored in situ fusion for spondylolysis with or without low-grade spondylolisthesis in affected patients. The aim of this study was to determine the effects of in situ posterolateral fusion in the treatment of refractory cases with spondylolysis.

Methods

A prospective before and after study was uti-

lized to analyze patients treated with in situ fusion. All surgical procedures were performed by the senior author (HB) from October 2002 to June 2005. All patients underwent a work up, which included a minimum of anteroposterior, lateral, dynamic (flexion-extension) radiographs, and MRI. Once operative management was indicated, a standard informed consent was obtained; patients were taken to the operating room and then placed in a standard prone position on a surgical table. Prior to induction of general anesthesia, all patients were given 1 gram intravenous cefazolin.

Surgical technique

Once the L5 level was localized through intra-operative plain radiographs, a midline longitudinal skin incision with two paraspinous fascial incisions was done. Blind dissection with the index finger could direct one to the transverse processes in the interval between multifidus and lateral (longissimus and iliocostalis) muscles. The transverse processes, pars interarticularis and facets of the levels to be fused were exposed. The spondylolytic vertebra and the one below were fused, in situ. We did not violate the facet joint at the highest transverse process exposed.

In the cases with associated disc herniation, we performed discectomy also, but in this peculiar group we performed midline longitudinal approach without intermuscular dissection for both discectomy and fusion. Iliac crest bone graft was harvested, prepared in corticocancellous strips and placed over the decorticated bony surface for fusion.

Postoperation

All patients were placed in a thoracolumbosacral orthosis for a period of 12 weeks, and the patients underwent a post-operative physical therapy prior to returning to full activity. After achieving of a solid posterolateral fusion, patients were permitted to return to their normal activities. Radiographic follow-up included dy-

| Evaluation | Functional Capacity |
|------------------|---|
| Excellent | No pain Able to return to his former occupation with no restriction Sports or recreational activities are not restricted |
| Good | Occasional pain, no more than 12 hours after extraordinary strenuous activity Able to return his occupation |
| Fair | Not restricted from engaging in less strenuous sports Less pain than preoperatively, but it remains a problem Must either wear an external support or is restricted to lighter work than before |
| Poor | Sports and recreation are restricted No better than preoperatively Unable to work Continues to seek medical help for pain |

Table 1. Henderson functional capacity.

dynamic radiographs and in doubtful cases, oblique views, Ferguson view or CT scanning were also performed to assess the presence or degree of bony fusion.

In this study, clinical outcome was evaluated by Henderson’s functional capacity [13] (Table 1). Functional capacity was evaluated as excellent, good, fair or poor. Recovery rates of clinical symptoms were calculated by Oswestry Disability Index (ODI) version 2.114 pre- and post-operatively at the time of latest follow-up. Statistical analysis was carried out with Wilcoxon-signed rank test and p value of less than 0.05 was interpreted as statistically significant.

Results

In this study, we described our experience

in 13 patients (8 males and 5 females) managed by in situ fusion after failing multimodality non-operative treatment. The mean duration of non-operative management was 36 months, ranging from 12 to 72 months. At the time of surgery patients’ age ranged from 8 to 40 years, with an average of 22.2. All patients complained of low back pain and 7 patients (53.8%) complained of thigh and/or leg pain. The pain was produced by lumbar spine extension and flexion in 6 patients (46.2%) and 2 (15.4%) respectively.

Radiographically, in addition to L5 spondylolysis, minimum spondylolisthesis with an average slip of 15.6% (range, 10 - 35%) was observed in 9 patients, and lumbar disc herniation was detected in 4 patients, which thought to be the main cause of symptoms in these patients. A




Fig. 1. Case No. 8: A. Preoperative lateral view, B. After 45 months from surgery.

| No | Patient ID | Length of stay (d) | Clinical Outcome | Post operative ODI (%) | Fusion | Residual symptoms | Follow up (months) |
|----|------------|--------------------|--|--------------------------------|---|--|--------------------|
| 1 | MM | 4 | Excellent | 2 | Yes, by plain radiography | - | 30 |
| 2 | RK | 4 | Excellent | 0 | Yes, by plain radiography | - | 43 |
| 3 | MRR | 5 | Good | 12 | Yes, by plain radiography and CT scan | Minimal LBP* | 62 |
| 4 | HRS | 4 | Good | 6 | Yes, by plain radiography and CT scan | Minimal LBP | 49 |
| 5 | AR | 4 | Excellent | 8 | Yes, by plain radiography and CT scan | - | 40 |
| 6 | SC | 9 | Excellent (poor after first operation) | 0 (28%, after first operation) | Pseudoarthrosis by plain radiography and CT scan (successfully revised) | None (after 2 nd operation) | 48 |
| 7 | RGR | 5 | Excellent | 0 | Yes, by plain radiography | - | 30 |
| 8 | DA | 3 | Excellent | 0 | Yes, by plain radiography | - | 45 |
| 9 | AD | 5 | Excellent | 4 | Yes, by plain radiography | - | 43 |
| 10 | PE | 3 | Excellent | 0 | Yes, by plain radiography | - | 40 |
| 11 | PA | 4 | Excellent | 0 | Yes, by plain radiography | - | 30 |
| 12 | MP | 4 | Excellent | 2 | Yes, by plain radiography | - | 40 |
| 13 | HS | 3 | Excellent | 2 | Yes, by plain radiography | - | 50 |

*LBP: Low Back Pain

Table 3. The patients' outcome.

low-up visit. 15.4% of patients had no symptoms at rest, but developed minor symptoms (slight LBP) with prolonged or strenuous physical activity. There were no immediate or delayed complications (including neurologic deterioration or nerve root deterioration) in any patient after the procedure. Dynamic radiographs follow-up at 24 months demonstrated solid fusion in all except 1 patient (Fig. 1). CT scanning was performed on several patients. We did not have any significant progression of the slip in our patients in the time we followed them up.

Discussion

After Kimura [15] from Japan in 1968 who

first described a method for treatment of spondylolysis by direct repair of the defect by interposing spongy bone, a variety of management strategies for patients with symptomatic spondylolysis with or without accompanying low-grade isthmic spondylolisthesis have been advocated [9,10,16-18]. The successful outcome after direct repair of the pars interarticularis ranged from 63% (by Jeanneret19) to 100% of patients (by Tokuhashi and Matsuzaki20); depend upon the client age, presence of significant disc degeneration, rigidity of the construct or the surgical technique itself [9,16,21-24]. The accuracy of plain radiographs to predict the status of a surgical union

at the pars area has been corrected in 69% of cases, especially in the presence of the instrumentation [25]. CT scanning in comparison with plain radiography reveals higher rate of surgical repair for lumbar spondylolysis [11].

Failure of direct pars repair has been attributed to the presence of significant disc degeneration or segmental instability [9,16]. Szypryt et al [26] recommended segmental fusion instead of direct pars repair in patients >25 years due to the increased incidence of disc degeneration and less satisfactory results in older patients.

Segmental spinal fusion is the traditional method for treatment of symptomatic isthmic spondylolisthesis in patients not responding to non-operative measures. It is a safe procedure with a high success rate and few complications [27]. De Loubresse and co-authors reported the outcome for 25 patients in whom low grade spondylolisthesis with radicular pain was treated with posterolateral fusion alone with an average follow-up of 32 months. Posterolateral fusion in situ provided excellent or good results in 88% of patients according to the modified classification of Stauffer and Coventry. Radicular pain disappeared at exertion and rest in 92% and 88% of the patients, respectively [28].

According to Schlenzka D [23], who compared the direct repair procedure with segmental in situ fusion for a mean follow-up of 14.8 years; although repair of the pars defect seems an appealing option for symptomatic patients who have failed to respond to conservative measures, this procedure has several disadvantages:

It is generally said direct repair requires pre-operative MRI to exclude more proximal disc degeneration (Schlenzka D [23] found no association between disc degeneration on MRI and the outcome of the patients).

It requires instrumentation, with its inheritance increased complications.

The duration of the procedure is much longer.

Lumbar spine mobility decreases after direct repair and the procedure does not seem to be ca-

pable of preventing the olisthetic disc from degeneration.

The probability of re-operation is greater in the direct repair group.

After direct repair, the ODI deteriorated with time leading to a clinically moderate but statistically significant difference in favour of segmental fusion.

Other disadvantages of the repair procedures were a lack of space for bone grafting, the difficulty of treating patients with more displacement or canal stenosis and increment in the price with the use of instrumentation [29]. With instrumentation in place, it is obvious that detecting the nonunion or solid union across the pars defect, may be more cumbersome [21].

Therefore, the expected theoretical benefits of the direct repair procedure (preservation of lumbar spine motion and protection of the adjacent segment above) could not be proven when compared to a group of patients treated by uninstrumented posterolateral fusion [23]. Lamberg TS et al [30] in the study of long-term clinical, functional and radiological outcome 21 years after standard uninstrumented posterolateral spinal fusion in childhood and adolescence spondylolysis and low-grade isthmic spondylolisthesis concluded this operation gives a satisfactory long-term fusion rate and good functional outcome and patient satisfaction. Although, the lumbar flexion is diminished, the patients perform, on average, as well as the general population in nondynamometric trunk strength measurements.

Conclusion

We accept that the number of cases is not high enough but it can be claimed that in situ fusion is a safe and effective modality to treat symptomatic patients with spondylolysis and low-grade spondylolisthesis. A trial of conservative management is always warranted, after which further studies, including dynamic radiographs, CT scanning, MRI or even pars injections can help discriminate which patients will

benefit from in situ fusion. This technique may be ineffective in patients with high grade unstable spondylolisthesis, severe degenerative disease, and L4 spondylolysis. For more potent recommendation, a study with much more cases is strongly suggested.

References

1. Fredrickson BE, Baker D, McHolick WJ, Yuan HA, Lubicky JP. The natural history of spondylolysis and spondylolisthesis. *J Bone Joint Surg Am* 1984 Jun; 66(5):699-707.
2. Micheli LJ, Wood R. Back pain in young athletes. Significant differences from adults in causes and patterns. *Arch Pediatr Adolesc Med* 1995 Jan; 149(1):15-8.
3. Motley G, Nyland J, Jacobs J, Caborn DN. The pars interarticularis stress reaction, spondylolysis and spondylolisthesis progression. *J Athl Train* 1998 Oct; 33(4):351-8.
4. Standaert CJ, Herring SA. Spondylolysis: A critical review. *Br J Sports Med* 2000 Dec; 34(6):415-22.
5. Standaert CJ, Herring SA, Halpern B, King O. Spondylolysis. *Phys Med Rehabil Clin N Am* 2000 Nov; 11(4):785-803. Review.
6. Blanda J, Bethem D, Moats W, Lew M. Defects of pars interarticularis in athletes: a protocol for nonoperative treatment. *J Spinal Disord* 1993 Oct; 6(5):406-11.
7. Shaffer B, Wiesel S, Lauerman W. Spondylolisthesis in the elite football player: an epidemiologic study in the NCAA and NFL. *J Spinal Disord* 1997 Oct; 10(5):365-70.
8. Meyerding HW. Spondylolisthesis. *Surg Gynecol Obstet* 1932; 54:371-377.
9. Dai LY, Jia LS, Yuan W, Ni B, Zhu HB. Direct repair of defect in lumbar spondylolysis and mild isthmic spondylolisthesis by bone grafting, with or without facet joint fusion. *Eur Spine J* 2001 Feb; 10(1):78-83.
10. Gillet P, Petit M. Direct repair of spondylolysis without spondylolisthesis, using a rod-screw construct and bone grafting of the pars defect. *Spine* 1999 Jun; 24(12):1252-6.
11. Backburne JS, Velikas EP. Spondylolisthesis in children and adolescents. *J Bone Joint Surg Br* 1977 Nov; 59-B(4):490-4.
12. Moller H, Hedlund R. Instrumented and noninstrumented posterolateral fusion in adult spondylolisthesis- a prospective randomized study: Part 2. *Spine* 2000 Jul; 25(13):1716-21.
13. Henderson ED: Results of the surgical treatment of spondylolisthesis. *J Bone J Surg Am* 1966 Jun; 48(4):619-42.
14. Fairbank JC, Pynsent PB. The Oswestry Disability Index. *Spine* 2000 Nov; 25(22):2940-52.
15. Kimura M. My method of filling the lesion with spongy bone in spondylolysis and spondylolisthesis. *Seikei Geka*. 1968 Mar; 19(4):285-96.
16. Lundin DA, Wiseman D, Ellenbogen RG, Shaffrey CI. Direct repair of pars interarticularis for spondylolysis and spondylolisthesis. *J Pediatr Neurosurg* 2003 Oct; 39(4):195-200.
17. Hefti F. Direct screw repair spondylolysis with the hooked screw [in German]. *Orthopade* 1997 Sep; 26(9):769-73.
18. Kakiuchi M. Repair of the defect in spondylolysis. Durable fixation with pedicle screws and laminar hooks. *J Bone Joint Surg* 1997 June; 79(6):818-25.
19. Jeanneret B. Direct repair of spondylolysis. *Acta Orthop Scand Suppl.* 1993; 251:111-5.
20. Tokuhashi Y, Matsuzaki H. Repair of defects in spondylolysis by segmental pedicular screw hook fixation. A preliminary report. *Spine* 1996 Sep; 21(17):2041-5.
21. Pai VS, Hodgson B. Assessment of bony union following surgical stabilization for lumbar spondylolysis: a comparative study between radiography and computed tomography. *J Orthop Surg (Hong Kong)*. 2006 Apr; 14(1):17-20.
22. Debnath UK, Freeman BJ, Gregory P, de la Harpe D, Kerslake RW, Webb JK. Clinical outcome and return to sport after the surgical treatment of spondylolysis in young athletes. *J Bone Joint Surg Br* 2003 Mar; 85(2):244-9.
23. Schlenzka D, Remes V, Helenius I, Lamberg T, Tervahartiala P, Yrjonen T, et al. Direct repair for treatment of symptomatic spondylolysis and low-grade isthmic spondylolisthesis in young patients: no benefit in comparison to segmental fusion after a mean follow-up of 14.8 years. *Eur Spine J* 2006 Oct; 15(10):1437-47.
24. Peer KS, Fascione JM. Spondylolysis: A Review and Treatment Approach. *Orthop Nurs* 2007 Mar-Apr; 26(2):104-11.
25. Kant AP, Daum WJ, Dean SM, Uchida T. Evaluation of lumbar spine fusion. Plain radiographs versus direct surgical exploration and observation. *Spine* 1995 Nov; 20(21):2313-7.
26. Szypryt EP, Twining P, Mulholland RC, Worthington BS. The prevalence of disc degeneration associated with neural arch defects of the lumbar spine assessed by magnetic resonance imaging. *Spine* 1989 Sep; 14(9):977-81.
27. Sherman FC, Rosenthal RK, Hall JE. Spine fusion for spondylolysis and spondylolisthesis in children.

Spine 1979 Jan-Feb; 4(1):59-69.

28. DeLoubresse CG, Bon T, Deburge A, Lassale B, Benoit M. Posterolateral fusion for radicular pain in isthmic spondylolisthesis. Clin Orthop Relat Res 1996 Feb; (323):194-201.

29. Phillips FM. The argument for noninstrumented posterolateral fusion for patients with spinal stenosis and degenerative spondylolisthesis. Spine 2004 Jan; 29(2):170-2.

30. Lamberg TS, Remes VM, Helenius IJ, Schlenzka DK, Yrjonen TA, Osterman KE, et al. Long-term clinical, functional and radiological outcome 21 years after posterior or posterolateral fusion in childhood and adolescence isthmic spondylolisthesis. Eur Spine J 2005 Sep; 14(7):639-44.