

## Inequality of obesity and socioeconomic factors in Iran: a systematic review and meta- analyses

Shirin Djalalinia<sup>1</sup>, Niloofar Peykari<sup>2</sup>, Mostafa Qorbani<sup>3</sup>, Bagher Larijani<sup>4</sup>  
Farshad Farzadfar<sup>5</sup>\*

Received: 11 January 2015

Accepted: 4 April 2015

Published: 25 July 2015

### Abstract

**Background:** Socioeconomic status and demographic factors, such as education, occupation, place of residence, gender, age, and marital status have been reported to be associated with obesity. We conducted a systematic review to summarize evidences on associations between socioeconomic factors and obesity/overweight in Iranian population.

**Methods:** We systematically searched international databases; ISI, PubMed/Medline, Scopus, and national databases Iran-medex, Irandoc, and Scientific Information Database (SID). We refined data for associations between socioeconomic factors and obesity/overweight by sex, age, province, and year. There were no limitations for time and languages.

**Results:** Based on our search strategy we found 151 records; of them 139 were from international databases and the remaining 12 were obtained from national databases. After removing duplicates, via the refining steps, only 119 articles were found related to our study domains. Extracted results were attributed to 146596 person/data from included studies. Increased ages, low educational levels, being married, residence in urban area, as well as female sex were clearly associated with obesity.

**Conclusion:** Results could be useful for better health policy and more planned studies in this field. These also could be used for future complementary analyses.

**Keywords:** Obesity, Socioeconomic factors, Iran.

*Cite this article as:* Djalalinia Sh, Peykari N, Qorbani M, Larijani B, Farzadfar F. Inequality of obesity and socioeconomic factors in Iran: a systematic review and meta- analyses. *Med J Islam Repub Iran* 2015 (25 July). Vol. 29:241.

### Introduction

The prevalence of obesity and its health adverse outcomes continues to rise worldwide with alarming rates in developed and developing countries (1-3). Obesity and overweight are considered as one of the leading preventable causes of attributable morbidity and death worldwide (1, 4, 5).

Co-morbidities are either directly caused by increased weight or indirectly related to mechanisms sharing a common cause such as unhealthy behaviors that are mostly affected by socio-economic factors (SEFs) and other related determinants (6).

Evidences revealed that substantial proportion of excess weight mortality intricate-

<sup>1</sup>. PhD, Non-communicable Diseases Research Center, Endocrinology and Metabolism Research Institute, Tehran University of Medical Sciences, Tehran, Iran, & Endocrinology and Metabolism Research Center, Endocrinology and Metabolism Research Institute, Tehran University of Medical Sciences, Tehran, Iran, & Development of Research & Technology Center, Deputy of Research and Technology, Ministry of Health and Medical Education, Tehran, Iran. sh\_djalalinia@razi.tums.ac.ir

<sup>2</sup>. PhD, Non-communicable Diseases Research Center, Endocrinology and Metabolism Research Institute, Tehran University of Medical Sciences, Tehran, Iran, & Endocrinology and Metabolism Research Center, Endocrinology and Metabolism Research Institute, Tehran University of Medical Sciences, Tehran, Iran, & Development of Research & Technology Center, Deputy of Research and Technology, Ministry of Health and Medical Education, Tehran, Iran. n-peykari@razi.tums.ac.ir

<sup>3</sup>. PhD, School of Medicine, Community Medicine Department Alborz University of Medical Sciences, Karaj, Iran, & Non-communicable Diseases Research Center, Endocrinology and Metabolism Population Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran. mqorbani1379@yahoo.com

<sup>4</sup>. MD, Endocrinology and Metabolism Research Center, Endocrinology and Metabolism Research Institute, Tehran University of Medical Sciences, Tehran, Iran. emri@tums.ac.ir

<sup>5</sup>. (**Corresponding author**) MD, MPH, DSc, Non-communicable Diseases Research Center, Endocrinology and Metabolism Research Institute, Tehran University of Medical Sciences, Tehran, Iran. farzadfar@tums.ac.ir

ly is associated with SEFs (6, 7). These findings suggest the importance of social inequalities effects on unhealthy behaviors (8, 9). It is noticeable that some modifiable lifestyle behaviors and factors relating to SEFs may influence these emerging health programs (10).

Recent evidences have shown that, in developing countries, the burden of obesity tend to shift toward some specific socioeconomic groups (2, 11, 12). Socioeconomic status and demographic factors, such as education, occupation, place of residence, gender, age, and marital status have been reported to be associated with excess weight (7). These patterns are mostly complex, dynamic, and may be differ between countries and sub-groups populations (11).

It is clear that decisions about the design, feasibility, planning, funding, implementation, and management of preventive or controlling programs require the most accurate

and reliable evidence provided through related researches (3, 13-15). Health researchers, professionals and policy-makers call for valid evidences to monitor, prevent, and control of obesity problems (12, 13).

Despite priority of the problem, there is an evident gap in the related literature on these topics (1, 2, 15-18). This study aimed to assess the association between SEFs and obesity/overweight in Iranian population. We followed a comprehensive approach to conducting an up-to-date systematic review and meta-analytic comparison of all available studies.

## Methods

### Search strategy

To assess papers on obesity and/or overweight and related socio-economic factors in Iranian population, the relevant empirical literature searched through several electronic databases, including: main domestic da-

Table 1. The Search strategy

Domain	Search strategy
Obesity/overweight	<p>Search strategy in PubMed/Medline            ("Body Mass Index"[Mesh] OR "Body Mass Index"[All Fields] OR "Overweight"[Mesh] OR "Overweight"[All Fields] OR "Obesity"[Mesh] OR "Obesity"[All Fields] OR "Quetelet* Index"[All Fields] AND ("Iran"[Mesh] OR "iran"[All Fields]) OR Iranian[All Fields] OR I.R.Iran[All Fields] OR "I.R Iran"[All Fields] OR ("persia"[MeSH Terms] OR "persia"[All Fields AND "humans"[MeSH Terms])</p> <p>Search strategy in ISI Web of Science            Topic= ("Body Mass Index" OR "Overweight" OR "Obesity" OR "Quetelet* Index") AND (("Iran" OR Iranian OR I.R.Iran OR "persia") OR Address= (Iran))</p> <p>Search strategy in Scopus            (TITLE-ABS-KEY (Body Mass Index" OR "Overweight" OR "Obesity" OR "Quetelet* Index")) AND (TITLE-ABS-KEY (Iran OR Iranian OR I.R.Iran OR Persia) OR (AFFIL (Iran)))</p> <p>IranMedex, SID and Irandoc            "Body Mass Index", "BMI", "Overweight", "Obesity", "Quetelet* Index", "Chaghi", "Shak-hese tudeh e badani", "ezafe vazn", "Dore kamar", in Persian language</p>
Socioeconomic factors	<p>(((((("Socioeconomic Factors"[Mesh] OR "Poverty"[Mesh]) OR "Social Class"[Mesh]) OR "Educational Status"[Mesh]) OR "Employment"[Mesh]) OR "Family Characteristics"[Mesh]) OR "Income"[Mesh]) OR "Occupations"[Mesh]) OR "Social Conditions"[Mesh] OR "Standard of Living"[All Fields] OR "living standard"[All Fields] OR "land tenure"[All Fields] OR "High-Income Population"[All Fields] OR "High Income Population"[All Fields] OR ("socioeconomic factors"[MeSH Terms] OR ("socioeconomic"[All Fields] AND "factors"[All Fields]) OR "socioeconomic factors"[All Fields] OR "inequality"[All Fields]) OR ("socioeconomic factors"[MeSH Terms] OR ("socioeconomic"[All Fields] AND "factors"[All Fields]) OR "socioeconomic factors"[All Fields] OR "inequalities"[All Fields])</p>
Geographic area	<p>((("iran"[MeSH Terms] OR "iran"[All Fields]) OR iranian[All Fields] OR I.R.Iran[All Fields] OR "persia"[MeSH Terms]) OR ((("iran"[MeSH Terms] OR "iran"[All Fields]) OR iranian[All Fields] OR I.R.Iran[All Fields] OR persia[Title/Abstract])) OR ((("iran"[MeSH Terms] OR "iran"[All Fields]) OR iranian[All Fields] OR I.R.Iran[All Fields] OR persia[Text Word])</p>

tabases, Iran-Medex, Scientific Information Database (SID), Irandoc, and also international databases including: PubMed and NLMGateway (for MEDLINE), Institute of Scientific Information (ISI), and SCOPUS. To obtain the most comprehensive results, we searched these data sources using Medical Subject Headings (MeSH) terms, Emtree, and related equal Persian key words for Iranian databases. The references and citations of relevant articles were also assessed. There was no limitation on age, time and language (Table 1).

### Definitions

Overweight and obesity were defined as  $25 \leq \text{BMI} < 30$  and  $\text{BMI} \geq 30$ , respectively (19). SEFs considered as social and economic factors that differentiate the individuals or groups within the social structure. These factors included the social and economic experiences and realities that help mold one's personality, attitudes, and lifestyle (20) (Table 2).

### Study selection and eligibility criteria

We included all available related original and systematic review papers that met our search protocol criteria and excluded article with duplicate citation. Aim to study selection, at the first phase, a reviewer check out all searched titles. Non-relevant papers deleted and those were crossed research ob-

jectives' were kept for more evaluation in the next level. In abstract review phase papers were evaluated, in more detailed, considering reported values or related indexes. In final step, remained papers exactly reviewed based on their full-text contents. When there were multiple publications on the same population, only the largest study or the main source of data was included.

### Quality assessment and data extraction

The quality assessment and data extraction of eligible papers has been followed independently by two research experts and probable discrepancy between them resolved based on third expert opinion. Using Cohen's kappa statistic, agreement of them in quality assessment was 0.92.

Quality assessment form has three parts: general information about the study, sampling quality, and measurement quality. Each study had a unique code and its general information such as the name and characteristics of the corresponding author have been inserted at the top of the form. The sampling quality refers to response rate, sample size, and sampling design and the measurement quality includes type of measurement tools, and accuracy of measurement. The final decision was based on the total scores obtained by each paper in ranking scale of: excellent (13-19), good (6-12) or poor ( $\leq 5$ ). Poor quality papers

Table 2. Socioeconomic factors' definitions

Socioeconomic factors	Classic Definition
Age group	Persons classified by age from birth (INFANT, NEWBORN) to octogenarians and older (AGED, 80 AND OVER).
Sex	The totality of characteristics of reproductive structure, functions, PHENOTYPE, and GENOTYPE, differentiating the MALE from the FEMALE organism.
Educational level	Educational attainment or level of education of individuals.
Marital status	A demographic parameter indicating a person's status with respect to marriage, divorce, widowhood, singleness, etc.
Occupation	Crafts, trades, professions, or other means of earning a living.
Income	Revenues or receipts accruing from business enterprise, labor, or invested capital.
Residence characteristics	Elements of residence that characterize a population. They are applicable in determining need for and utilization of health services.
Urbanization	The process whereby a society changes from a rural to an urban way of life. It refers also to the gradual increase in the proportion of people living in urban areas.
Social condition	The state of society as it exists or in flux. While it usually refers to society as a whole in a specified geographical or political region, it is applicable also to restricted strata of a society.
Social class	A stratum of people with similar position and prestige; includes social stratification. Social class is measured by criteria such as education, occupation, and income.

have been deleted and two other categories considered for data extraction processes.

Data were collected according to a standard protocol including the citation of paper, study design and setting, study year, participants and their recruitment, sex, age, reported Socio-economic Factors, OR (95% CI), main conclusions, and authors' recommendations.

All included papers have been reviewed and the required information have been extracted and inserted in data extraction sheet. In brief, the data extraction sheet contains detailed information on prevalence or mean, standard deviation, standard error of mean, RO (95% CI), sample size, age, sex, province/district of the country, year, coverage of the study (representativeness), scope of study (rural/urban/both), and other related reported values.

#### Statistical analysis

The reported results are presented as Odds Ratio (OR) and 95% confidence interval (CI). For more precise comparison, the Odds Ratio (OR), Confidence Interval for 95% (CI 95%) were calculated for possible cases. Heterogeneity of reported values between studies was assessed by the

Chi-square based Q test and I square statistics. The result of Q test was regarded to be statistically significant at  $P < 0.001$ . Due to severe heterogeneity among studies regarding reported values, overall OR was estimated using random-effect meta-analysis model (using the Der-Simonian and Laird method) (21). Forest plot also was used to present result of meta-analysis schematically. The analyses were conducted using STATA 11 software.

#### Ethical considerations

Present study has been approved by the ethical committee of Tehran University of Medical Science. All of included studies in our review would be cited in all reports and all publications of our study. Whenever we needed more information about a certain study, for obtaining required information, we contacted the corresponding author.

#### Results

We refined data for association between SEFs and obesity/overweight in Iranian population. Based on our search strategy we found 151 records; of them 139 were from international databases and the




Fig. 1. The flow diagram of the study selection process is shown in the

remaining 12 were obtained from national databases. After removing duplicates, via the refining steps, only 119 articles were found related

to our study domain. The flow diagram of the study selection process is shown in the Fig. 1.

Table 3. The results of including papers on socioeconomic factors and obesity/overweight

No	Ref	Study design and Setting	Study year	Participants and their recruitment	Sex (F/M or B:both)	Age (Year)	Socioeconomic Factors	Reported values	Main Conclusion	Recommendations
1	Amani R. et al.(29)	Cross sectional, Population based/Ahvaz	2003	General Population, random sample/ n=637	F	Mean ± sd: 26.9±5.8 ys (18–40 years)	educational levels	educational levels illiterate (n =39) BMI:25.2±4.9 primary(n=242) BMI:26.6±4.9 secondary (n=217) BMI:25.8±4.7 higher (university) (n =139) 25±4.1 (p<0.02)	Women with less educational grades tend to have more body fat. There was Significant difference between BMI and body fat of primary and higher educational levels (P<0.02).	Further studies on related fields with a greater number of subjects of different minority groups are needed.
2	Asgari F, et al.(30)	Cross sectional, Population based/ (STEPS Survey)	2009	General Population, random sample/ n=20917 (51.8% female)	B	20≤ years	age/ sex/ place of residence / employment status	Obesity odds ratio age OR:30-40:2.48(2.20-2.80) female sex: OR:2.63:(2.32-2.78) rural place of residence: OR:1.46(1.33-1.61)  Employment status: Other:1.00 Public employment:0.79(0.68-0.92) Private employment:0.76(0.60-0.99)	OR of obesity went up with increasing age and then it decreased for ages above 50 years. men had lower BMI values than women Rural residents had higher odds of being obese compared with urban residents. Obesity may be more acceptable among unemployed people.	Results may provide better insights of the factors associated with obesity and can be used as a basis to reinforce health programs to prevent obesity in Iran.
4	Azizi F, et al.(31)	Cross sectional, Population based/ Tehran	1999-2001	General Population, random sample/ n=15005 Tehranian children, adolescents, and adults, 44% males and 56% females.	B/ F (56%)	Nearly, 5 % of the study population was between 3–6 years, 6 % between 7–10 years, 9% between 11–14 years, 19% between 15–24 years, 17% between 25–34 years, 16% between 35–44 years, 12% between 45–54 years, and 10% between 55–64 years, and 7% over 64	sex	obesity was significantly higher in women than in men (29.5% vs 14.4 %, p < 0.001) the prevalence of overweight was greater in men than in women (42.5% vs 38.1%, p < 0.01). In adult population, the prevalence of central obesity (high WHR) was greater in women than in men (67.2% vs 33.0 %, p<0.001) Obes female:OR:2.72(2.42-3.07) Over weight female:OR:1.19(1.09-1.31)	More than 60% of adults in the present study were obese or overweight. Moreover, it is worth mentioning that more than 17% of the children and adolescents in our study suffered from obesity or overweight..	The results suggest a need for special attention to health status in Tehranian children and adolescents. In case of comparing TLGS data w

5	Bakhshi E, et al.(32, 33)	Cross sectional, Population based/ National Health Survey (NHS)	1999-2000	General Population, cluster-stratified sampling/ 2728 women and men 1364	B/ F (50 %)	years. 20-75 (mean: 36.9 y)	age/ Economic Index / educational level/ residence / marital status	Obesity odds ratio Obesity odds ratio age OR: 1.02(1.00-1.05) education high education: 0.60 (0.37-0.7) residence in city 0.70(0.47-1.04) marital status Or:1.13(0.96-130)  Economic Index female: 1.002 (0.99-1.01) male: 1.02 (1.01-1.04)Male vs. female: Active workforce:1.21(0.32-4.56) male vs. female: non active workforce:0.40(0.16-1.00)	Among women, factors that increased obesity included age, low education. Among men, these factors included high economic index, low education.	better understanding of the social and cultural mechanisms of obesity in couples.  The preventive strategies based on the affective factors.
6	Behzadnia S, et al.(34)	Cross sectional, Population based/Sari	2009-2010	General Population, multi-stage and stratified randomization /, n: 653	B/ F (55.5 %)	7-12	socio economic status	Higher prevalence of obesity in the children with good socio economic status was found (p=0.001).	In children with moderate SES the prevalence of overweight was 23% (151) and in children with good SES it was 34.9% (227) (p = 0.001).	the need for serious attention to the issue of childhood obesity, performance of more extensive studies, identification of underlying factors precisely and designing the implementation of needed interventions.
7	Dastgiri S, et al.(35)	Cross sectional, Population based/ Tabriz	2006	General Population, Random sampling/ n:200	B/ F (56%)	18≤	sex/ age/ education/ monthly income	Obesity odds ratio female sex OR: 1.81 (0.96-2.66) education 12+:0.41(0.31-0.63) age OR:2.29(1.96-2.99)  Monthly income Female Monthly income(\$US) ≤65:1 66-130:0.87(0.76-0.96) 131-220:0.72(0.62-0.81) ≥221:0.58(0.41-0.67) Male Monthly income(\$US) ≤65:1 66-130:0.75(0.62-0.93) 131-220:0.41(0.32-0.68) ≥221:0.32(0.24-0.53)	Mean BMI was significantly lower in men than in women (25.2 ^ 4.9 vs. 27.1 ^ 5.3 kg m <sup>2</sup> , P, 0.001). Total prevalence of obesity in the area was 22.4% (95% CI: 18,0-27.6), with 24% (95% CI: 18.5-31.4) of women and 18% (95% CI: 12.5-25.6) of men being obese. For both women and men obesity prevalence showed a positive association with age (P for trend ,0.001), while there was a negative correlation of obesity with education and income (P for trend ,0.001). Overweight and obese was significantly more prevalent among women than men and urban- compared to rural-dwellers. A high prevalence of overweight and obesity was seen among individuals who were	crucial necessity of establishing a population-based centre for obesity in the area. More population-based investigations on dietary choices are needed to develop effective preventive strategies to control overweight and obesity disorders in different regions.
8	Esmaeily H, et al.(16)	Cross sectional, Population based/ Great Khorasan	2009	General Population, Cluster-stratified sampling/ n: 4977	B/ F (50.2%)	15-65	sex/ residence in city/ marital status	Obesity odds ratio ever married OR:3.50 (2.72-4.49) Residence in city 2.58(2.13-3.13)	Overweight and obese was significantly more prevalent among women than men and urban- compared to rural-dwellers. A high prevalence of overweight and obesity was seen among individuals who were	A community-based approach using multiple strategies including appropriate education will be required to address this problem



								not high education( $\leq 12y$ ) OR:0.71(0.41-1.0.1) unemployed OR:2.30(1.92-2.76)	divorced or widowed and among house- wives, or individuals with poor education. Urbanization, age, illiteracy, female gender and divorced, or widowed status were significant predictors of obesity ( $p < 0.001$ ).	
								Overweight odds ratio ever married OR:2.71 (2.29-3.21) Residence in city OR:1.80(1.57-2.05) not high education( $\leq 12y$ ) OR:0.89(0.77-1.03) unemployed OR:1.81(1.61-2.04)		
9	Hajian- Tilaki KO, et al. (36)	Cross sectional, Population based/ /Mazandran	2004	General Population, cluster sampling/ n:3600	B/ F (50%)	20-70 M: 38.5 $\pm$ 14.3, F: 37.5 $\pm$ 13.0)	education	Obesity odds ratio Obes high education( $\geq 12y$ ) OR:0.55(0.45-0.71)	The findings of this study indicate that education level is inversely associated with general obesity in both sexes but with abdominal obesity only in women.	Further studies are needed to explore lifestyle factors that are influenced or modified by education in men or women.
10	Janghorbani M, et al. (37)	National Cross- sectional sur- vey/National	2004- 2005	General Population, stratified probabili- ty cluster sam- pling/n:89440	B/ F (60.8%)	15-65 (Mean:39.2)	sex/marit al status	Obesity odds ratio ever married 3.64(3.31-3.99)  Over weight odds ratio ever married male OR:2.24(2.08-2.41) Female OR: 2.36(2.20-2.53)	The marital status appears to influence the likelihood of developing overweight, obesi- ty, and abdominal obesity in both sex ma	There is a need to assess the mechanisms for this associa- tion.
11	Shahraki M, et al. (38)	Cross sectional, Population based/ Sistan and Baluchestan Province	2004- 2006	General Population, random sampling/ n:888	F	$\geq 20$	educa- tion/ marital status	Obesity odds ratio level of education university graduates OR:1.00 intermediary & high school OR:4.20(1.64-10.83) illiterate or low literacy OR:3.70(1.31-10.50) marital status Single OR:0.15(0.02-1.20)	Significant factors associated with obesity by a logistic regression model were education level (OR for university graduates v. illiterate or low literacy levels: 1.00 vs3.7 ( $P \leq 0.001$ ), and OR for married v. single subjects: 1.00 vs.0.15; ( $P \leq 0.001$ )	Obesity prevention should be a relevant topic on the public health agenda in developing countries such as Iran. Without developing effective strategies to modi- fy the current situation, it is likely that the obesity epi- demic will continue in the future.
12	Tavassoli AA, et al. (39)	Cross sectional, Population based/ Islamshahr dis- trict	2003	General Population, random survey/n :1003	F	10-65	place of resi- dence/ educa- tion/ marital status	Obesity odds ratio Place of residence no sig diff Literacy ( year) 0-3 n:166 mean BMI 30(6.0) 4-7 n:288 mean BMI 28.4(5.7) 8 $\leq$ n:268 mean BMI 25.3(5.4) marital status OR:1.18(0.90-1.46)	The mean BMI was significantly higher in married women and in women with less than 8 years of formal education. The mean WHR was significantly higher in women with less than 8 years of education or with more than 6 parity female adults.	The prevention of over- weight and obesity through a healthy diet and increased physical activity should now be an important priority area.
13	Nematy M, et al. (19)	Cross sectional, Population based/ Razavi-Khorasan Province	-	General Population, cluster random sampling/ /n	B/(917 m/1045 fe)	$\geq 60$ years /70.2 $\pm$ 7.8	sex/ education	Obesity odds ratio female sex OR:1.71 (1.43-2.07) illiterate vs literate OR:0.47(0.33-0.66)	Regression analysis results indicated that gender ( $p < 0.001$ ), place of residence ( $p < 0.001$ ), literacy ( $p = 0.01$ ), and source of income ( $p < 0.001$ ) were significantly associated with the incidence	Results reinforce the need to plan strategies for primary prevention of this fast- growing public health prob- lem.

:2000										
14	Sotoudeh G, et al.(10)	Cross sectional, Population based/ / Isfahan, Najaf-Abad, and Arak	2010	General Population, random sampling/ n:12514	B/(6,123 M, 6,391 F)	≥19 38.89±14.93 y	Educa- tion/inco me/ owner- ship of car/ job/	Obesity odds ratio Urban vs rural residence OR:2.32 (1.86-2.86) education OR:0.37(0.26-0.41) married vs single OR:2.65(1.90-3.40) Family members >4 vs <4 OR:1.195(0.998-1.432) Income 25-75% vs <0.25% OR:1.134(0.906-1.418) Income >0.75% vs <0.25% OR:1.294(0.997-1.679) Ownership of car OR:2.044(1.711-2.442) Government job vs retired/not working OR:0.700(0.511-0.958) Private job vs retired/not working OR:0.599(0.482-0.744)	of overweight or obesity. This study showed that 40.6% of elderly subjects were over- weight or obese. The results of the present study showed significant differences in some factors associated with obesity among men and women; such differences should be taken into account for interventional pro- grammes at the individual level and for providing long-term public-health policies.	The association of different socioeconomic and lifestyle factors and their gender differences should be considered for culturally- appropriate intervention strategies to be implemented at the population level for tackling obesity and associ- ated cardio- metabolic risk factors.
15	Maddah M, et al. (40)	Cross sectional, Population based/ / Rasht	2005	General Population, multistage sam- pling meth- od/n:1054	F	14-17	age	Obesity odds ratio Age group OR: 2.0( 0.18-3.09)	These data indicate that overweight is highly prevalent among adolescent girls, especially in lower social groups in Rasht, and the rate is exceeding those reported in other parts of the country.	Preventive strategies need to be adopted to combat the epidemic of overweight and obesity in this population.
16	Navadeh S,et al. (41)	Cross sectional, Population based/ /National	2005	General Population, random sampling/ n:36252	F	20-65	Age/emp loyment/ residence place/ marital sta- tus/educa tion	Obesity odds ratio HWs vs. employed OR: 1.39 ( 1.18-1.63) ever married OR: 2.15(1.84-2.51) residence in urban OR:1.37(1.21-1.55) Low SEE OR :1.22(1.02-1.45) high education OR: 0.87(0.83-0.91) age OR:1.38(1.33-1.43)	HWs vs. employed women had the adjusted OR 1.39 (CI95%, 1.18-1.63) for obesity. Older women, with higher educational level and socioeco- nomic status and those living in urban areas were at risk of obesity. In comparison to HWs, working as an Official Clerk (OR=0.66) associated with a decrease in odds of obe- sity significantly, while others did not.	Being as HW is an inde- pendent significant factor for obesity in women. Pre- ventive health care programs to reduce risk of obesity in women should be applied, considering their occupation for achieving more effec- tiveness.
17	Maddah M, etal.(42)	Cross sectional, Population based/ / Rasht	2008	General Population, random sampling/ n:4035	B/(6028 M ,6223F)	F: 38.6±5.9 M: 43.1±6.4 years	education	Obesity odds ratio High Education OR:0.97(0.74-1.27)  Overweight odds ratio High Education OR: 0.77(0.65-0.92)	These results highlight the importance of socioeconomic indicators in obesity research in Iran. Overall prevalence of overweight and obesity was positively related to educational level in men. The lowest rate of overweight/obesity was observed in very low educated men and very high educated women.	-



Table 4. The results of heterogeneity test (I squares) and Estimated OR for related SEFs

No	Variable	Practical definition	Number of studies	Model of meta-analyses	Estimated OR	95% CI of OR	Results of heterogeneity test (I squares)
1	Sex	Female vs. male	3	Random-effect meta-analysis model	2.06	1.45-2.93	88.9%, p=0.00
2	Age	30< vs. ≤30 yo	4	Random-effect meta-analysis model	1.66	1.24-2.22	99.2%, p=0.00
3	Marital status	Ever married vs. never married	6	Random-effect meta-analysis model	2.14	1.36-3.38	97.7%, p=0.00
4	Education	High education (more than 12yos) vs. low education	7	Random-effect meta-analysis model	0.60	0.43-0.83	92.9%, p=0.00
5	Living place	Urban vs. rural residence	5	Random-effect meta-analysis model	1.58	1.18-2.11	93.4%, p=0.00

Table 3, shows the association of SEFs and obesity/overweight based on reported factors in eligible population-based studies. For more precise comparison, the OR and 95% confidence interval were calculated for possible cases. We have also included studies' methodology information.

Considering the systematic review results, all included studies were cross-sectional ones. Eight articles were population-based studies and the others were clinical- and hospital-based. Regarding the geographic distribution, we found 4 national, 6 provincial, 5 community, and 1 district level studies. All participants were from general population. All of searched articles were in English or Persian language. Although, we haven't limit the search strategy on certain time; retrieving articles were between 1973 and 2014. Included articles were published between 2002 and 2014 and the studies were conducted between 1999 and 2010. In general, these results were attributed to 146596 person/data from included studies. Excluding 5 studies that focused on female sex, the remained studies covered both sex.

Only one study considered inequality assessment index about obesity. Based on first run of non-communicable disease sur-

veillance study data (STEPS study, 2005) in Shahroud, concentration index for obesity was  $(0.038 \pm 0.036)$  for both sex. It was reported for female  $(-0.005 \pm 0.039)$  and male  $(0.194 \pm 0.070)$ . It showed that obesity was more prevalent in men with high socioeconomic levels (22).

Other studies focused on different components of SEFs. The findings are scattered, with different ranges of values for Odds. On the other hand, non-standard classifications of SEFs led to greater complexity in estimation and comparability of the results. For instance, regarding the education, results have been presented as years of schooling through continue variables or different frame of categorization of educational levels. This factor, only in three studies has been reported by sex; among them the highest OR were reported for university educated persons vs. primary education (for women  $0.41(0.31-0.63)$  and men  $1.2+0.62(0.36-0.71)$ ). Severe heterogeneity of overweight results, did not allow us to include them in meta-analyses.

The results of heterogeneity test (I squares) and estimated OR for related SEFs are presented in Table 4. Considering the severe heterogeneity among reported val-




Fig. 2. The forest plot of OR of obesity according to sex in Iran using random-effect mode




Fig. 3. The forest plot of OR of obesity according to age in Iran using random-effect mode

ues, to overall estimation of OR, for possible variables, of SEFs, we used random-effect model.

Figs. 2-6 present the forest plot of eligible articles for estimating overall OR of obesity according to related SEFs in Iran.

### Discussion

This is an updated systematic review on reported association between SEFs and

obesity/overweight among the Iranian population according to sex, age, and levels of reported values, from 1973 to 2014. In our study, a total of 17 studies were eligible for inclusion. Our findings provide evidence-based data on the association between SEFs and obesity/overweight.

The analysis revealed that female sex (OR: 2.06, 95%CI: 1.45-2.93), age (OR: 1.66, 95%CI: 1.24-2.22), married status




Fig. 4. The forest plot of OR of obesity according to marital status in Iran using random-effect mode




Fig. 5. The forest plot of OR of obesity according to education in Iran using random-effect mode

(OR: 2.14, 95%CI: 1.36-3.8), and urban residency (OR: 1.58, 95%CI: 1.18-2.11) increase the risk of obesity. It is considerable that academic education (OR: 0.60, 95%CI: 0.43-0.83) have a reverse associa-

tion with obesity. Other related studies showed a significant reverse association between low socio-economic situation and obesity/overweight, and direct association between factors such as: female sex, mari-




Fig. 6. The forest plot of OR of obesity according to living place in Iran using random-effect mode

tal status, low education, non-employment and obesity (23-26).

There are few studies on association between SEFs and obesity/overweight in Iran. Moreover the scarcity of data, different studies followed different approaches in their methodology. We also faced a large variety of combinations of unequal determinant factors. Considering above, a case by case, conservative approach was followed to make an accurate conclusion (17, 18, 26).

According to our analysis, age, sex, place of residence, employment, and marital status was studied based on available data. Based on them, age  $\geq 30$ , female sex (vs. male), residence in rural areas (vs. urban), high education (vs. non-academic education), unemployment (vs. employment), and ever married situation (vs. never married) were independent significant factor for obesity (Table 3).

It is mentionable that, only one study considered inequality assessment index about obesity. Based on first run of non-communicable disease surveillance study data (STEPS study, 2005) in Shahroud, concentration index for obesity was  $(0.038 \pm 0.036)$  for both sex. It was reported

$-0.005 \pm 0.039$  and  $0.194 \pm 0.070$  for female and male, respectively; it showed that obesity is most probable in high socioeconomic levels men (22).

Using principal component analysis, in a comprehensive analyses of national and sub-national mortality effects of metabolic risk factors and smoking in Iran, the sub-national regions were defined based on a combination of geography SES. SES was measured using an index constructed from variables from the 2006 census, including years of schooling, employment rates, and family assets (26).

In another national survey of school student with 5,528 student participant, aged 10–18 years, structural equation modeling (path analysis) was applied to evaluate the association between SES and BMI. In this study, higher socioeconomic status was directly associated with BMI in both sexes. In this study, indexes for evaluation of family's SES were defined based on questions on the parental level of education, parental occupational status, number of inhabitants in home, and possessing a family private car (27).

In another related experience, to assess the prevalence of Iranian adolescents'

growth disorders, sub-national geographical and social classification of the country was considered as the base of SES for definition of living regions (28). It is also important that, there was no comprehensive study for sub-national evaluation of association between obesity / overweight and its related SEFs in Iran.

Considering the previous studies, the present study has several achievements. This study presents scientific evidences to depict association between SEFs and obesity/overweight among the Iranian population. All available sources of data and domestic data-bases were searched using English/Persian equivalent search terms. As the main limitation, the validity and applicability of our systematic review depends on the quality of the primary studies that are included. As another point, heterogeneity of searched results limits the generalization of our findings.

### Conclusion

According to our knowledge, this is the first systematic review of association between SEFs and obesity/overweight in Iran that provide practical information on associations between several socio-demographic factors and obesity. Increased ages, low educational levels, being married, residence in urban area, as well as female sex were clearly associated with obesity. Results could be useful for better health policy and more planned studies in this field. These also can be used for future complementary analyses.

### Conflict of interests

The authors declare that they have no competing interests.

### Acknowledgements

This study was a part of a Ph.D. dissertation completed by Shirin Djalalinia. Project granted by Ministry of Health and Medical Education of Islamic Republic of Iran and Setad-e-Ejraie Farmane Imam. We would like to express our thanks to NCDRC. FF, SD, and NP participated in study design.

SD, NP, and MQ drafted the manuscript. SD, MQ and NP participated in statistical analysis and interpretation of results. SD and NP participated in literature review and data extraction. FF, BL and SD participated in revision of paper. All authors read and approved the final manuscript.

### References

1. Finucane MM, Stevens GA, Cowan MJ, Danaei G, Lin JK, Paciorek CJ, et al. National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9.1 million participants. *The Lancet* 2011; 377(9765):557-67.
2. Stevens GA, Singh GM, Lu Y, Danaei G, Lin JK, Finucane MM, et al. National, regional, and global trends in adult overweight and obesity prevalences. *Population Health Metrics* 2012;10:22.
3. Jafari-Adli S, Jouyandeh Z, Qorbani M, Soroush A, Larijani B, Hasani-Ranjbar S. Prevalence of obesity and overweight in adults and children in Iran; a systematic review. *Journal of Diabetes & Metabolic Disorders* 2014;13(1):121.
4. Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United States, 2000. *Jama* 2004;291(10):1238-45.
5. Djalalinia S, Qorbani M, Peykari N, Kelishadi R. Health impacts of obesity. *Pakistan Journal of Medical Sciences* 2015;31(1):4.
6. Bray G. Medical consequences of obesity. *The Journal of clinical endocrinology and metabolism* 2004;89(6):2583.
7. Wang Y, Beydoun M. The obesity epidemic in the United States-gender, age, socioeconomic, racial/ethnic, and geographic characteristics: a systematic review and meta-regression analysis. *Epidemiologic reviews* 2007;29:6.
8. Nandi A, Glymour M, Subramanian S. Association among socioeconomic status, health behaviors, and all-cause mortality in the United States. *Epidemiology (Cambridge, Mass)* 2014; 25(2):170.
9. Vakili M, Abedi P, Sharifi M, Hosseini M. Dietary diversity and its related factors among adolescents: a survey in Ahvaz-Iran. *Global journal of health science* 2013;5(2):181-6.
10. Tavassoli AA, Gharipour M, Khosravi A, Kelishadi R, Siadat ZD, Bahonar A, et al. Gender differences in obesogenic behaviour, socioeconomic and metabolic factors in a population-based sample of Iranians: the IHHP study. *Journal of health, population, and nutrition* 2010;28(6):602-9.
11. Soon H. Pattern of dietary behaviour and obesity in Ahwaz, Islamic Republic of Iran. *Eastern Mediterranean Health Journal* 2001;7(1-2):163-70.

12. Danaei G, Singh GM, Paciorek CJ, Lin JK, Cowan MJ, Finucane MM, et al. The global cardiovascular risk transition: Associations of four metabolic risk factors with national income, urbanization, and western diet in 1980 and 2008. *Circulation* 2013;127(14):1493-502.
13. Djalalinia S, Kelishadi R, Qorbani M, Peykari N, Kasaeian A, Saeedi Moghaddam S, et al. Suggestions for Better Data Presentation in Papers: An Experience from a Comprehensive Study on National and Sub-national Trends of Overweight and Obesity. *Archives of Iranian medicine* 2014;17(12):830-6.
14. Di Cesare M, Khang Y, Asaria P, Blakely T, Cowan M, Farzadfar F, et al. Inequalities in non-communicable diseases and effective responses. *Lancet* 2013;381(9866):585.
15. Mirzazadeh A, Haghdoost A, Sadeghirad B, Bahreini F, Ashrafi K, Chack LE, et al. The Prevalence of Obesity in Iran in Recent Decade; a Systematic Review and Meta-Analysis Study. *Iranian journal of public health* 2009;38:1-2.
16. Esmaily H, Azimi-Nezhad M, Ghayour-Mobarhan M, Parizadeh MR, Safarian M, Parizadeh MJ, et al. Association between socioeconomic factors and obesity in Iran. *Pakistan Journal of Nutrition* 2009;8(1):53-6.
17. Peykari N, Sepanlou S, Djalalinia S, Kasaeian A, Parsaeian M, Ahmadvand A, et al. National and Sub-national Prevalence, Trend, and Burden of Metabolic Risk Factors (MRFs) in Iran: 1990-2013, Study Protocol. *Archives of Iranian medicine* 2014;17(1):54.
18. Kelishadi R, Hovsepian S, Qorbani M, Jamshidi F, Fallah Z, Djalalinia S, et al. National and sub-national prevalence, trend, and burden of cardiometabolic risk factors in Iranian children and adolescents, 1990-2013. *Archives of Iranian medicine* 2014;17(1):71.
19. Nematy M, Sakhdari A, Ahmadi-Moghaddam P, Aliabadi M, Kimiagar M, Ilaty AA, et al. Prevalence of obesity and its association with socioeconomic factors in elderly Iranians from Razavi-Khorasan province. *The Scientific World Journal* 2009;9:1286-93.
20. <http://www.ncbi.nlm.nih.gov/mesh/68012959>.
21. Gardiner J, Luo Z, Roman L. Fixed effects, random effects and GEE: what are the differences? *Statistics in medicine* 2009;28(2):221.
22. Emamian MH, Alami A, Fateh M. Socioeconomic inequality in non-communicable disease risk factors in Shahroud, Iran. *Iranian J Epidemiology*. 2011; 7:44-51.
23. Monteiro CA, Moura EC, Conde WL, Popkin BM. Socioeconomic status and obesity in adult populations of developing countries: a review. *Bulletin of the World Health Organization* 2004;82(12):940-6.
24. McLaren L. Socioeconomic status and obesity. *Epidemiologic reviews* 2007;29(1):29-48.
25. Brennan SL, Henry MJ, Nicholson GC, Kotowicz MA, Pasco JA. Socioeconomic status, obesity and lifestyle in men: the Geelong Osteoporosis Study. *Journal of men's health* 2010;7(1):31-41.
26. Farzadfar F, Danaei G, Namdaritabar H, Rajaratnam JK, Marcus JR, Khosravi A, et al. National and subnational mortality effects of metabolic risk factors and smoking in Iran: a comparative risk assessment. *Population Health Metrics* 2011;9.
27. Kelishadi R, Motlagh ME, Roomizadeh P, Abtahi SH, Qorbani M, Taslimi M, et al. First report on path analysis for cardiometabolic components in a nationally representative sample of pediatric population in the Middle East and North Africa (MENA): the CASPIAN-III Study. *Annals of nutrition & metabolism* 2013;62(3):257-65.
28. Bahreynian M, Motlagh ME, Qorbani M, Heshmat R, Ardalan G, Kelishadi R. Prevalence of growth disorders in a nationally representative sample of Iranian adolescents according to socioeconomic status: the CASPIAN-III Study. *Pediatrics & Neonatology* 2014.
29. Amani R. Comparison between bioelectrical impedance analysis and body mass index methods in determination of obesity prevalence in Ahvazi women. *European journal of clinical nutrition* 2007;61(4):478-82.
30. Asgari F, Biglarian A, Seifi B, Bakhshi A, Miri HH, Bakhshi E. Using quadratic inference functions to determine the factors associated with obesity: findings from the STEPS Survey in Iran. *Annals of epidemiology* 2013;23(9):534-8.
31. Azizi F, Rahmani M, Emami H, Mirmiran P, Hajipour R, Madjid M, et al. Cardiovascular risk factors in an Iranian urban population: Tehran lipid and glucose study (phase 1). *Sozial- und Preventivmedizin* 2002;47(6):408-26.
32. Bakhshi E, Eshraghian MR, Mohammad K, Foroushani AR, Zeraati H, Fotouhi A, et al. The positive association between number of children and obesity in Iranian women and men: results from the National Health Survey. *BMC public health* 2008;8:213.
33. Bakhshi E, Mohammad K, Eshraghian MR, Seifi B. Factors related to obesity among Iranian men: results from the National Health Survey. *Public health nutrition* 2010;13(9):1389-94.
34. Behzadnia S, Vahidshahi K, Hamzeh Hosseini S, Anvari S, Ehteshami S. Obesity and related factors in 7-12 year-old elementary school students during 2009-2010 in Sari, Iran. *Medicinski glasnik: official publication of the Medical Association of Zenica-Doboj Canton, Bosnia and Herzegovina* 2012;9(1):86-90.
35. Dastgiri S, Mahdavi R, TuTunchi H, Faramarzi E. Prevalence of obesity, food choices and socio-economic status: a cross-sectional study in the north-west of Iran. *Public health nutrition*



2006;9(8):996-1000.

36. Hajian-Tilaki KO, Heidari B. Association of educational level with risk of obesity and abdominal obesity in Iranian adults. *Journal of public health (Oxford, England)* 2010;32(2):202-9.

37. Janghorbani M, Amini M, Rezvanian H, Gouya MM, Delavari A, Alikhani S, et al. Association of body mass index and abdominal obesity with marital status in adults. *Archives of Iranian medicine* 2008;11(3):274-81.

38. Shahraki M, Shahraki T, Ansari H. The effects of socio-economic status on BMI, waist:hip ratio and waist circumference in a group of Iranian women. *Public health nutrition* 2008;11(7):757-61.

39. Sotoudeh G, Khosravi S, Khajehnasiri F, Khalkhali HR. High prevalence of overweight and

obesity in women of Islamshahr, Iran. *Asia Pacific journal of clinical nutrition* 2005;14(2):169-72.

40. Maddah M. Overweight and obesity among Iranian female adolescents in Rasht: more overweight in the lower social group. *Public health nutrition* 2007;10(5):450-3.

41. Navadeh S, Sajadi L, Mirzazadeh A, Asgari F, Haghazali M. Housewives' obesity determinant factors in Iran; national survey - stepwise approach to surveillance. *Iranian journal of public health* 2011;40(2):87-95.

42. Maddah M, Solhpour A. Obesity in relation to gender, educational levels and living area in adult population in Rasht, northern Iran. *International journal of cardiology* 2010;145(2):310-1.