

## Prevalence, etiology, and types of dental trauma in children and adolescents: systematic review and meta-analysis

Saber Azami-Aghdash<sup>1</sup>, Farbod Ebadifard Azar<sup>\*2</sup>, Fatemeh Pournaghi Azar<sup>3</sup>  
Aziz Rezapour<sup>4</sup>, Mohammad Moradi-Joo<sup>5</sup>, Ahmad Moosavi<sup>6</sup>, Sina Ghertasi Oskouei<sup>7</sup>

Received: 18 May 2015

Accepted: 1 July 2015

Published: 10 July 2015

### Abstract

**Background:** Dental traumas are common among children and adolescents in many societies posing health and social problems. The aim of this study was to conduct a systematic review and meta-analysis on prevalence, etiology, types, and other epidemiologic aspects of dental trauma in children and adolescents (0-18 years old).

**Methods:** In this systematic meta-analytical review, data were collected searching for key words including traumatic dental injuries, dental trauma, dental injury, dental trauma, tooth injuries, tooth trauma, traumatized teeth, dentoalveolar trauma, oral trauma, epidemiology, etiology, prevalence, incidence, occurrence, child\*, and adolescence in the following databases: Scopus, CINAHL, Science Direct, PubMed and Google scholar.

**Results:** From the total of 3197 articles, 44 completely relevant papers were included in the study. The prevalence of dental trauma was variable based on geographical area and was estimated 17.5% in the population, with higher prevalence in boys. Falling was the major cause for dental trauma, and the most frequent location was home. The most frequent type of trauma was enamel fracture.

**Conclusion:** A relatively high prevalence was detected for dental trauma, which calls for effective planning and intervention to prevent the occurrence in children and adolescents. These may include special care for children, eliminating fall-prone areas, installing safety measures at homes, using protective appliances in sports, education, and raising the knowledge and availability of services to address enamel fracture. Region-specific criteria should be taken into account in programs and interventions.

**Keywords:** Dental Trauma, Prevalence, Etiology, Type, Children and Adolescents, Systematic Review, Meta-analysis.

**Cite this article as:** Azami-Aghdash S, Ebadifard Azar F, Pournaghi Azar F, Rezapour A, Moradi-Joo M, Moosavi A, Ghertasi Oskouei S. Prevalence, etiology, and types of dental trauma in children and adolescents: systematic review and meta-analysis. *Med J Islam Repub Iran* 2015 (10 July). Vol. 29:234.

### Introduction

Traumatic injuries not only pose a health risk worldwide, but are also regarded among serious social problems (1). One important category is dental trauma (2),

accounting for a major part of health problems in children and adolescents (3-6). Studies conducted in different countries report various prevalence rates for traumatic dental injuries among children and ado-

<sup>1</sup>. PhD Student, Road Traffic Injury Research Center, Tabriz University of Medical Sciences, Tabriz, Iran. [saberazami@yahoo.com](mailto:saberazami@yahoo.com)

<sup>2</sup>. (**Corresponding author**) Professor, Health Management and Economics Research Center, Iran University of Medical Sciences, Tehran, Iran. [f\\_ebadi@yahoo.com](mailto:f_ebadi@yahoo.com)

<sup>3</sup>. Assistant Professor, Road Traffic Injury Research Center, Tabriz University of Medical Sciences, Tabriz, Iran. [pournaghiazarm@yahoo.com](mailto:pournaghiazarm@yahoo.com)

<sup>4</sup>. Assistant Professor, Department of Health Economics, Health Management and Economics Research Center, Iran University of Medical Sciences, Center of Excellence in Health Management and Economics, Health Management and Information Sciences, Iran University of Medical Sciences, Tehran, Iran. [rezapoor\\_a57@yahoo.com](mailto:rezapoor_a57@yahoo.com)

<sup>5</sup>. MSc, Cancer Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran. [moradijoo@gmail.com](mailto:moradijoo@gmail.com)

<sup>6</sup>. Assistant Professor, Department of Health and Community Medicine, Dezful University of Medical Sciences, Dezful, Iran. [dr\\_ahmad\\_mosavi@yahoo.com](mailto:dr_ahmad_mosavi@yahoo.com)

<sup>7</sup>. MD, Department of Pediatric Dentistry, Faculty of Dentistry, Tabriz University of Medical Sciences, Tabriz, Iran. [ghertasis@gmail.com](mailto:ghertasis@gmail.com)

lescents (7-10). Dental trauma occurs in children mainly because of their weak balance and just having learnt to walk (11).

Demographic evaluations indicate a higher prevalence for trauma in males compared to females (12-14). Falling, fight, sports, accidents, and hitting items or people are among common etiologic factors (15-18). Home setting, school and street are places with the highest frequency of dental traumatic injuries (19,20), which most importantly include enamel fracture and enamel and dentin fracture without pulp exposure (21-24).

According to the heterogeneity among the individual study results and the importance of preventing dental trauma, it is of importance to accurately determine the prevalence of dental trauma using appropriate research methodology. Also identifying the influencing factors is essential for better planning, decision-making and intervention.

Therefore, the present study aimed to conduct a systematic review and meta-analysis on the prevalence of dental trauma among children and adolescents in different countries and other related variables.

## Methods

This study was a systematic and meta-analytical review. The required data were collected searching for key words including traumatic dental injuries, dental trauma, dental injury, dental trauma, tooth injuries, tooth trauma, traumatized teeth, dentoalveolar trauma, oral trauma, epidemiology, etiology, prevalence, incidence, occurrence, child\*, and adolescence in Scopus, CINAHL, Science Direct, PubMed, Google scholar. Manual journal searching and an extensive hand search of the Gray Literature were also conducted. Articles from the reference list of the studies which were found to be relevant were also considered. Article publication time limit was set 1995 onward.

Articles in English and Persian that reported dental trauma (injury to the teeth and/or periodontium, and nearby soft tis-

ues) and those studies (2) that were conducted on children and adolescents with age range of 0-18 years and cross-sectional studies were also included. Exclusion criteria were as follows: studies on trauma from sport activities, studies on trauma from medical interventions such as endoscopy, studies on individuals with medical or special conditions, conference presentations, case reports and interventional studies (such as randomized controlled trials). Two reviewers evaluated the articles according to the checklist of strengthening the Reporting of Observational Studies in Epidemiology (STROBE) (Appendix. 1) and controversies were referred to a third party. First, articles with non-relevant titles to the subject of this review were excluded. Then, the abstracts and the full texts of the articles were reviewed respectively to exclude those articles that matched the exclusion criteria of the study, or had a weak relevance to the subject of the study. Computer software for reference management (Endnote X5) was used for organizing and assessing the titles and abstracts as well as recognizing the repetitive items.

The searches returned 3197 articles, but those articles that were non-relevant, and repetitive between databases, with weak relevance to the study or matched the exclusion criteria were excluded; finally, 44 relevant articles were selected (Fig. 1). These articles were fully read, and the required data for the systematic review were extracted into the extraction table designed for the purpose of the study in spreadsheet computer software (Excel, Microsoft Office; Microsoft, US). SPSS 17 was used to analyze some of the quantitative data.

To determine the overall prevalence of dental traumatic injuries and perform the meta-analysis, computer software (CMA-Comprehensive Meta-Analysis; Englewood, NJ, USA) was employed. The results were reported using forest plots, in which the size of each square indicates the sample size and the lines drawn on each side of the square indicate the confidence interval (CI) of approximately 95% for den-




Fig.1. Literature review and retrieval flow diagram

tal trauma.

The overall prevalence of dental traumatic injuries was estimated using the random model. Either all or some of the variables were treated as if they were random. Random model was in contrast with the fixed model that demonstrates the observed quantities with respect to the explanatory variables which are treated as if the variable were not random.

### Results

From a total of 3197 relevant articles, 44 completely relevant articles were included in the study. Most studies were conducted in Brazil (16 articles), and the oldest was conducted in Jordan (1995); most of the studies were done in dental clinics, and only two out of the 44 (4%) studies were conducted in IRAN. The overall number of participants was 69502 children and adolescents aged 0-18 years (mean, 1579 each study). The mean±SD prevalence of dental trauma was 10.2% ± 4.70 among boys and 6.5% ± 3.40 girls, which was statistically significant (p< 0.001).

The mean of falling was the most frequent cause of dental trauma. The most important etiological factors of traumatic dental injuries are shown in Fig. 1.

Places with the highest frequency of traumatic dental injuries are demonstrated in Table 1.

The mean of enamel fracture had the highest frequency (55%) among the types of dental trauma in children and adolescents (Fig. 3).

To perform the meta-analysis, the heterogeneity index was determined within the articles using the heterogeneity I<sup>2</sup> test. After confirmation of heterogeneity of the




Fig. 2. Trauma place of occurrence




Fig. 3. Type of dental trauma

Table 1. Extraction table for Prevalence, etiology, and types of dental trauma in children and adolescents

Au-Cu-Ye	Sample	prevalence	Gender	Etiology	place	Type/teeth affected
Schuch et al [25], Brazil,2012	1210-children aged 8–12	12.6%	M (6.8), F(5.8)	Fall (35), Assault (40), Accident (8), Sport (10), other (7)	Home (55), School (18) Street (15),other (12)	Enamel fracture (10.6), dentin Fracture (3.2), other (.5)
Martins et al [26], Brazil,2012	590 children aged 7-14 years	12.7%	M (7.9), F(4.8)	-	-	-
Piovesan et al, Brazil,2012	441 children-of 12- to 59-month-old	31.7%	M(18.6), F(13.1)	-	-	fracture of enamel only (86.91), fracture of enamel and dentin(4.20),fracture involving pulp(3.27),other (5.69)
Dame'-Teixeira et al[27]. Brazil,2012	1528-12-year-old	34.8%	-	-	-	Enamel fracture only(71.16),Enamel-dentin fracture (25.84), other (2.99)
Teixeira et al [28], Brazil,2012	1528-12-year-old Schoolchildren.	34.79%	M(40.8) F(28.6)	Not remember the cause (53.64%), Falls (15.46%), sports (11.08%), collision against objects or people (10.60%), violence (.7%), traffic accidents (.5%), other causes (7.98%)	Home (22.32%), school (8.52%) elsewhere (15.50%)	
Piovesan et al [29], Brazil, 2011	792 12-yearold schoolchildren,	9.7%				Crown fracture of enamel only ( 94.5), Crown fracture of enamel and dentin ( 4.4), Crown fracture involving pulp ( 1.1)
Jorge et al [30], Brazil, 2011	891 Adolescents from schools.	24.7%	M(27.6) F(22.8)	Unknown (33.2), Falls (17.7), Playing(12.7), Collision (10), Violent incident (7.3), Sports(6.4), Use of teeth for functions other than eating (5.9), Chewing (5), Traffic accidents (1.8),		
Aldrigui, et al [31], Brazil, 2011	260 children aged from 2 to 5 years	33.5%				
Norton and O'Connell [32], Ireland,2012	839- children between 9 and 84 months	25.6%	M(26.7) F(24.6)		Most injuries occurred within and around the home, 46.9% and 35.7%.	enamel fracture (39.4%), discoloration of the crown (20.2%)
Bendoand et al [33], Brazil,2010	1612 children aged 11 to 14	17.1%	M (19.9%) F (15%)	Falls (43.6), Sports (20.4), Unknown (25.5), Others (10.5)	Home (41.8), School (14.2), Street (10.5), Unknown (24.4), Others (9.1)	Enamel fracture (63.6), Enamel-dentin fracture (15.3), Complicated crown fracture (1.8), Lateral luxation (0.4), Avulsion(0.7), Restoration (23.3)
Diaz et al [34], Chile,2010	1719 from 1 to 15 years of age,	37.9%				
Altun et al, Turkey[35],2009	4956 children aged 6–12 years	9.5%	M (5.4%) F (4.1%)	Fall (40.3), Impact with a hard object (30.5), Bicycle/tricycle accident (25.4), Other 18 (3.8)		Enamel fractures (44.6), Enamel/dentin fracture (19.0), Intrusive luxation (13.4), Lateral luxation (7.3), Subluxation( 6.1), Enamel/dentin/pulp fracture (5.5), Crown discoloration(5.2), other (7.1)

Noori and Al-Obaidi [36], Iraq, 2009	4015, 6- to 13-year-old children	6.1%	M (3.7%) F (2.4%)	Falls (60.9), Playing (21.4), Impact with a hard object (9.1), other (8.6)	At home (69.9), school (17.7), on street (18.9), Unknown (2.5)	Enamel fracture (36.6%), enamel-dentine fracture (35.4%), and concussion (11.5%). Other (16.5)
Avsar and Topaloglu [37], Turkey, 2009	563- children 0–3 years	17.4%	M (10.3%) F (7.1%)	Falls (73.5), Striking objects (14.2), Traffic accident (1), Child abuse (2), Unknown (9.2)		
Naidoo and et al [38], South Africa, 2009	1665 children aged 11–13 years	6.4%	M (4.2%) F (2.2%)	Falls (43.4%), Sport (13.2%), collision With objects (9.4%), Unknown (8.5%)	Homes (46.3%), schools (35.8%), on a street (5.7%)	Enamel fracture (69.1%), enamel and dentine (22.8%), enamel, dentine and pulp (5.7%)
Jorge et al [30], Brazil, 2009	519 infants and toddlers between 1 and 3 years of age	41.6%		Falls (29.8%), collisions (6.8%)	Home (32%), street (1.7%), day care centers (0.8%)	Enamel fractures (37.2%), enamel–dentin fractures (5.7%),
Fakhruddin et al [39], Canada, 2008	2422 12- to 14-year-old Ontario schoolchildren	11.4%		Falls (24.8%), Sport (23.7%), collision with objects (10%), violence (5.9%), other (35.6)	School (24.4), Home (21.1), street (7.4)	
Pattussi et al, Brazil [40], 2006	1302 14- to 15-year-old adolescents	16%	M (9.6%) F (6.4%)	Playing (48.1), Sports (13.3), Teeth misuse (9.0), Violence (5.7), Other causes (14.8), Don't know (9.0)	Home (44.3), Street/walkway (26.7), School (10.0), Other places (10.0), Don't know (9.0)	
Locker [41], Canada, 2005	3010 grade 8 children	18.5%				
Traebert et al [42], Brazil, 2003	307, 12-year-old school Children	18.9%	M (11.7%) F (7.2%)	Fall (47.9), Collision (37.5), Traffic accident (2.1), Eating (2.1), Unknown (10.4)	Home (60.4), School (18.4), street (18.6), Swimming pool (2.4)	
Nicolau et al [43], UK, 2001	652, 13-year-old adolescents	20.4%	M (13.9%) F (6.5%)	Fall (24.1), Collisions (15.0), Traffic accidents (10.5), Misuse of teeth (6.0), Sports (2.3), Violence (1.5), Unknown (40.6)		
Cortes et al [44], Brazil, 2001	3702-schoolchildren aged 9–14 years	12%	M (7.2%) F (4.8%)			
Vanderas and Papagiannoulis [4], Greece, 1999	199 children aged 8 to 10 years	16.6%	M (11.5%) F (5.1%)			Enamel fractures (75.8) Infractions (24.2)
Marcenes and et al, Syria [45], 1999	1087 children aged 9 to 12 years,	8%	M (5.3%) F (2.7%)	Fall (9.1), Collision (32), Traffic accidents (24.1), Violence (42.5), Other (3.4), Missing information (4.6)		
Chen et al [46], Taiwan, 1999	1200 second grade students	16.5%	M (10.2%) F (6.3%)	Collision (65.3), Fall (26.9), Violence (2.6), Sport (3.6), Other (1.6),	Home (63.7), School (23.8), other (12.4)	
Petti et al [47], Italy, 1997	938, 6- to 11-year-old children	21.3%		Indoor play (31.5), Outdoor play (31), Sport (9), Fall (8), Traffic accident (1.5), Chewing (1), Unknown (18)		

Huang et al [17], Taiwan, 2009	6312 15- to 18-year-old senior high school	19.9%	M (12.4%) F (7.5%)	Sports(30.8), Eating (20.5), Falls (19.4), Traffic Accidents (10.2), Collisions (7.1), other (12)	Home (29.1%), school (23.7%), street (11.1%)	
Kovács et al [48], Romania., 2012	4638 -children and adolescents aged between 1 and 18 years	24.8%	M (14.6%) F (10.2%)	Fall (42.5), Sports(23.5), accidents (9.5), violence (7.5),Collision (5.5), Other (6.5),Unknown(4.7)		Uncomplicated coronal fracture (26.1), Enamel fracture (12.5), Lateral luxation (11.2), Contusion (7.7) , other (42.5)
Govindarajan et al [49], India, 2012	3200 school children in the age 3-13 years	10.13	M (6.03%) F (4.1%)	Fall (41.9), Sports(19.1), accidents (8.64), violence (5.8), Collision (.3), Unknown(24.07)		
Dua and Sharma [50], India, 2012	880 children in the age 7-12 years	14.5%				Enamel fracture (50%), crown fracture without pulpal involvement (20.3), crown fracture with pulpal involvement (13.2), avulsion (4.6), fracture unspecified(11.9)
Abdulaziz and et al[51], 2010, Kuwait	500 preschool children(2-6 years)	11.2%		Fall (94.6%),	Home (87.5%), street (7.1%), school (5.4%).	Enamel fractures (29.4), enamel- dentin fractures (26.5), crown fractures (14.7),
Malikaew et al [52], Thailand, 2006	2,725 of 11-13 year old Thai children.	35.0%	M (22.1%) F (12.9%)	Fall (24.8%), Collision(21.1), Traffic accident(1), Misuse of teeth(18.7), Unknown(21.7) , other (12.7)	Homes (31.7), schools (28.0), street (4.7), swimming pool (1.2), other (10.7)	Enamel fracture (83), enamel- dentin fractures (4.8)
-Hamdan and Rock[53], Jordan, 1995	459 schoolchildren aged 10-12 years	17.3	M (10.4%) F (6.9%)			
Vejdani. And Mohammad Alizadeh, Iran [54]; 2006	700 fifth and sixth grade schoolchildren	10.3	M (5.4%) F (4.9%)	Falls (37.5), Collision (22.2), Traffic accident (4.2), Sports (19.4), other (17.7),		Enamel fracture(76.4), Enamel and dentine fracture(19.4) Enamel, dentin and exposure of the pulp (4.2)
Vejdani et al [55], Iran; 2011	1042 fifth and sixth grade schoolchildren	15.2	M (9.6%) F (5.6%)			Enamel fracture (8.3) , Enamel and dentine fracture (16.9) Enamel, dentin and exposure of the pulp (2.1)
Sgan-Cohen et al [56],Israel, 2005	1195 fifth and sixth grade schoolchildren	29.6%		Falls (30.3), violence (30.3), Sports (27.3)	Homes (31.4), schools (32.2), street/ outside (36.4)	
Nicolau et al [43], Brazil, 2001	652 ,13-year-old adolescents	20.4%	M (13.7%) F (6.7%)	Falls (24.1) collisions (15), traffic accidents (10.5), misuse of the teeth (6), sports (2.3) violence (1.5).		
Rajab [57], Jordan,2003	2751 children	14.2%	M (9.2%) F (5%)	Falls (49.9) collisions (30), sports (8.7) violence (7.2). traffic accidents (3)	Homes (63.17), schools (25.5), street (11.25)	Enamel and dentine fracture (43.8), Enamel, dentin and exposure of the pulp (37.5), Enamel fracture (14.6)
Soriano et al [58], Brazil, 2007	1046 boys and girls aged 12	10.5%		Falls (27.3) collisions (18.2), sports (8.2) violence (6.4). traffic accidents (2.7), Non-specified accidents (22.7) other (14.5)	Homes (25.5), schools (13.6), street (23.6), Do not remember (20.9), other(16.4)	Enamel fracture(47.3), Enamel and dentin fracture without pulp exposure (34.5), Crown fracture without pulp exposure (10), other(8.2)
Sgan-Cohen et al [59], Israel, 2008	453 5th and 6th grade	33.8%		Falls (29.1), sports (16.4) violence		

David et al [60], South India,2009	838 schoolchildren 12-year-old school children	6.1%		(20), playing (20)
Wendt et al [61], Brazil, 2010	571 preschool children	36.6%	M (19.7%) F (16.9%)	
Thelen and Ba^rdsen, Albania[62], 2010	2789, adolescents aged 16–18 years	9.9%	M (5.8%) F (4.1%)	Collision (27.5), Falls (13.4), sports (14.1) violence (4.0), Traffic accidents ( 4.7), Do not remember( 10.5), other(25.8)
Livny et al [63], Israel, 2010	804 sixth grade Children of ages 11 and 12 years.	17.7%	M (11.1%) F (6.6%)	

studies, the best estimation of the prevalence of trauma was determined based on the random effect model (Fig. 4).

The prevalence of dental trauma in children and adolescents based on the random effect was determined to be 17.5% (95% CI: 14.6%-20.4 %); 95% CI for the prevalence was drawn for each study in the horizontal line format ( $Q= 3631.17$ ,  $df= 43$ ,  $p<0.001$ ).

The frequency distribution for dental trauma based on the location variable (continent) is shown in Figs. 5-7.

To assess the publication bias, funnel plot was used (Fig. 8). The results of funnel plot revealed that there was publication bias among studies. The shape of funnel plots was not symmetrical.

### Discussion

Dental trauma is a major health problem in many societies (26,29,30), with higher prevalence rates among children and adolescents (25,26,29,48,64). The present study revealed the prevalence of dental trauma in children and adolescents (under 18 years of age) to be 17.5%, but with variances among different geographic regions.

In the present study, the prevalence of dental traumas for all subjects was determined to be 17.5%, which necessitates efficient plan-

ning and intervention to prevent their incidence. Moreover, considering the negative influence of dental traumas on individuals' quality of life (39,65), psychological and social problems (66,67), and direct and indirect costs of treatment (68), developing effective interventional strategies to deal with this issue is of prime importance. The difference in the prevalence of dental trauma among the studied geographic regions might be a reflection of their different cultural, sanitary, and economic conditions, type of activities and professions practiced, and ecological characteristics like the fluoride concentration of drinking tap water. For instance, the prevalence of dental trauma was found to be higher in the continent of America compared to Asia or Europe, and this could be the result of sports activities of youth, as most studies in the region pertained to Brazil. Furthermore, the fluoride concentration of water in regions with mild and humid weather condition is usually low (69), resulting in lower fluoride content in teeth which can render teeth more prone to dental trauma. This notion highlights the importance of local and regional considerations during planning and interventions by policy makers.




Fig. 4. The prevalence of dental trauma in children and adolescents based on the random effect model.

The prevalence of dental trauma was higher in boys compared to girls ( $10.2 \pm 4.7$  vs.  $6.5 \pm 3.4$ ). Previous studies also agree on the higher prevalence of dental trauma in boys than girls (27,51,70-72), which might be due to boys performing more activities like sports, bicycle riding, fighting, working, etc. compared to girls (73).

In the present study, the most important

cause for dental trauma was falling, which is in agreement with previous literature (74-78). This can be explained by the imbalance of the newly walking children leading to falling accidents. Thus, children should be cared more prudently, reinforced by measures to eliminate the risk of falling. The other important cause of dental trauma in this study was sports, which is in line






Fig.5. The prevalence of dental trauma in children and adolescents in the continent of America was estimated based on the random effect of 21.2% (95% CI:16.6%-26.7%); 95% CI was drawn for each prevalence in the horizontal line format ( $I^2 = 98.7$ ).




Fig. 6. The prevalence of dental trauma in children and adolescents in Europe was estimated based on the random effect of 19.1% (95% CI:13.7%-24.1%); 95% CI was drawn for each prevalence in the horizontal line format ( $I^2 = 97.9$ ).




Fig. 7. The prevalence of dental trauma in children and adolescents in Asia was estimated based on the random effect of 14.7% (95% CI: 11.3%-19.0 %); 95% CI was drawn for each prevalence in the horizontal line format ( $I^2 = 98.4$ ).

with previous research (79,80). Ensuring the existence of sound safety measures in sports venues accompanied by the use of protective athletic appliances such as mouth guards, and educating children and adolescents who engage in sports activities are among suggestions that can be effective in reducing the incidence of dental trauma in this age range (81,82). Fights and violence were also among the causes of dental trauma in children and adolescents in the present study, which is indicative of the behavioral characteristics of this age range, with dominant, violent, emotional, and illogical traits. In this regard, special attention should be given to social education at home and school, fostering a culture of self-control and social respect towards others.

The findings of this study revealed that dental traumatic accidents occurred most frequently at home, and this is in line with many previous studies (83-85). This is explainable by the fact that children and adolescents spend a significant part of their time at home; and therefore, special safety measures should be considered at houses in which families live with their children.

Based on the analysis of the study data, enamel fracture was the most frequent type of dental traumatic accidents, confirming the findings of other studies (34,42,56,86). Therefore, dentists and other health-care professionals as well as lay people should have adequate knowledge regarding the management of dental traumatic events (87-90).

The analyses performed in the present study were limited by age under 18 years. Thus, it is suggested to include adults in future studies. The study was also limited by the fact that it only included articles in English and Persian and only included studies conducted since 1995.

### Conclusion

The results of this study indicated a significant prevalence for dental traumatic accidents among children and adolescents, which necessitates efficient planning and




Fig 8. Funnel plot of standard error by event rate

interventions in order to prevent their incidence. Planning for reducing the risk of falling in children would be of utmost importance. Safety measures would be most effective when implemented at home where dental traumatic accidents occur most frequently.

## References

1. Leung YY, Cheung LK. Can coronectomy of wisdom teeth reduce the incidence of inferior dental nerve injury? *Ann R Australas Coll Dent Surg* 2008;19:50-1.
2. Vanderas AP, Papagiannoulis L. Urinary catecholamine levels and incidence of dentofacial injuries in children: a 2-year prospective study. *Endod Dent Traumatol* 2000;16(5):222-8.
3. Bemelmans P, Pfeiffer P. [Incidence of dental, mouth, and jaw injuries and the efficacy of mouthguards in top ranking athletes]. *Sportverletz Sportschaden* 2000;14(4):139-43.
4. Vanderas AP, Papagiannoulis L. Incidence of dentofacial injuries in children: a 2-year longitudinal study. *Endod Dent Traumatol*. 1999;15(5):235-8.
5. Panagakos FS, Silverstein J. Incidence of percutaneous injuries at a dental school: a 4-year retrospective study. *Am J Infect Control* 1997;25(4):330-4.
6. Hamilton FA, Hill FJ, Holloway PJ. An investigation of dento-alveolar trauma and its treatment in an adolescent population. Part 1: The prevalence and incidence of injuries and the extent and adequacy of treatment received. *Br Dent J* 1997;182(3):91-5.
7. Hardt N, Fellmann W. [Nerve injuries during dental and orthodontic interventions. Their causes, incidence, sequelae and legal assessment]. *Schweiz Monatsschr Zahnmed* 1996;106(1):31-44.
8. Luz JG, Di Mase F. Incidence of dentoalveolar injuries in hospital emergency room patients. *Endod Dent Traumatol* 1994;10(4):188-90.
9. Schatz JP, Hausherr C, Lang R, Joho JP. [Dental and dentoalveolar injuries: their etiology, incidence and distribution in a specialized university service]. *Schweiz Monatsschr Zahnmed* 1994; 104(7):843-7.
10. Bolhuis JH, Baarda DB, Leurs JM, Fogel GE. [Incidence and prevention of dental injuries in sports hockey]. *Ned Tijdschr Tandheelkd* 1986;93(7-8):283-7.
11. Sassen H. [Incidence of clinically manifest functional disorders in partial dentition injury]. *Dtsch Zahnärztl Z* 1982;37(12):969-74.
12. Schutzmansky G. [Statistics on the incidence and severity degree of the accidental injuries on the corona dentis in the frontal teeth of children and adolescents. Examination material of the adolescent dental clinic of the town and district Halle (Saade)]. *Z Gesamte Hyg* 1970;16(2):133-5.
13. Marchiori EC, Santos SE, Asprino L, de Moraes M, Moreira RW. Occurrence of dental avulsion and associated injuries in patients with facial trauma over a 9-year period. *Oral Maxillofac Surg* 2012;7:7.
14. Garbin CA, Guimaraese Queiroz AP, Rovida TA, Garbin AJ. Occurrence of traumatic dental injury in cases of domestic violence. *Braz Dent J* 2012;23(1):72-6.
15. Traebert J. Accidents, sports, and physical leisure activities are the most frequent causes of traumatic dental injury and the rate of pulp necrosis is high following its occurrence in Pilsen, The Czech Republic. *J Evid Based Dent Pract* 2011; 11(2):102-4.
16. Thoren H, Numminen L, Snall J, Kormi E, Lindqvist C, Iizuka T, et al. Occurrence and types of dental injuries among patients with maxillofacial fractures. *Int J Oral Maxillofac Surg* 2010; 39(8):774-8.
17. Huang B, Marcenes W, Croucher R, Hector M. Activities related to the occurrence of traumatic dental injuries in 15- to 18-year-olds. *Dent Traumatol* 2009;25(1):64-8.
18. Harry GJ, Lefebvre d'Hellencourt C. Dentate gyrus: alterations that occur with hippocampal

injury. *Neurotoxicology* 2003;24(3):343-56.

19. Andreasen FM, Zhijie Y, Thomsen BL, Andersen PK. Occurrence of pulp canal obliteration after luxation injuries in the permanent dentition. *Endod Dent Traumatol* 1987;3(3):103-15.

20. Maidwell-Smith MA. The role of the dental surgeon in a case of suspected non-accidental injury occurring in a child. Its aetiology, recognition and management. *Apex* 1980;12(1):11-2.

21. Jarvinen S. [Studies on causes and occurrence of dental injuries in children]. *Proc Finn Dent Soc* 1972;68(1):27-31.

22. Krishna A, Malur MH, Swapna DV, Benjamin S, Deepak CA. Traumatic dental injury-an enigma for adolescents: a series of case reports. *Case Rep Dent* 2012;756526(10):7.

23. Wennervaldt K, Melchior J. Risk of perforation using rigid oesophagoscopy in the distal part of oesophagus. *Dan Med J* 2012;59(11).

24. Jabbar NS, Aldrigui JM, Braga MM, Wanderley MT. Pulp polyp in traumatized primary teeth - a case-control study. *Dent Traumatol* 2012;20(10):12019.

25. Schuch HS, Goettens ML, Correa MB, Torriani DD, Demarco FF. Prevalence and treatment demand after traumatic dental injury in South Brazilian schoolchildren. *Dent Traumatol* 2012; 16(10):12003.

26. Martins VM, Sousa RV, Rocha ES, Leite RB, Paiva SM, Granville-Garcia AF. Dental trauma among Brazilian schoolchildren: prevalence, treatment and associated factors. *Eur Arch Paediatr Dent* 2012;13(5):232-7.

27. Dame-Teixeira N, Alves LS, Susin C, Maltz M. Traumatic dental injury among 12-year-old South Brazilian schoolchildren: prevalence, severity, and risk indicators. *Dent Traumatol* 2012;27(10):1600-9657.

28. Dame-Teixeira N, Alves LS, Susin C, Maltz M. Traumatic dental injury among 12-year-old South Brazilian schoolchildren: prevalence, severity, and risk indicators. *Dent Traumatol* 2013;29(1):52-8.

29. Piovesan C, Abella C, Ardenghi TM. Child oral health-related quality of life and socioeconomic factors associated with traumatic dental injuries in schoolchildren. *Oral Health Prev Dent* 2011; 9(4):405-11.

30. Jorge KO, Moyses SJ, Ferreira E, Ramos-Jorge ML, de Araujo Zarzar PM. Prevalence and factors associated to dental trauma in infants 1-3 years of age. *Dent Traumatol* 2009;25(2):185-9.

31. Aldrigui JM, Abanto J, Carvalho TS, Mendes FM, Wanderley MT, Bönecker M, et al. Impact of traumatic dental injuries and malocclusions on quality of life of young children. *Health Qual Life Outcomes* 2011; 9(78):1477-7525.

32. Norton E, O'Connell AC. Traumatic dental injuries and their association with malocclusion in the primary dentition of Irish children. *Dent*

*Traumatol* 2012;28(1):81-6.

33. Bendo CB, Paiva SM, Oliveira AC, Goursand D, Torres CS, Pordeus IA, et al. Prevalence and associated factors of traumatic dental injuries in Brazilian schoolchildren. *J Public Health Dent* 2010;70(4): 313-8.

34. Diaz JA, Bustos L, Brandt AC, Fernandez BE. Dental injuries among children and adolescents aged 1-15 years attending to public hospital intemuco, Chile. *Dent Traumatol* 2010; 26:254-261.

35. Altun C, Ozen B, Esenlik E, Guven G, Gürbüz T, Acikel C, et al. Traumatic injuries to permanent teeth in Turkish children, Ankara. *Dental Traumatology* 2009;25:309-313.

36. Noori AJ, Al-Obaidi WA. Traumatic dental injuries among primary school children in Sulaimani city, Iraq. *Dent Traumatol* 2009;25(4):442-6.

37. Avsar A, Akbas S, Ataibis T. Traumatic dental injuries in children with attention deficit/hyperactivity disorder. *Dent Traumatol* 2009; 25(5):484-9.

38. Naidoo S, Sheiham A, Tsakos G. Traumatic dental injuries of permanent incisors in 11 to 13-year-old South African schoolchildren. *Dent Traumatol* 2009;25(2):224-8.

39. Fakhruddin KS, Lawrence HP, Kenny DJ, Locker D. Impact of treated and untreated dental injuries on the quality of life of Ontario school children. *Dent Traumatol* 2008;24:309-313.

40. Pattussi MP, Hardy R, Sheiham A. Neighborhood Social Capital and Dental Injuries in Brazilian Adolescents. *Am J Public Health* 2006;96: 1462-1468.

41. Locker D. Prevalence of traumatic dental injury in grade 8 children in six Ontario communities. *Can J Public Health* 2005;96(1):73-6.

42. Traebert J, Bittencourt DD, Peres KG, Peres MA, de Lacerda JT, Marcenes W. Aetiology and rates of treatment of traumatic dental injuries among 12-year-old school children in a town in Southern Brazil. *Dent Traumatol* 2006;22:173-8.

43. Nicolau B, Marcenes W, Sheiham A. Prevalence, causes and correlates of traumatic dental injuries among 13-year-olds in Brazil. *Dent Traumatol* 2001;17(5):213-7.

44. Cortes M, Marcenes W, Sheiham A. Prevalence and correlates of traumatic injuries to the permanent teeth of schoolchildren aged 9-14 years in Belo Horizonte, Brazil. *Dent Traumatol* 2001;17: 22-26.

45. Marcenes W, Al Beiruti N, Tayfour D, Issa S. Epidemiology of traumatic injuries to the permanent incisors of 9-12-year-old schoolchildren in Damascus, Syria. *Endod Dent Traumatol* 1999;15: 117-123.

46. Chen YL, Tsai TP, See LC. Survey on incisor Trauma in Second Grade Students Of Central Taiwan. *Chang Gung Medical Journal* 1999;22: 212-219.

47. Petti S, Cairella G, Tarsitani G. Childhood

obesity: a risk factor for traumatic injuries to anterior teeth. *Endod Dent Traumatol* 1997;13:285-288.

48. Kovacs M, Pacurar M, Petcu B, Bukhari C. Prevalence of Traumatic Dental Injuries in Children Who Attended Two Dental Clinics in TA centrgru MureA masculine Between 2003 and 2011. *Oral Health Dent Manag* 2012;11(3):116-24.

49. Govindarajan M, Reddy VN, Ramalingam K, Durai KS, Arun Rao P, Prabhu A. Prevalence of traumatic dental injuries to the anterior teeth among three to thirteen-year-old school children of Tamilnadu. *Contemp Clin Dent* 2012;3(2):164-7.

50. Dua R, Sharma S. Prevalence, causes, and correlates of traumatic dental injuries among seven-to-twelve-year-old school children in Dera Bassi. *Contemp Clin Dent* 2012;3(1):38-41.

51. Hasan AA, Qudeimat MA, Andersson L. Prevalence of traumatic dental injuries in preschool children in Kuwait - a screening study. *Dent Traumatol* 2010;26(4):346-50.

52. Malikaew P, Watt RG, Sheiham A. Prevalence and factors associated with traumatic dental injuries (TDI) to anterior teeth of 11-13 year old Thai children. *Community Dent Health* 2006;23(4):222-7.

53. Hamdan MA, Rock WP. A study comparing the prevalence and distribution of traumatic dental injuries among 10-12-year-old children in an urban and in a rural area of Jordan. *Int J Paediatr Dent* 1995;5(4):237-41.

54. Vejdani J, Mohammad Alizadeh N. The Prevalence and Etiology of Anterior Crown Fracture of Anterior Permanent Teeth. *Journal of Medical Faculty Guilan University of Medical Sciences* 2006;15(58):87-92 [In Persian].

55. Vejdani J, Bahramnejhad E, Rezaie M. Prevalence and Etiology of Anterior Permanent Teeth Crown Fracture in Elementary Students in Rasht in 2007. *JRDS* 2011;8(1):15-19 [In Persian].

56. Sgan-Cohen HD, Megnagi G, Jacobi Y. Dental trauma and its association with anatomic, behavioral, and social variables among fifth and sixth grade schoolchildren in Jerusalem. *Community Dent Oral Epidemiol* 2005;33:174-80.

57. Rajab LD. Traumatic dental injuries in children presenting for treatment at the Department of Pediatric Dentistry, Faculty of Dentistry, University of Jordan, 1997-2000. *Dent Traumatol* 2003;19(1):6-11.

58. Soriano EP, Caldas Jr AF, Go'és PSA. Prevalence and risk factors related to traumatic dental injuries in Brazilian schoolchildren. *Dent Traumatol* 2007;23(4):232-40.

59. Sgan-Cohen HD, Yassin H, Livny A. Dental trauma among 5th and 6th grade Arab schoolchildren in Eastern Jerusalem. *Dent Traumatol* 2008;24(4):458-61.

60. David J, Astrom AN, Wang NJ. Factors associated with traumatic dental injuries among 12-

year-old schoolchildren in South India. *Dent Traumatol* 2009;25(5):500-5.

61. Wendt FP, Torriani DD, Assunção MC, Romano AR, Bonow ML, da Costa CT, et al. Traumatic dental injuries in primary dentition: epidemiological study among preschool children in South Brazil. *Dent Traumatol* 2010;26(2):168-73.

62. Thelen DS, Bardsen A. Traumatic dental injuries in an urban adolescent population in Tirana, Albania. *Dent Traumatol* 2010;26(5):376-82.

63. Livny A, Sgan-Cohen HD, Junadi S, Marcenes W. Traumatic dental injuries and related factors among sixth grade schoolchildren in four Palestinian towns. *Dent Traumatol* 2010;26(5):422-6.

64. Kumar A, Bansal V, Veerasha KL, Sogi GM. Prevalence of traumatic dental injuries among 12- to 15-year-old schoolchildren in Ambala district, Haryana, India. *Oral Health Prev Dent* 2011;9(3):301-5.

65. Ramos-Jorge ML, Bosco VL, Peres MA, Nunes AC. The impact of treatment of dental trauma on the quality of life of adolescents – a case-control study in southern Brazil. *Dent Traumatol* 2007;23:114-119.

66. Marcenes W, Murray S. Social deprivation and traumatic dental injuries among 14-year-old schoolchildren in Newham, London. *Dent Traumatol* 2001;17:17-21.

67. McGrath C, Broder H, Wilson-Genderson M. Assessing the impact of oral health on the quality of life of children: implications for research and practice. *Commun Dent Oral Epidemiol* 2004;32:81-5.

68. Glendor U, Jonsson D, Halling A, Lindqvist K. Direct and indirect costs of dental trauma in Sweden: a 2-year prospective study of children and adolescents. *Community Dent Oral Epidemiol* 2001;29:150-60.

69. Azami-Aghdash S, Ghofazadeh M, Pournaghi Azar F, Naghavi-Behzad M, Mahmoudi M, Jamali Z. Fluoride Concentration of Drinking Waters and Prevalence of Fluorosis in Iran: A Systematic Review. *Dental Research, Dental Clinics, Dental Prospects* 2013;7(1):1-7.

70. de Vasconcelos Cunha Bonini GA, Marcenes W, Oliveira LB, Sheiham A, Bonecker M. Trends in the prevalence of traumatic dental injuries in Brazilian preschool children. *Dent Traumatol* 2009;25(6):594-8.

71. Jagger RG, Abbasbhai A, Patel D, Jagger DC, Griffiths A. The prevalence of dental, facial and head injuries sustained by schoolboy rugby players. A pilot study. *Prim Dent Care* 2010;17(3):143-6.

72. Elhammali N, Bremerich A, Rustemeyer J. Demographical and clinical aspects of sports-related maxillofacial and skull base fractures in hospitalized patients. *Int J Oral Maxillofac Surg* 2010;39(9):857-62.

73. Rocha MJ, Cardoso M. Traumatized permanent teeth in Brazilian children assisted at the

Federal University of Santa Catarina, Brazil. *Dent Traumatol* 2001;17:245-9.

74. Zhou HH, Ongodia D, Liu Q, Yang RT, Li ZB. Dental trauma in patients with maxillofacial fractures. *Dent Traumatol* 2012;12(10):1600-9657.

75. Gupta S, Kumar-Jindal S, Bansal M, Singla A. Prevalence of traumatic dental injuries and role of incisal overjet and inadequate lip coverage as risk factors among 4-15 years old government school children in Baddi-Barotiwala Area, Himachal Pradesh, India. *Med Oral Patol Oral Cir Bucal* 2011;16(7):e960-5.

76. Ravishankar TL, Kumar MA, Ramesh N, Chaitra TR. Prevalence of traumatic dental injuries to permanent incisors among 12-year-old school children in Davangere, South India. *Chin J Dent Res* 2010;13(1):57-60.

77. Diaz JA, Bustos L, Brandt AC, Fernandez BE. Dental injuries among children and adolescents aged 1-15 years attending to public hospital in Temuco, Chile. *Dent Traumatol* 2010;26(3):254-61.

78. Lin H, Naidoo S. Causes and prevalence of traumatic injuries to the permanent incisors of school children aged 10-14 years in Maseru, Lesotho. *Sadj* 2008;63(3):154-6.

79. Kumamoto DP, Maeda Y. A literature review of sports-related orofacial trauma. *Gen Dent* 2004; 52(3):270-280.

80. Shirani G, Kalantar Motamedi MH, Ashuri A. Prevalence and patterns of combat sport related maxillofacial injuries. *J Emerg Trauma Shock* 2010; 3(4):314-317.

81. Labella CR, Smith BW, Sigurdsson A. Effect of mouthguards on dental injuries and concussions in college basketball. *Med Sci Sports Exerc* 2002; 34(1):41-44.

82. Winters Sr JE. Commentary: role of properly fitted mouthguards in prevention of sport-related concussion. *J Athl Train* 2001; 36(3):339-341.

83. Garcia-Godoy F, Sanchez R, Sanchez JR. Traumatic dental injuries in a sample of dominican schoolchildren. *Community Dent Oral Epidemiol* 1981;9:193-7.

84. Onetto JE, Flores MT, Garbarino ML. Dental trauma in children and adolescents in Valparaiso, Chile. *Endod Dent Traumatol* 1994;10:223-7.

85. Blinkhorn FA. The etiology of dento-alveolar injuries and factors influencing attendance for emergency care of adolescents in the north west of England. *Endod Dent Traumatol* 2000;16:162-5.

86. Tovo MF, dos Santos PR, Kramer PF, Feldens CA, Sari GT. Prevalence of crown fractures in 8-10 years old schoolchildren in Canoas, Brazil. *Dent Traumatol* 2004;20:251-4.

87. Cohenca N, Forrest JL, Rostein I. Knowledge of oral health professionals of treatment of avulsed teeth. *Dent Traumatol* 2006; 22(6):296-301.

88. Diaz J, Bustos L, Herrera S, Herrera S. Knowledge of the management of paediatric dental traumas by non-dental professionals in emergency rooms in South Araucania, Temuco, Chile. *Dent Traumatol* 2009; 25(6):611-619.

89. Skeie MS, Audestad E, Bardsen A. Traumatic dental injuries-knowledge and awareness among present and prospective teachers in selected urban and rural areas of Norway. *Dent Traumatol* 2010; 26(3):243-247.

90. Abu-Dawoud M, Al-Enezi B, Andersson L. Knowledge of emergency management of avulsed teeth among young physicians and dentists. *Dent Traumatol* 2007;23(6):348-355.