



Assessment of length of stay in a general surgical unit using a zero-inflated generalized Poisson regression

Roghayeh Farhadi Hassankiadeh¹, Anoshirvan Kazemnejad^{1*}, Mohammad Gholami Fesharaki¹, Siamak Kargar Jahromi², Nasim Vahabi¹

Received: 30 Nov 2016 Published: 17 Dec 2017

Abstract

Background: The effective use of limited health care resources is of prime importance. Assessing the length of stay (LOS) is especially important in organizing hospital services and health system. This study was conducted to identify predictors of LOS among patients who were admitted to a general surgical unit.

Methods: In this cross-sectional study, the sample included all patients who were admitted to the general surgical unit of Shariati hospital in 2013 (n= 334). To determine the factors affecting LOS, Zero-inflated Poisson (ZIP), zero-inflated negative binomial (ZINB), and zero-inflated generalized Poisson (ZIGP) regression models were fitted using R software, and then the best model was selected.

Results: Among all 334 patients, the mean (\pm SD) age of the patients was 45.2 (\pm 16.47) years and 220 (65.9%) of them were male. The results revealed that based on ZIGP model, type of surgery (appendicitis, abdomen and its contents, hemorrhoids, lung, and skin), type of insurance, comorbid diseases (hypertension, heart disease, and hyperlipidemia), place of residence (local and non-local), age, and number of tests had significant effects on the LOS of GS patients.

Conclusion: According to the Akaike information criterion (AIC) in each fitted model, it was found that ZIGP regression model is more appropriate than ZIP and ZINB regression models in assessing LOS in GS patients, especially due to the presence of excess zeros and overdispersion in count data.

Keywords: Length of stay, General surgical unit, Zero-inflated generalized Poisson regression

Copyright© Iran University of Medical Sciences

Cite this article as: Farhadi Hassankiadeh R, Kazemnejad A, Gholami Fesharaki M, Kargar Jahromi S, Vahabi N. Assessment of length of stay in a general surgical unit using a zero-inflated generalized Poisson regression. *Med J Islam Repub Iran.* 2017 (17 Dec);31:91. <https://doi.org/10.14196/mjiri.31.91>

Introduction

One of the main objectives of applied and medical statistics is modeling data to analyze and explain medical phenomena. Modeling of length of stay (LOS) is an efficient way to determine the factors that affect LOS. Moreover, LOS is one of the simplest indicators of hospital activity used for various purposes including hospital care management, quality control, appropriate use of hospital services, hospital planning, determining efficiency, and using hospital resources (1-4). Furthermore, many studies have identified LOS as a risk factor for 30-day readmission (5-7). Lim et al. collected effective factors on LOS for patients who died from 2000 to 2003. The results revealed that the older the patients the longer the LOS will

be. Old patients with chronic diseases had long LOS, while young patients with mainly acute disease had a short one (8). Lobato et al. identified prolonged-stay risk factors after colorectal surgery (the colon and rectum). They found that congestive heart failure, high levels of albumin, and anemia are 3 contributing factors that prolonged LOS after colorectal surgery (9). To determine the associated factors with length of stay, Huang JQ studied 2757 adults' data who had been admitted during 2 years. Patients' age, functional status, comorbid disease, and weight were considered effective factors in LOS (10). Arab et al., in a study in Lorestan public hospitals, found that age, gender, marital status, place of residence, job,

Corresponding author: Dr Anoshirvan Kazemnejad, kazem_an@modares.ac.ir

¹ Department of Biostatistics, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran.

² Shariati Hospital, Medical Education Research Center, Tehran, Iran.

↑What is "already known" in this topic:

LOS is one of the simplest indicators of hospital activity, and many studies have identified LOS as a risk factor for 30-day readmission.

→What this article adds:

We used GLM to identify the effective factors on LOS in a general surgical unit.

type of referral, type of insurance, causes of hospitalization, and patient condition on discharge affect the average length of stay. LOS is also longer among males than females. Recovered or relatively recovered patients on discharge have shorter LOS (11). Hosseini et al. conducted an analytic cross-sectional study on a sample of 570 hospitalized patients for appendicitis in 25 provinces and 50 social security hospitals in Iran in March 21st, 2009 to September 22nd, 2009. The recorded age, gender, type of insurance, and hospital stay variables for each single patient were obtained from the hospitals' information management system. The findings revealed that the number of active beds, gender, age, and type of insurance did not affect the increase or decrease of LOS (12).

Because the data of LOS are count data, statistical techniques based on the normal distribution are not appropriate, instead generalized linear models including negative binomial regression or Poisson regression should be used (13-17).

Mean and variance equality of response variable is the main condition for using Poisson model. If the condition is not fulfilled, then, generalized Poisson and negative binomial models are appropriate (18-20).

Generalized Poisson regression and negative binomial regression models are, to some extent, capable of determining dispersion. When the zero data are exceeded, data such as LOS of these models will not be efficient. These methods cannot be used to explain and analyze overdispersion in LOS data, so one of the proper approaches to analyze them is zero-inflated regression models (21). In this method, response distribution consists of 2 components: zero and non-zero, and these models depend on the non-zero component. For example, if non-zero Poisson is considered, then, it is called ZIP (Zero Inflated Poisson). However, if negative binomial distribution is taken into account, it is called ZINB (Zero Inflated Negative Binomial), and if a generalized Poisson distribution is considered, it is called ZIGP (Zero Inflated Generalized Poisson).

Considering the importance of LOS for surgical patients, this study was conducted to determine the factors affecting the LOS of general surgical patients using zero-inflated regression models (ZIP, ZINB and ZIGP).

Methods

Data

Table 1. Frequency Distribution of the General Surgical Patients

Variable	Category	Number	%
Gender	Female	114	34.1
	Male	220	65.9
Type of insurance	Have insurance	278	83.2
	Do not have Insurance	56	16.8
Place of residence	Non-local	71	21.3
	Local	263	78.7
Age (year)	< 14	6	1.8
	14-44	152	45.5
	45-65	136	40.7
	> 65	40	12
length of stay (day)	0	179	53.6
	1	62	18.6
	2	27	8
	≥3	66	19.8

A cross-sectional study was conducted and data were collected from patients admitted to the general surgical unit of Shariati hospital during March 21st, 2013 and June 21st, 2013. The number of records was reported to be 334. The features consisted of type of surgery, which included appendicitis, digestive disease, abdomen and its contents, skeletal and muscle system, blood and lymphatic system, glands, urinary system, hemorrhoids, cardiovascular system, female reproductive system, breast, lung, and skin. This classification is based on California Tariff. Other features included number of surgery, type of insurance, history of smoking, pulse, different types of diseases (hypertension, diabetes, hyperlipidemia, kidney failure, and other comorbid diseases), blood hemoglobin levels, age, gender, number of tests, and type of admission.

Statistical Analysis

In this study, many patients were hospitalized just for one day, which resulted in excess zeros in LOS data, so the zero-inflated models were proper choices.

The general form of the ZIP model is as follows:

$$Y_i = \begin{cases} 0 & \text{with probability } p_i \\ \text{Poisson}(\lambda_i) & \text{with probability } (1 - p_i) \end{cases}$$

In which the parameters $P = (p_1, \dots, p_N)^T$ and $\lambda = (\lambda_1, \dots, \lambda_N)^T$ are modeled via canonical link in GLMs family as $\log(\lambda) = B\beta$ and $\text{logit } p = G\gamma$ to design matrices B and G.

The probability mass function is as follows:

$$P(Y_i = y_i) = \begin{cases} p_i + (1 - p_i)e^{-\lambda_i} & x = 0 \\ (1 - p_i) e^{-\lambda_i} \lambda_i^{y_i} / y_i! & x = 1, 2, \dots \end{cases}$$

where $0 < p < 1$ and $\lambda_i \geq 0$.

Three models including zero-inflated Poisson, zero-inflated negative binomials, and zero-inflated generalized Poisson were fitted to data. Then, these models were compared using the AIC (smaller AIC means better fit). Data were analyzed using R software and $p < 0.05$ was considered as statistically significant.

Results

Most patients were local males aged 14 to 44 years with

Table 2. The Fitness Of Zero-Inflated Poisson and Zero-Inflated Negative Binominals Regression Models

Parameter (category)	ZIP		ZINB	
	Estimation	p	Estimation	p
Fixed value	0.14	0.83	0.24	0.838
Appendicitis (No*/Yes)	-3.71	<0.001	-3.80	<0.001
Digestive disease (No*/Yes)	0.0992	0.65	0.02	0.95
Abdomen and its Contents (No*/Yes)	-0.79	<0.001	-0.78	0.023
Skeletal and muscle system (No*/Yes)	0.15	0.604	.004	0.99
Blood and lymphatic system (No*/Yes)	-0.11	0.74	-0.16	0.747
Glands (No*/Yes)	0.38	0.101	0.38	0.317
Urinary system (No*/Yes)	0.05	0.854	-0.12	0.802
Hemorrhoids (No*/Yes)	-2.16	0.001	-1.97	0.003
Cardiovascular system (No*/Yes)	-0.03	0.91	-0.14	0.772
Female reproductive system (No*/Yes)	0.58	0.175	0.39	0.569
Breast (No*/Yes)	-0.62	0.292	-0.77	0.275
Lung (No*/Yes)	-1.54	0.004	-1.69	0.047
Skin (No*/Yes)	-2.24	0.004	-2.08	0.041
Type of insurance (No*/Yes)	0.54	0.001	0.01	0.047
History of smoking (Non-smoker*/Smoker)	0.18	0.187	0.28	0.242
Hypertension (No*/Yes)	0.53	0.000	0.55	.019
Diabetes (No*/Yes)	0.04	0.763	-0.14	0.593
Heart disease (No*/Yes)	0.43	0.006	0.41	0.049
Hyperlipidemia (No*/Yes)	0.61	0.001	0.39	0.062
Kidney disease (No*/Yes)	-0.01	0.963	-0.042	0.905
Comorbid disease (No*/Yes)	-0.01	0.957	-0.09	0.709
Hemoglobin (Not natural*/Natural)	-0.06	0.551	-0.10	0.60
Gender (Female*/male)	0.04	0.751	0.01	0.944
Marital status (Single*/Married)	0.18	0.21	0.19	0.38
Place of residence (Non-local*/Local)	-0.51	0.000	-0.64	0.001
Type of admission (Low risk*/Medium Risk)	0.03	0.872	0.173	0.992
Type of admission (Low risk*/High Risk)	0.58	0.023	0.066	0.90
Type of admission (Emergency/ hospitalization)	-0.26	0.029	-0.29	0.162
Number of tests	0.17	0.000	.1589	0.000
Age	0.24	.0424	0.25	.0456
Number of surgery	0.06	0.231	0.143	0.120
Pulse	-0.01	0.415	-0.001	0.965
AIC		1184		1072

*Reference group

insurance. Table 1 displays demographic characteristics. From the perspective of model selection, AIC indicated that the ZIGP model (AIC= 1070) offers the best fit for this data set. Tables 2 and 3 demonstrated the results of ZIP, ZINP, and ZIGP models. According to the results of the ZIGP model (Table 3), some factors such as type of surgery, appendicitis (p= 0.008), abdomen and its contents

(p= 0.036), hemorrhoids (p= 0.011), lung (p= 0.059), skin (p= 0.04), type of insurance (p= 0.008), type of comorbid disease, hypertension (p= 0.023), heart disease (p=0.0470), hyperlipidemia (p= 0.047), type of residence (local or non-local) (p= 0.019), number of tests (p<0.001), and age (p= 0.045) had a significant effect on the LOS in the general surgical unit.

Table 3. Frequency Distribution and Result of the ZIGP Model

Parameter (category)	Number	%	Estimation	SE	p
Fixed value	334	-	0.15	1.139	0.89
Appendicitis (No*/Yes)	30	9	-3.70	1.411	0.008
Digestive disease (No*/Yes)	123	36.8	0.06	0.360	0.871
Abdomen and its contents (No*/Yes)	53	15.9	-0.74	0.353	0.036
Skeletal and muscle system (No*/Yes)	19	5.7	0.14	0.468	0.769
Blood and lymphatic system (No*/Yes)	13	3.9	-0.12	0.532	0.827
Glands (No*/Yes)	40	12	0.35	0.382	0.365
Urinary system (No*/Yes)	7	2.1	0.05	0.457	0.908
Hemorrhoids (No*/Yes)	21	6.3	-2.14	0.847	0.011
Cardiovascular system (No*/Yes)	25	7.5	-0.042	0.453	0.926
Female reproductive system (No*/Yes)	3	0.9	0.59	0.750	0.435
Breast (No*/Yes)	8	2.4	-0.61	0.826	0.458
Lung (No*/Yes)	5	1.5	-1.56	0.814	0.049
Skin (No*/Yes)	13	3.9	-2.23	1.088	0.040
Type of insurance (No*/Yes)	278	83.2	0.53	0.266	0.046
History of smoking (Non-smoker*/Smoker)	50	15	0.19	0.226	0.407
Hypertension (No*/Yes)	47	14.1	0.49	0.215	0.023
Diabetes (No*/Yes)	39	11.7	-0.004	0.239	0.987
Heart disease (No*/Yes)	38	11.4	0.44	0.260	0.047
Hyperlipidemia (No*/Yes)	16	4.8	0.58	0.305	0.047
Kidney disease (No*/Yes)	25	7.5	-0.03	0.309	0.916

Cnrd Table 3

Comorbid disease (No*/Yes)	41	12.3	-0.002	0.234	0.993
Hemoglobin (Not natural*/Natural)	243	72.8	-0.07	0.178	0.697
Gender (Female*/male)	220	65.9	0.055	0.192	0.774
Marital status (Single*/Married)	252	75.4	0.12	0.225	0.606
Place of residence (Non-local*/Local)	71	21.3	-0.43	0.183	0.019
Type of admission (Low risk*/Medium Risk)	56	16.8	0.001	0.275	0.997
Type of admission (Low risk*/High Risk)	13	3.9	-0.57	0.430	0.181
Type of admission (Emergency/ hospitalization)	219	65.6	-0.20	0.204	0.323
Number of tests	4.82	3.13	0.17	0.023	<0.001
Age	45.174	16.47	0.25	0.006	0.046
Number of surgery	1.59	1.02	0.06	0.090	0.496
Pulse	81.03	7.42	-0.005	0.012	0.701

*Reference group

For example, appendicitis is a significant variable on LOS and its estimated value is 3.71. Thus, we can interpret that the log of expected LOS is expected to be 3.71 days lower for patients who have appendicitis compared to those who don't, while holding the other variables constant in the model.

Discussion

Nowadays improving the quality and efficiency of health care services necessitates the application of efficient and quality indicators. Length of stay (LOS), one of these indicators, has been used as a source of hospital costs and quality indicator (2).

Length of stay, like other count data, needs count models to be analyzed. Poisson distribution has been widely used by researchers. Not having any overdispersion larger than expected, variability is the important assumption of the Poisson regression model. Until recent years, the NB (Negative Binomial) and GP (Generalized Poisson) models have been used to describe overdispersion caused by observed heterogeneity in distribution. The distribution of LOS involves a large proportion of zeros that can lead to overdispersion. Thus, zero-inflated models were fitted to the data. According to AIC criterion, ZIGP was the best model for analyzing this data. Effective demographic and hospital factors (clinical and non-clinical) were studied in the general surgery ward at Shariati hospital in Tehran using zero-inflated generalized Poisson regression model to analyze many zero-stay days. The results of zero-inflated generalized Poisson regression model revealed that type of surgery (appendicitis, abdomen and its contents, hemorrhoids, lung, and skin), type of insurance, type of comorbid disease (hypertension, heart disease, and hyperlipidemia), place of residence (local and non-local), and number of tests affected hospital stay. Non-local patients stayed longer than the local ones, as these patients referred to advanced referral hospitals in Tehran when their diseases reached severe stages and they needed more complex treatment. This delay lies in the lack of definitive diagnosis of multiple visits to doctors or shortcomings on essential and necessary facilities in their hometowns. Lack of accommodation in Tehran is another contributing factor for long hospital stay. These patients refer to the hospital for earlier hospitalization and diagnostic and therapeutic measures, which lead to prolonged hospital stay (22). Ravangard et al., who divided the place of residence in to Tehran, Tehran suburb, Western, Central, Northern,

Southern, and Southeast of Iran, concluded that patients coming from Southern and Southeastern regions of Iran have longer hospital stay than others (22). Age is found to be directly associated with length of stay, which is consistent with the results of the study conducted by Arab et al. (11). In the present study, type of insurance was found to be directly associated with length of stay. Patients without insurance had shorter length of hospital stay than those with insurance. This result is consistent with that of Arab et al. (11). The model was separately fitted for each of the 13 types of surgery. The results revealed that appendicitis, abdomen and its contents, and hemorrhoids were found to be significantly associated with length of stay. According to the negative coefficient of type of surgery (appendicitis, abdomen and its contents, hemorrhoids, and skin), such patients had shorter length of stay. The other effective factor was type of comorbid disease (hypertension, heart disease, and hyperlipidemia); the positive coefficients showed longer length of stay, as heart patients require long CCU, ICU, and surgical ward stay. Therefore, heart disease is found to be directly associated with length of hospital stay. Finally, the results of zero-inflated Poisson regression model indicated that increased number of tests led to longer hospital stay. This was due to many routine tests at the beginning for diagnosis as well as delay in response to inpatient wards, which is consistent with results of Ravangard et al. (22).

Conclusion

According to AIC, we conclude that zero-inflated generalized Poisson regression is more appropriate than zero-inflated Poisson regression and zero-inflated negative binomial models based on this dataset.

Acknowledgments

This study was financially supported by Tarbiat Mo-dares University. We also thank health professionals and staff in Shariati Hospital.

Conflict of Interests

The authors declare that they have no competing interests.

References

- Clarke A. Length of in-hospital stay and its relationship to quality of care. *Quality Safety Health Care*. 2002;11(3):209-10.
- Jimenez R, Lam R, Marot M, Delgado A. Observed-predicted length of stay for an acute psychiatric department, as an indicator of inpatient

- care inefficiencies. Retrospective case-series study. *BMC Health Service Res.* 2004;4(1):4.
3. Atienza N, García-Heras J, Muñoz-Pichardo J, Villa R. An application of mixture distributions in modelization of length of hospital stay. *Statistic Med.* 2008;27(9):1403-20.
 4. Ghoreishi Nejad S. Agent simulation in healthcare. Canada: The University of Regina Canada. 2008.
 5. Brooke BS, Goodney PP, Powell RJ, Fillinger MF, Travis LL, Goodman DC, et al. Early discharge does not increase readmission or mortality after high-risk vascular surgery. *J Vasc Surg* 2013;57:734-40.
 6. Lucas DJ, Haider A, Haut E, Dodson R, Wolfgang CL, Ahuja N, et al. Assessing readmission after general, vascular, and thoracic surgery using ACS-NSQIP. *Ann Surg.* 2013;258:430-9.
 7. Schneider EB, Hyder O, Wolfgang CL, Hirose K, Choti MA, Makary MA, et al. Patient readmission and mortality after surgery for hepatopancreato-biliary malignancies. *J Am Coll Surg.* 2012;215:607-15.
 8. Lim A, Tongkumchum P. Methods for analyzing hospital length of stay with application to inpatients dying in Southern Thailand. *Global Journal of Health Science.* 2009;1(1):p27.
 9. de Campos Lobato LF, Ferreira PCA, Wick EC, Kiran RP, Remzi FH, Kalady MF, et al. Risk factors for prolonged length of stay after colorectal surgery. *J Coloproctology.* 2013;33(1):22-7.
 10. Huang JQ, Hooper PM, Marrie TJ. Factors associated with length of stay in hospital for suspected community-acquired pneumonia. *Canadian respiratory journal: journal of the Canadian Thoracic Society.* 2006;13(6):317.
 11. Arab M, Zarei A, Rahimi A, Rezaiean F, Akbari F. Analysis of factors affecting length of stay in public hospitals in Lorestan Province, Iran. *Hakim Research Journal.* 2010;12(4):27-32.
 12. Hosseini AF, Gohari MR. Application of multilevel model in determining the effective factors in the length of stay among appendectomy patients. *Razi J Med Sci.* 2014;20(115):70-7.
 13. Marazzi A, Paccaud F, Ruffieux C, Beguin C. Fitting the distributions of length of stay by parametric models. *Medical care.* 1998;36(6):915-27.
 14. Famoye F, Wulu J, Singh KP. On the generalized Poisson regression model with an application to accident data. *J Data Sci.* 2004;2:287-95.
 15. Skrondal A, Rabe-Hesketh S. Some applications of generalized linear latent and mixed models in epidemiology: repeated measures, measurement error and multilevel modeling. *Norsk epidemiologi.* 2003;13(2).
 16. Wang K, Yau KK, Lee AH. A zero-inflated Poisson mixed model to analyze diagnosis related groups with majority of same-day hospital stays. *J Comp Method Prog Biomed.* 2002;68(3):195-203.
 17. Wang W, Famoye F. Modeling household fertility decisions with generalized Poisson regression. *J Population Econ.* 1997;10(3):273-83.
 18. Karlis D, Xekalaki E. Mixed poisson distributions. *Int Statist Rev.* 2005;73(1):35-58.
 19. Ng S, Yau K, Lee A. Modelling inpatient length of stay by a hierarchical mixture regression via the EM algorithm. *Math Comp Model.* 2003;37(3):365-75.
 20. Wulu J, Singh K, Famoye F, McGwin G. Regression analysis of count data. *J Indian Soc Agri Stat.* 2002;55(2):220-31.
 21. Hilbe J. *Negative binomial regression*: Cambridge University Press; 2011.
 22. Ravangard R, Arab M, Zeraati H, Rashidian A, Akbarisari A, Mostaan F. Patients' length of stay in women hospital and its associated clinical and non-clinical factors, Tehran, Iran. *Iranian Red Crescent Med J.* 2011;13(5):309.