



Study of *Porphyromonas gingivalis* in periodontal diseases: A systematic review and meta-analysis

Mohammad Rafiei¹, Faezeh Kiani², Fatemeh Sayehmiri³, Kourosh Sayehmiri⁴, Abdolkarim Sheikhi^{5*}, Mona Zamanian Azodi³

Received: 9 August 2016

Published: 12 Sep 2017

Abstract

Background: The mouth cavity hosts various types of anaerobic bacteria including *Porphyromonas gingivalis*, which causes periodontal inflammatory diseases. *P. gingivalis* is a gram-negative oral anaerobe and is considered as a main etiological factor in periodontal diseases. Several studies have reported a relationship between *P. gingivalis* in individuals with periodontal diseases and a critical role of this bacterium in the pathogenesis of periodontal diseases. The present study aimed at estimating this probability using a meta-analysis.

Methods: We searched several databases including PubMed, Scopus, Google Scholar, and Web of Science to identify case-control studies addressing the relationship between *P. gingivalis* with periodontal diseases. A total of 49 reports published from different countries from 1993 to 2014 were included in this study. I^2 (heterogeneity index) statistics were calculated to examine heterogeneity. Data were analyzed using STATA Version 11.

Results: After a detailed analysis of the selected articles, 49 case-control studies with 5924 individuals fulfilled the inclusion criteria for the meta-analysis. The healthy controls included 2600 healthy individuals with a Mean±SD age of 36.56±7.45 years. The periodontal diseases group included 3356 patients with a mean age of 43.62±8.35 years. There was a statistically significant difference between *P. gingivalis* in periodontal patients and healthy controls; 9.24 (95% CI: 5.78 to 14.77; $P = 0.000$). In the other word, there was a significant relationship between the presence of *P. gingivalis* and periodontal diseases.

Conclusion: Analyzing the results of the present study, we found a strong association between the presence of *P. gingivalis* and periodontal diseases. This result suggests that another research is needed to further assess this subject.

Keywords: *Porphyromonas gingivalis*, Periodontal Diseases, Chronic Periodontitis, Aggressive Periodontitis, Gingivitis

Copyright© Iran University of Medical Sciences

Cite this article as: Rafiei M, Kiani F, Sayehmiri F, Sayehmiri K, Sheikhi A, Zamanian Azodi M. Study of *Porphyromonas gingivalis* in periodontal diseases: A systematic review and meta-analysis. *Med J Islam Repub Iran*. 2017 (12 Sep);31:62. <https://doi.org/10.14196/mjiri.31.62>

Introduction

Periodontal diseases are complicated oral diseases, which are specified by bacterial- induced inflammatory destruction of tooth-supporting tissues (1). Periodontal diseases can be described as one of the predominant polymicrobial infections of humans, which can advance and lead to gum retrogression, soft tissue harm, weakness of the bone, eventual osteoporosis and tooth loss (severe periodontitis). Periodontal disease have many risk factors including smoking and diabetes, moreover, several bacteria have been connected to the intensity and progress of periodontitis (2, 3).

The oral cavity is a source of different microorganisms

that cause a class of infections and inflammation inside the cavity (4). So far, more than 700 bacterial taxa have been identified in sample s taken from oral cavities (2). Evidence for periodontal etiology relies on the performance of several criteria described by Socransky (5).

Of the bacteria believed to be pathogenic in periodontal disease, *P. gingivalis* has been extensively studied due to its unique ability to evade the immune response (6). *P. gingivalis* is a gram-negative oral anaerobe and considered as a main etiological factor in periodontal diseases by produc-

Corresponding author: Dr Abdolkarim Sheikhi, sheikhi.a@dums.ac.ir
sheikh@queensu.ca

¹ Department of Biostatistics and Epidemiology, School of Medicine, Arak University of Medical Sciences, Arak, Iran.

² Student Research Committee, Ilam University of Medical Sciences, Ilam, Iran.

³ Proteomics Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

⁴ Department of Social Medicine, School of Medicine, Ilam University of Medical Sciences, Ilam, Iran.

⁵ Department of Immunology and Microbiology, Dezfoul University of Medical Sciences, Dezfoul, Iran.

↑What is “already known” in this topic:

More than 700 bacteria are found in samples taken from the oral cavity. In several studies, qualitative information of the *p.gingivillis* has been done in oral diseases.

→What this article adds:

A variety of qualitative studies have been done on Gengivals, but there are just a few conducted quantitative studies on it which their results are poor and inconsistent as well. So a Meta Analysis method is used in this study to integrate the results of the all previous quantitative investigations into a unique and statistically valid conclusion.

ing a number of virulence factors and extracellular proteases such as lipopolysaccharide, fimbria, gingipain etc., resulting in destruction of periodontal tissues (7–11). The various surface components of *P. gingivalis* enable the bacterium to interact with the external medium and simplify its growth, nutrient gain, colonization, and formation of a biofilm that protects it against the host's defense (12, 13). In addition to being painful, persistent dental disease is linked to diabetes, heart disease, high blood pressure, and MS in later years of life; therefore, extensive studies have been conducted to control the bacteria causing dental diseases (14).

The pathogenicity of *P. gingivalis* has been investigated in a variety of experimental animal models such as rat, mouse, rabbit, drosophila, and cell models, showing complicated mechanisms of *P. gingivalis* -host interplay in the expansion of periodontal diseases (15-19).

Although many studies aimed to specify the macrobiotic dependents of specific disease types and the extent of periodontal destruction, there is yet no quantitative data on the levels of *P. gingivalis* in periodontal diseases. To authenticate the studies, performing a meta-analysis seemed necessary. Thus, this study aimed at evaluating the prevalence of *P. gingivalis* in patients with periodontal diseases.

Methods

Search Strategies

A database was built for the prevalence of *P. gingivalis* periodontal diseases from 1993 to 2014 using PubMed, Web of Science, Google scholar Medline, Embase, the Cochrane Library, and Scopus databases. The search was restricted to original articles published in English that presented the prevalence or incidence of *P. gingivalis* among patients with periodontal diseases. The following keywords from medical subject headings, titles, or abstracts were used with the help of Boolean operators (and, or): *P. gingivalis*, chronic periodontitis, aggressive periodontitis, and gingivitis. We also searched bibliographies of the retrieved articles for additional references. The titles from the search results were examined closely and determined to be suitable for potential inclusion into the study. In addition, the references from the selected articles were examined as a further search tool. Relevant trials noted in the reference lists of each selected article were also evaluated for inclusion. All papers whose keywords were present in their titles or abstracts were used in the initial list and other unrelated articles were eliminated.

Inclusion and Exclusion Criteria

All original articles presenting case-control studies on the prevalence of *P. gingivalis* in periodontal diseases were considered. The selection of articles for review was completed based on 3 stages: titles, abstracts, and full texts. When necessary, authors were contacted for additional information. Studies were excluded if they presented insufficient data, if they were not epidemiologic studies, and if they focused on the prevalence of *P. gingivalis* in diseases other than periodontal diseases. Review articles, congress abstracts, studies reported in languages other than English

or Persian, meta-analyses or systematic reviews, and duplicate publication of the same study were also excluded. The STROBE (strengthening the reporting of observational studies in epidemiology) statement was used for quality control of the studies. We assessed the quality of studies according to variables related to the study objectives, characteristics of the study population, clearly explained inclusion/exclusion criteria, data collection method, as well as the validity, explicit findings, and appropriate data analysis methods of the studies.

Data Extraction

For all studies, the following data were extracted: first author, year of publication, location, sample size, sample age, *P. gingivalis* screening method, and sample specimens, percentage of *P. gingivalis* in patients and healthy individuals. Abstracts and full articles were reviewed independently by 2 of the authors, and if results were discordant, papers were reviewed jointly until the differences were resolved.

Data Synthesis and Analysis

Studies were combined based on the sample size, mean and standard deviation. The difference between the average variance of the normal distribution was calculated using the formula of 2 integrated variance. To assess the heterogeneity of the studies, Cochran test and the I^2 (heterogeneity index) were used. Due to the significant heterogeneity in the studies, random effects model was used. The findings are described in forest plots (the point estimations and their 95% CI). To examine publication bias, Begg plot and regressions method were used. P value less than 5% was considered as a significant heterogeneity test. Sensitivity analyses were prespecified. Statistical analyses were performed using STATA version 11.

Results

Our initial search strategy yielded 172 potential articles for inclusion; in a secondary screening, 48 of them were excluded based on title and abstract evaluation, and 124 were retained for detailed full text evaluation. We excluded another 75 articles (9 review articles, 18 articles with other diseases, 13 for lack of enough information, 23 articles with percentage of *P. gingivalis* only in patients group, and 12 duplicates). After a detailed analysis of the selected articles, 49 case-control studies with 5924 individuals (3356 patients and 2600 healthy individuals) fulfilled the inclusion criteria for the meta-analysis (Fig. 1). The characteristics of the 49 trials and the quality scores included in the meta-analysis are summarized in Table 1.

The healthy controls comprised 2600 healthy individuals, with the age range of 14 to 67 years and a mean age of 36.56 ± 7.45 years. The periodontal diseases group included 3356 patients with periodontal diseases, ranging from 14 to 59 years, and with a mean age of 43.62 ± 8.35 years.

In the present study, a statistically significant difference was found between *P. gingivalis* in periodontal patients and healthy controls; 9.24 (95% CI: 5.78 to 14.77; $p < 0.001$) (Fig. 2). In the other word, on analyzing the results of the present study, a strong association was found between the

Table 1. Study characteristics

First author (Reference)	Country (year)	Case	Control	Mean age		OR (95% CI)		Type of disease	Sample specimens	Methods of Copper measurement
				Case	Control	Lower	Upper			
Michalowicz.B.S(52)	Jamaica(2000)	35	65	14-18	14-18	2.75	16.91	Periodontitis	subgingival plaque	PCR method
Kato.A (38)	Japan(2013)	85	20	57.4±13.1	45.9±17.0	0.14	1.50	Chronic periodontitis	plaque samples	PCR method
Lee.S.M (53)	Korea(2005)	17	19	52 ± 11.1	49 ± 10.2	0.19	3.92	Gingivitis	subgingival plaque	PCR method
R. Mikuls .T(54)		39	40	52 ± 11.1	49 ± 10.2	1.88	13.29	Periodontitis	subgingival plaque	stabilization P. gingivalis antibody seropositivity
L. Griffen.A (55)	USA(2009) Ohio State (1998)	130	181	51.4 ± 9.3	49.2 ± 9	6.52	19.21	Periodontitis	subgingival plaque	PCR method
Avila-Campos .Mj(40)	Brazil (2002)	50	50	45.5 ± 9.7	32.3 ± 8.9	0.68	3.56	Periodontitis	subgingival plaque	PCR method
Winkelhoff AJ(27)	Netherlands (2002)	116	94	42.9 ± 9.8	40.4 ± 11.9	5.81	26.19	Periodontitis	subgingival plaque	Anaerobic cultivation
Ruben.L (56)	Chile (2007)	20	6	27 ± 5.2	22.7 ± 4.9	0.16	6.20	Periodontitis/ gingivitis	subgingival plaque	Bacterial culture
Amano.A(33)	Japan (2013)	139	380			6.73	19.72	periodontitis	Dental plaque	PCR method
Takeuchi.Y(57)	Japan (2001)	103	20			100.27	21997	periodontitis	Saliva and subgingival plaque	PCR method
Missailidis. C.G(44)	Brazil (2004)	57	25			45.12	1211.10	periodontal attachment loss	subgingival plaque	PCR method
Missailidis. C.G (44)	Brazil (2004)	20	25			0.67	22.11	gingivitis	subgingival plaque	PCR method
Zhao.L (58)	Chile (2007)	115	136			8.48	29.49	Chronic periodontitis	subgingival samples	PCR method
Yang. H-W (59)	Taiwan (2004)	407	91			11.44	35.16	periodontal disease	subgingival plaque	indirect immunofluorescent assay
Choi.B-K (60)	Korea (2000)	29	20			11.5	1090.36	periodontitis	subgingival plaque	PCR method
Lyko.K (39)	Brazil (2013)	48	24	11.1 ± 3.52	10.54 ± 3.75	0.29	24.28	periodontitis	saliva samples	PCR method
Carinci.F (37)	Italy (2013)	66	46	48.9±18.2	31.6 ± 18.6	0.09	0.55	periodontitis	periodontal pocket	PCR method
Wilson.M (41)	USA (1993)	28	18	18-59	18-59	0.06	0.23	periodontitis	subgingival plaque	ELISA
M. Souccar.N (61)	Lebanon (2010)	20	20	34.3±5.36	26.10 ± 4.57	2.98	93.89	periodontitis	Oral plaque	PCR method
Riep.B (20)	Germany (2009)	46	21	55.2 ± 11.2	66.6 ± 1.5	0.64	5.95	Chronic periodontitis	subgingival plaque	PCR method
Riep.B (20)	Germany (2009)	44	21	34.4 ± 6.5	66.6 ± 1.5	0.40	3.50	aggressive periodontitis	subgingival plaque	PCR method
Yoneda.M (62)	Japan (2012)	150	60	54.6 ± 1.2	52.9 ± 2.4	1.58	6.33	periodontitis	saliva samples	PCR method
Feng. X-H (63)	China (2009)	48	25	38.9 ± 9.9	23.6 ± 1.8	7.56	134.46	periodontitis	subgingival plaque	PCR method
Zhou.T (32)	China (2013)	27	20			6.63	555.49	Chronic periodontitis	subgingival	PCR and reverse hybridization assay
Lu. RF (34)	China (2013)	80	56			60.69	1157.09	Aggressive periodontitis	gingival crevicular fluid	PCR method
Condorelli. F (64)	Italy (1998)	33	21			3.25	226.97	severe periodontal disease	subgingival plaque	culture
Wu.Y-M(65)	China (2007)	61	30	42.4±8.7	37.35 ± 7.3	4.05	54.58	chronic periodontitis	subgingival plaque	PCR method
Puig-Silla.M (12)	Spain (2012)	33	37	43.39 ± 7.4	40.68 ± 7.1	1.94	15.05	chronic periodontitis	subgingival plaque	PCR method
Puig-Silla.M (12)	Spain (2012)	16	37	38.81 ± 6.9	40.68 ± 7.1	0.47	5.63	Gingivitis	subgingival plaque	PCR method
Liu .Y (15)	China (2013)	25	20			3.58	118.36	periodontitis	gingival crevicular fluid	PCR method
Mayorga-Fayad.I (66)	Colombia (2007)	143	40	39.5±9.85	32.6±10.6	6.53	75.76	periodontitis	subgingival plaque	culture
Scapoli.L (35)	Italy (2012)	127	66	48.9 ± 18.2	31.6 ± 18.6	2.66	9.60	periodontitis	periodontal pocket microbiota	PCR method
Wang. P (30)	China(2014)	25	29			4.21	64.69	periodontitis	subgingival plaque	PCR method
Wara-aswapati. N (47)	Thailand (2009)	20	20			5.13	383.34	chronic periodontitis	subgingival plaque	PCR method
Faghri. J (67)	Iran (2007)	61	40	43 ± 11	41.35 ± 9.8	3.03	19.33	chronic periodontitis	subgingival plaque	PCR method
Feng. X-H (68)	China (2006)	55	17			5.06	87.12	Aggressive periodontitis	subgingival plaque	PCR method
Zhan .DF (69)	China (2005)	152	30			19.58	1289.11	chronic periodontitis	Periodontal pocket and gingival sulcus	PCR method
Amano.A (70)	Japan (2000)	139	380			6.73	19.72	periodontitis	Dental plaque	PCR method
Lau .L (45)	Spain (2004)	32	30	49.4 ± 8.99	37.8 ± 7.5	7.11	111.62	periodontitis	subgingival plaque	PCR method
Lau .L (45)	Spain (2004)	30	30	46.6 ± 9.82	37.8 ± 7.5	0.75	10.33	Gingivitis	subgingival plaque	PCR method
Yong. X (1)	China (2014)	42	32			0.50	3.29	chronic gingivitis	Gingival crevicular fluid	PCR method
Yong. X (1)	China (2014)	95	32			2.15	15.63	chronic periodontitis	Gingival crevicular fluid	PCR method
Tomita. S (31)	Japan (2013)	20	10	43.6 ± 11.1	28.7 ± 3.2	2.95	1187.72	chronic periodontitis	subgingival plaque	PCR method
Moon. J-H (7)	Korea (2013)	284	128	48.3 ± 9.5	42.3 ± 13.5	12.62	66.10	chronic periodontitis	subgingival plaque	PCR method
SuzanaStingu.C (36)	Germany (2012)	33	20	33.39 ± 10.47	37.65 ± 10.88	1.91	47.96	chronic periodontitis	subgingival plaque	PCR method
Enrique Botero. J (71)	Colombia (2007)	80	30	33.91 ± 9.32	26.90 ± 7.17	9.02	123.31	periodontitis	subgingival plaque	PCR method
R.van der Ploeg. J (72)	Switzerland(2004)	17	33	53.1 ± 8.53	26.8 ± 5.3	4.94	116.54	chronic periodontitis	subgingival plaque	PCR method
Takeuchi.Y (73)	Japan (2003)	35	18	51.8 ± 7.29	27.3 ± 3.71	34.03	9816.94	chronic periodontitis	subgingival plaque	PCR method
Suzuki.N (74)	Japan (2008)	21	305	43.8 ± 12.1	45.4 ± 14.9	1.97	15.46	chronic periodontitis	saliva samples	PCR method

presence of *P. gingivalis* and periodontal diseases (Fig. 2).

According to publication bias figure, the effect of bias in these studies was not significant. In fact, most studies were located inside the Funnel Plot, and thus the results of most relevant studies, considering the title, were included into the analysis ($p = 0.005$) (Fig. 3).

Discussion

It is widely accepted that the etiology of periodontal dis-

eases is polymicrobial in nature. Worsening or improvement of periodontal situation goes along by a change in the bacterial composition of subgingival plaque (20). It has, therefore, been suggested that microbial testing be used to diagnose and optimize periodontal therapy and assess its outcome. However, this strategy may be confusing as the beginning and the progress of periodontal diseases are impressed by interaction of myriad genetic, environmental, host, and microbial factors (20-23). Furthermore, molecular studies reveal an unexpectedly high diversity of microorganisms and progression of disease remains to be investigated. Nevertheless, current microbiological testing




Fig. 1. The flow diagram of studies identified the systematic review and meta-analysis

mostly involves the classic doubtful oral pathogens (20).

The aim of this study was to evaluate the association between the presence of *P. gingivalis* and periodontal diseases. *P. gingivalis* has been known to be a risk factor for periodontal diseases although its exact roles in the initiation and progression of the oral diseases remain unclear (15,

24).

In the present study, a significantly strong positive correlation was found between the presence of *P. gingivalis* and periodontal diseases. Published data suggests that the flora associated to chronic periodontal diseases is dominated by




Fig. 2. Meta-analysis of the association of *P. gingivalis* with periodontal diseases. Square represents effect estimate of individual studies with their 95% confidence intervals with size of squares proportional to the weight assigned to the study in the meta-analysis. In this chart, studies are stored in order of the year of publication and author's names, based on a random effects model.




Fig 3. Egger's funnel plot for publication bias. The diameter of each circle represents the weight in the meta-analysis. Each circle represents the RDs according to the standard error of each RDs. The diameter of each circle represents the weight in the meta-analysis.

the presence of Gram-negative anaerobic bacteria, and especially *P. gingivalis* (25). Slots and Ting described a synergistic association of *P. gingivalis* in presentations with evolutive lesions (26). Van Winkelhoff et al. concluded that *P. gingivalis* was the marker of a destructive lesion (27). Authors of previous studies reported similar results. The authors of various studies on *P. gingivalis* have reported that these pathogens are able to pollute soft tissues and flee the surgical debridement of periodontal lesions (28, 29).

Many epidemiological studies report a positive association between the presence of *P. gingivalis* and periodontal diseases (7, 25, 30-36), whereas in some studies, no association was found between the presence of *P. gingivalis* and periodontal diseases (20, 37-41). This has also been shown by Kumar et al. (42). The fact that all types of bacteria studied could be detected, even in healthy controls, might indicate that the presence of periodontal pathogens does not necessarily lead to periodontal diseases (1). However, these species show just a small percentage of whole bacteria, and the open-ended approach of their study may not be geared to indicate an association of this species with the disease (20). In another study, Kumar et al. demonstrated a significant association of *P. gingivalis* with chronic periodontitis using PCR amplification of 16S rRNA genes (43).

Some studies have reported a strong association between the presence of *P. gingivalis* and periodontitis, but did not find any association between the presence of *P. gingivalis* and gingivitis (1, 12, 44, 45). Although a number of putative bacteria are considered to be associated with chronic periodontitis, it has become clear that other factors are involved in the etiology of several types of periodontitis. In the other word, *P. gingivalis* is one of the risk factors responsible for periodontitis.

P. gingivalis is one of the bacteria that form the classic 'red complex' described by Socransky et al. (46). It is the bacteria most frequently found in patients with periodontal disease (12). Wara-aswapati et al. (47) found that the mean age of patients was higher than that of the healthy participants. *P. gingivalis* has been reported to be related to adult periodontitis (48, 49). In a study by Liu et al (15), they analyzed the correlation of patients' age and occurrence of *P. gingivalis* and found that the age of *P. gingivalis* positive and negative was statistically different, implying that the

prevalence of *P. gingivalis* may increase as the patients' age increase.

The association of certain putative pathogens with periodontal diseases shown in many studies may be explained in part by deeper pockets in these patients. In the present study, the association between probing severely and the occurrence of periodontal disease has not studied because most previous studies conducted on the association of particular bacteria with periodontitis did not test the influence of probing depth. Some studies found that the search depth had a much greater impact on the species than did the diagnosis. In study of Riep et al. (20), the prevalence of *P. gingivalis* was highly associated ($p < 0.001$) with pocket depth. Ali et al. (50) studied the existence of *P. gingivalis* in very deep pockets (≥ 6 mm) and deep pockets (4–5 mm) of 36 Romanian patients presenting with chronic periodontitis. *P. gingivalis* was recognized in 75.8% of the patients and 63.6% of the sampled sites.

Listgarten and Loomer (51) also questioned whether microbial identification should be considered as a strategy in managing this disease. They found no strong evidence supporting the benefit of microbial testing, partly due to a lack of standardization among diagnostic methods.

In summary, on analyzing the results of the present study, a strong association was found between the presence of *P. gingivalis* and periodontal diseases. Furthermore, the results of the present study indicated that *P. gingivalis* is one of the risk factors responsible for periodontitis. This result suggests that further research is needed to investigate their pathogenicity.

Acknowledgments

The authors extend their gratitude to the Student Research Committee, Ilam University of Medical Sciences for its support.

Conflict of Interests

The authors declare that they have no competing interests.

References

1. Yong X, Chen Y. Periodontopathogens and human b-defensin-2 expression in gingival crevicular fluid from patients with periodontal disease in Guangxi, China. *J Periodontal Res* 2015;50(3):403-410.
2. Tomita S, Komiya-Ito A. Prevalence of *Aggregatibacter actinomycetemcomitans*, *Porphyromonas gingivalis* and *Tannerella forsythia* in Japanese patients with generalized chronic and aggressive periodontitis. *Microb Pathogene*. 2013;61-62:11-15.
3. S.Michaud D. Role of bacterial infections in pancreatic cancer. *Carcinogenesis*. 2013;34(10):2193–2197.
4. Varghese J, Kumar Tumkur V, Ballal V, Subraya Bhat G. Antimicrobial effect of *Anacardium occidentale* leaf extract against pathogens causing periodontal disease. *Adv Biosci Biotechno*. 2013;4(8): 15–18.
5. Socransky SS. Relationship of bacteria to the etiology of periodontal disease. *J Dent Res*. 1970;49:203-222.
6. Tribble G, E Kerr J, Wang B-Y. Genetic diversity in the oral pathogen *Porphyromonas gingivalis*: molecular mechanisms and biological consequences. *Future Microbiol*. 2013;8:607–620.
7. Moon JH, Herr Y, Lee HW, Shin SI, Kim Ch, Amano A, Lee JY. Genotype analysis of *Porphyromonas gingivalis* fimA in Korean adults using new primers. *J Med Microb*. 2013; 62:1290-1294.
8. Decaillet F, Giannopoulou C, Cionca N, Almaghouth A, Mombelli A. Microbial profiles of patients seeking treatment for periodontitis.

- Influence of origin, smoking and age? Schweiz Monatsschr Zahnmed. 2012;122(3):198-204
9. Bostanci N, Belibasakis GN. Porphyromonas gingivalis: an invasive and evasive opportunistic oral pathogen. FEMS Microbiol Lett. 2012;331(1):1-9.
 10. Hayashi F, Okada M, Oda Y, Kojima T, Kozai K. Prevalence of Porphyromonas gingivalis fimA genotypes in Japanese children. J Oral Sci. 2012;54(1):77-83.
 11. Perez-Chaparro PJ, Lafaurie GI, Gracieux P, Meuric V, Tamanai-Shacoori Z, Castellanos JE, et al. Distribution of Porphyromonas gingivalis fimA genotypes in isolates from subgingival plaque and blood sample during bacteremia. Biomedica. 2009;29(2):298-306.
 12. Puig-Silla M, Dasi-Fernandez F, Montiel-Company JM, Almerich-Silla JM. Prevalence of fimA genotypes of Porphyromonas gingivalis and other periodontal bacteria in a Spanish population with chronic periodontitis. Med Oral Patol Oral Cir Bucal. 2012; 17(6):e1047-53.
 13. Yoshimura F, Murakami Y, Nishikwa K, Hasegawa Y, Kawaminami S. Surface components of Porphyromonas gingivalis. J Periodontal Res 2009; 44:1-12.
 14. Patra JK, Kim ES, Oh K, Kim HJ, Dhakal R, Kim Y, et al. Bactericidal effect of extracts and metabolites of Robinia pseudoacacia L. on Streptococcus mutans and Porphyromonas gingivalis causing dental plaque and periodontal inflammatory diseases. Molecules. 2015;20(4):6128-6139.
 15. Liu Y, Zhang Y, Wang L, Guo Y, Xiao S. Prevalence of porphyromonas gingivalis four rag locus genotypes in patients of orthodontic gingivitis and periodontitis. PLoS ONE. 2013;8(4): e61028.
 16. Igboin CO, Moeschberger ML, Griffen AL, Leys EJ. Porphyromonas gingivalis virulence in a Drosophila melanogaster model. Infect Immun. 2011;79(1):439-448.
 17. Fitzpatrick RE, Aprico A, Wijeyewickrema LC, Pagel CN, Wong DM, Potempa J, et al. High molecular weight gingipains from porphyromonas gingivalis induce cytokine responses from human macrophage-like cells via a nonproteolytic mechanism. J Innate Immun. 2009;1(2):109-117.
 18. Haruyama K, Yoshimura A, Naito M, Kishimoto M, Shoji M, Abiko Y, et al. Identification of a gingipain-sensitive surface ligand of porphyromonas gingivalis that induces Toll-like receptor 2- and 4- independent NF-kappaB activation in CHO cells. Infect Immun. 2009;77(10):4414-4420.
 19. Grenier D, Tanabe S. Porphyromonas gingivalis gingipains trigger a proinflammatory response in human monocyte-derived macrophages through the p38a mitogen-activated protein kinase signal transduction pathway. Toxins (Basel). 2010;2(3):341-352.
 20. Riep B, Edesi-Neuss L, Claessen F, Skarabis H, Ehmke B, Flemmig TF, et al. Are putative periodontal pathogens reliable diagnostic markers? J Clin Microbiol. 2009;47(6):1705-1711.
 21. Nibali L, Ready DR, Parkar M, Brett PM, Wilson M, Tonetti MS, et al. Gene polymorphisms and the prevalence of key periodontal pathogens. J Dent Res. 2007;86(5):416-420.
 22. Page RC, Kornman KS. The pathogenesis of human periodontitis: an introduction. Periodontol. 1997; 2000(14):9-11.
 23. Umeda M, Chen C, Bakker I, Contreras A, Morrison J.L, Slots J. Risk indicators for harboring periodontal pathogens. J Periodontol. 1998;69:1111-1118.
 24. Shi X, Hanley SA, Kele MF, Fawell SC, Aduse-Opoku J, Whiley RA, et al. The rag locus of Porphyromonas gingivalis contributes to virulence in a murine model of soft tissue destruction. Infect Immun. 2007;75(4):2071-2074.
 25. Benrachadi L, Bouziane A, Azziman Z, Bouziane-Ouartini F, Ennibi O. Screening for periodontopathogenic bacteria in severe chronic periodontitis in a Moroccan population. Médecine et maladies infectieuses. 2012;42(12):599-602.
 26. Slots J, Ting M. Actinobacillus actinomycetemcomitans and Porphyromonas gingivalis in human periodontal disease: occurrence and treatment. Periodontol. 2000;22:82-121.
 27. Van Winkelhoff AJ, Loos BG, van der Reijden WA, van der Velden U. Porphyromonas gingivalis, Bacteroides forsythus and other putative periodontal pathogens in subjects with and without periodontal destruction. J Clin Periodontol. 2002;29:1023-8.
 28. Meyer DH, Lippmann JE, Five-Taylor PM. Invasion of epithelial cells by Actinobacillus actinomycetemcomitans: a dynamic, multi-step process. Infect Immun. 1996;64:2988-97.
 29. Haffajee AD, Cugini MA, Dibart S, Smith C, Kentjr RL, Socransky SS. The effect of SRP on the clinical and microbiological parameters of periodontal diseases. J Clin Periodontol. 1997;24:324-34.
 30. Wang P, Duan D, Zhou X, Li X, Yang J, Deng M, et al. Relationship between expression of human gingival beta-defensins and levels of periodontopathogens in subgingival plaque. J Periodontal Res. 2014;50(1):113-122.
 31. Tomitaa S, Komiya-Itoa A, Imamura K, Kitaa D, Otaa K, Takayama S, et al. Prevalence of Aggregatibacter actinomycetemcomitans, Porphyromonas gingivalis and Tannerella forsythia in Japanese patients with generalized chronic and aggressive periodontitis. Microbial Pathogenesis. 2013;61-62:11-15.
 32. Ting Zh, Hong X, Zhaohui Y. Relationships of five periodontal pathogens causing subgingival plaque in patients with chronic periodontitis under different periodontal conditions. West China J Stomatol. 2013;5: 518-521.
 33. Amano A, Kuboniwa A, Nakagawa I, Akiyama S, Morisaki I, Hamada S. Prevalence of specific genotypes of Porphyromonas gingivalis fimA and periodontal health status. J Dent Res. 2000; 79(9):1664-8
 34. Lu R, Feng L, Gao X, Meng H, Feng X. Relationship between volatile fatty acids and Porphyromonas gingivalis and Treponema denticola in gingival crevicular fluids of patients with aggressive periodontitis. Beijing Da Xue Xue Bao. 2013;45(1):12-16.
 35. Scapoli L, Girardi A, Palmieri A, Testori T, Zuffetti F, Monguzzi R, et al. Microflora and periodontal disease. Dent Res J (Isfahan). 2012; 9(2):S202-S206.
 36. Stingu CS, Jentsch H, Eick S, Schaumann R, Knofler G, Rodloff A. Microbial profile of patients with periodontitis compared with healthy subjects. Quintessence Int. 2012;43(2):e23-31.
 37. Carinci F, Scapoli L, Girardi A, Cura F, Lauritano D, Nardi GM, et al. Oral microflora and periodontal disease: new technology for diagnosis in dentistry. Ann Stomatol. 2013;5(2):170-173.
 38. Kato A, Imai K, Ochiai K, Ogata Y. Higher Prevalence of Epstein-Barr Virus DNA in Deeper Periodontal Pockets of Chronic Periodontitis in Japanese Patients. PLoS ONE. 2013;8(8): e71990.
 39. Kato A, Imai K, Ochiai K, Ogata Y. Salivary detection of periodontopathic bacteria in Fanconi's anemia patients. Anaerobe. 2013;24: 32-35.
 40. Avila-Campos MJ, Velasquez-Melendez G. Prevalence of putative periodontopathogens from periodontal patients and healthy subjects in São Paulo SP Brazil. Rev Inst Med trop S. Paulo. 2002;44(1):1-5.
 41. Wilson M, Lopatin D, Osborne G, Kieser JB. Prevalence of Treponema denticola and Porphyromonas gingivalis in plaque from periodontally- healthy and periodontally-diseased sites. J Med Microbiol. 1993;38:406-410.
 42. Kumar PS, Leys EJ, Bryk JM, Martinez FJ, Moeschberger ML, Griffen AL. Changes in periodontal health status are associated with bacterial community shifts as assessed by quantitative 16S cloning and sequencing. J Clin Microbiol. 2006;44:3665-3673.
 43. Kumar PS, Griffen AL, Barton JA, Paster BJ, Moeschberger ML, Leys EJ. New bacterial species associated with chronic periodontitis. J Dent Res. 2003;82:338-344.
 44. Missailidis CG, Umeda JE, Ota-Tsuzuki C, Anzai D, Mayer MP. Distribution of fimA genotypes of Porphyromonas gingivalis in subjects with various periodontal conditions. Oral Microb Immunol. 2004;19(4):224-229.
 45. Lau L, Sanz M, Herrera D, Morillo JM, Martín C, Silva A. Quantitative real-time polymerase chain reaction versus culture: a comparison between two methods for the detection and quantification of Actinobacillus actinomycetemcomitans, Porphyromonas gingivalis and Tannerella forsythensis in subgingival plaque samples. J Clin Periodontol. 2004;31:1061-1069.
 46. Socransky SS, Haffajee AD, Cugini MA, Smith C, Kent RL, Jr. Microbial complexes in subgingival plaque. J Clin Periodontol. 1998;25(2):134-44.
 47. Wara-aswapati N, Pitiphat W, Chanchaimongkon L, Taweechaisupapong S, Boch JA, Ishikawa I. Red bacterial complex is associated with the severity of chronic periodontitis in a Thai population. Oral Dis. 2009;15:354-9.
 48. Décaillon F, Giannopoulou C, Cionca N, Almaghouth AA, Mombelli A. Microbial profiles of patients seeking treatment for periodontitis. Influence of origin, smoking and age? Schweiz Monatsschr Zahnmed. 2012;122(3):198-204.
 49. Hayashi F, Okada M, Oda Y, Kojim T, Kozai K. Prevalence of Porphyromonas gingivalis fimA genotypes in Japanese children. J Oral Sci. 2012; 54(1):77-83.

50. Ali RW, Velcescu C, Jivanescu MC, Lofthus B. Prevalence of putative periodontal pathogens in subgingival plaque samples from adult periodontitis patients. *J Clin Periodontol.* 1996;23:133.
51. Listgarten MA, Loomer PM. Microbial identification in the management of periodontal diseases. A systematic review. *Ann Periodontol.* 2003;8:182-192.
52. Michalowicz BS, Ronderos M, Camara-Silva R, Contreras A, Slots J. Human herpesviruses and *Porphyromonas gingivalis* are associated with Juvenile periodontitis. *J Periodontol.* 2000;71:981-988.
53. Lee SM, Yoo SY, Kim HS, Kim KW, Yoon YJ, Lim SH, et al. Prevalence of Putative Periodontopathogens in Subgingival Dental Plaques from Gingivitis Lesions in Korean Orthodontic Patients. *The J Microb.* 2005;260-265.
54. Mikuls TR, Payne JB, Reinhardt RA, Thiele GM, Maziarz E, Cannella AC, et al. Antibody responses to *Porphyromonas gingivalis* (*P. gingivalis*) in subjects with rheumatoid arthritis and periodontitis. *Int Immunopharmacol.* 2009;9(1):38-42.
55. Griffen AL, Becker MR, Lyons SR, Moeschberger ML, Leys EJ. Prevalence of *Porphyromonas gingivalis* and Periodontal Health Status. *J Clin Microbiol* 1998; 36(11):3239-3242.
56. Leon R, Silva N, Ovalle A, Chaparro A, Ahumada A, Gajardo M, et al. Detection of *Porphyromonas gingivalis* in the Amniotic Fluid in Pregnant Women With a Diagnosis of Threatened Premature Labor. *J Periodontol.* 2007;78:1249-1255.
57. Takeuchi Y, Umeda M, Sakamoto M, Benno Y, Huang Y, Ishikawa I. *Treponema socranskii*, *Treponema denticola*, and *Porphyromonas gingivalis* Are Associated With Severity of Periodontal Tissue Destruction. *J Periodontol.* 2001;72(10):1354-1363.
58. Zhao L, Wu YF, Meng S, Yang H, OuYang YL, Zhou XD. Prevalence of *fimA* genotypes of *Porphyromonas gingivalis* and periodontal health status in Chinese adults. *J Periodont Res.* 2007;42(6):511-517.
59. Yang HW, Huang YF, Chou MY. Occurrence of *Porphyromonas gingivalis* and *Tannerella forsythensis* in Periodontally Diseased and Healthy Subjects. *J Periodontol.* 2004;75(8):1077-1083.
60. Choi BK, Park SH, Yoo YJ, Choi SH, Chai JK, Cho KS, et al. Detection of Major Putative Periodontopathogens in Korean Advanced Adult Periodontitis Patients Using a Nucleic Acid-Based Approach. *J Periodontol.* 2000;71(9):1387-1394.
61. Souccar NM, Chakhtoura M, Ghafari JG, Abdelnoor AM. *Porphyromonas gingivalis* in dental plaque and serum C-reactive protein levels in pregnancy. *J Infect Dev Ctries.* 2010;4(6):362-366.
62. Yoneda M, Naka S, Nakano K, Wada K, Endo H, Mawatari H, et al. Involvement of a periodontal pathogen, *Porphyromonas gingivalis* on the pathogenesis of non-alcoholic fatty liver disease. *BMC Gastroenterology.* 2012;12:16.
63. Feng XH, Zhang L, Meng HX, Xu L, Chen ZB, Shi D, et al. Detection of 3 anaerobic microorganisms in saliva and subgingival plaque of patients with periodontitis. 2009. *J Peking Uni.* 2009;41(1):44-48.
64. Condorelli F, Scalia G, Cali G, Rossetti B, Nicoletti G, Lo Bue AM. Isolation of *Porphyromonas gingivalis* and Detection of Immunoglobulin A Specific to Fimbrial Antigen in Gingival Crevicular Fluid. *J Clin Microbiol.* 1998;36(8):2322-2325.
65. Wu YM, Yan J, Chen LL, Gu ZY. Association between infection of different strains of *Porphyromonas gingivalis* and *Actinobacillus actinomycetemcomitans* in subgingival plaque and clinical parameters in chronic periodontitis. *J Zhejiang Uni Sci B.* 2007; 8(2):121-131.
66. Mayorga-Fayad I, Lafaurie GI, Contreras A, Castillo DM, Baron A, Aya Mdel R. Subgingival microbiota in chronic and aggressive periodontitis in Bogotá, Colombia: an epidemiological approach. *Biomédica.* 2007;27:21-33.
67. Faghri J, Moghim S, Abed AM, Rezaei F, Chalabi M. Prevalence of *Porphyromonas gingivalis* and *Bacteroides forsythus* in chronic periodontitis by multiplex PCR. *Pak J Biol Sci.* 2007;15;10(22):4123-7.
68. Feng XH, Zhang L, Meng HX, Xu L, Chen ZB, Shi D. Prevalence of putative periodontal microorganisms in Chinese patients with aggressive periodontitis. *Zhonghua Kou Qiang Yi Xue Za Zhi.* 2006; 41(6):344-7.
69. Zhan D, Liu Z, Xia X, Hu J, Chen L, Yan J. Study on the detection of *P. gingivalis*, *A. actinomycetemcomitans* and *T. denticola* and the correlation between coinfections of the microbes and levels of chronic periodontitis lesion. *Zhonghua Liu Xing Bing Xue Za Zhi.* 2005;26(2):120-3.
70. Amano A, Kuboniwa A, Nakagawa I, Akiyama S, Morisaki I, Hamada S. Prevalence of specific genotypes of *Porphyromonas gingivalis fimA* and periodontal health status. *J Dent Res.* 2000; 79(9):1664-8.
71. Botero JE, Contreras A, Lafaurie G, Jaramillo A, Betancourt M, Arce RM. Occurrence of Periodontopathic and Superinfecting Bacteria in Chronic and Aggressive Periodontitis Subjects in a Colombian Population *J Periodontol.* 2007;78:696-704.
72. van der Ploeg JR, Giertsen E, Lüdin B, Mörgeli Ch, Zinkernagel AS, Gmür R. Quantitative detection of *Porphyromonas gingivalis* ϵ mA genotypes in dental plaque. *FEMS Microbiology Letters.* 2004; 232: 31-37.
73. Takeuchi Y, Umeda M, Ishizuka M, Huang Y, Ishikawa I. Prevalence of periodontopathic bacteria in Aggressive Periodontitis patients in a Japanese population. *J Periodontol.* 2003;74(10):1460- 1469.
74. Suzuki N, Yoneda M, Naito T, Iwamoto T, Masuo Y, Yamada K, et al. Detection of *Helicobacter pylori* DNA in the saliva of patients complaining of halitosis. *J Med Microb.* 2008; 57:1553-1559.