

Brief Communication

SUPINE LENGTH, WEIGHT AND HEAD CIRCUMFERENCE OF NEONATES AT BIRTH IN URBAN AREAS OF ARAK AND FACTORS AFFECTING THEM

Keywords: Neonates, supine length, head circumference, weight at birth, NCHS
MJIRI, Vol. 19, No. 4, 363-366, 2006.

Supine length, weight and head circumferences of 10241 neonates (5241 boys, 5000 girls, sex ratio 105) born in Arak (central Iran) in 2002 are reported. Mean+SD of boys and girls (p -value for sex difference) supine length (mm), weight (gr) and head circumference (mm) were estimated as 501+30 and 497+31 ($p<0.001$); 3194+586 and 3100+532 ($p<0.001$); 351+18 and 345+18 ($p<0.001$), respectively. Supine length and weight of our subjects were significantly lower than that of their American counterparts ($p<0.01$), but our boys head circumferences were significantly higher than the NCHS reference data ($p<0.01$), while no significant differences was seen among girls.

Anthropometric measurements at birth are significant indicators for predicting neonatal health. Several studies have shown that different anthropometric measurements at birth can be used as valid indicators of low birth weight.¹⁻³

Two limited studies were carried out on birth weight in Iran over 30 years ago^{4,5} and several others recently.⁶⁻

¹⁰ The only population based study on healthy neonates was carried out in 1996,¹¹ which include supine length, weight, arm, chest and head circumferences at birth and provide reference data for sizes at birth in southern Iran.

At present, no data are available on sizes at birth in urban Arak areas (Iran). The present study reports supine length, weight and head circumferences of all neonates born in 2002 in Arak, the main city in the central province of Iran. The paper further compares the data with that of NCHS¹² and south Iranian reference data.¹¹

The data relate to a cross-sectional study of all the 10241 live neonates (5241 boys, 5000 girls, sex ratio 1.05) born in 2002 in Arak. Arak is an industrial city with a population of nearly 700,000 of whom a large proportion immigrated from western provinces of Iran. The city is located 270 km south west of Tehran, Iran's capital and is classified as a semi-developed city.

Supine length (SL) and head circumferences (HC) were measured to the nearest 1 mm with non-stretchable plastic coated tapes. Weight (WT) was measured by

Table I. Anthropometric measurements at birth of Arak neonates.

Variable	Boys			Girls			P
	Mean	SD	95% CI	Mean	SD	95% CI	
Weight (gr)	3194	586	3178-3210	3100	532	3085-3115	<0.001
Supine Length (mm)	501	30	500-502	497	30	496-497	<0.001
Head Circumference (mm)	351	18	350-351	345	18	345-347	<0.001

using baby weighing scales to the nearest 10g. Centile statistics and nonparametric tests were used to compare our data with local and universal reference data. Student's t-tests were used to compare differences between our data and other reference data. Pearson corre-

Correspondence: S.M.T. Ayatollahi, Ph.D, FSS, CStat., Department of Biostatistics and Epidemiology, School of Public Health, Shiraz University of Medical Sciences, Shiraz. E-mail: ayatolahim@sums.ac.ir, Telephone:+98-711-7251001, Fax: +98-711-7260225

Length, Weight and Head Circumference in Arak Infants

Table II. Centiles of anthropometric measurements at birth of Arak neonates.

Centiles Variable		5	10	25	50	75	85	90	95
		Boys	Weight (gr)	2250	2500	2900	3250	3500	3700
Supine Length (mm)	460		470	490	500	520	530	530	540
Head Circumference (mm)	320		330	340	350	360	365	370	370
Girls	Weight (gr)	2200	2500	2827	3150	3450	3550	3700	3828
	Supine Length (mm)	450	470	490	500	510	520	530	540
	Head Circumference (mm)	320	330	340	350	355	360	360	370

Table III. Anthropometric measurements at birth of Arak neonates & Shiraz study and NCHS.

Study	Variable	Boys				Girls				P
		Mean	SD	95% CI	P*	Mean	SD	95% CI	P**	
Arak	Weight (gr)	3194	586	3178-3210	—	3100	532	3085-3115	—	<0.001
	Supine Length (mm)	501	30	500-502	—	497	30	496-497	—	<0.001
	Head Circumference (mm)	351	18	350-351	—	345	18	345-347	—	<0.001
Shiraz	Weight (gr)	3300	500	2300-4300	<0.001	3150	450	2250-4000	<0.001	<0.005
	Supine Length (mm)	494	23	448-540	<0.001	448	21	446-530	<0.001	<0.05
	Head Circumference (mm)	346	15	316-376	<0.001	342	13	316-368	<0.001	<0.0001
NCHS	Weight (gr)	3309	430	—	<0.001	3239	410	—	<0.001	—
	Supine Length (mm)	500	20	—	0.004	496	18	—	0.202	—
	Head Circumference (mm)	345	18	—	<0.001	341	19	—	<0.001	—

P: p-value for comparing mean of the variables in boys and girls

P*: p-value for comparing of mean of boys in Arak & Shiraz and NCHS

P**: p-value for comparing of mean of girls in Arak & Shiraz and NCHS

NCHS values References: New pediatric growth charts provide a tool toward future weight problems. URL: <http://www.cdc.gov/nchs/releases/00news/grow.hrt/htm>(last reviewed in March 01,2001).

Table IV. Correlation coefficient between anthropometric measurements at birth of Arak neonates.

	Weight	Supine Length	Head Circumference
Weight	_____	0.509*	0.552*
Supine Length	0.518*	_____	0.471*
Head Circumference	0.536*	0.534*	_____

* $p < 0.01$.

lation coefficients were applied to find the correlations of the anthropometric measurements.

Table I presents mean (SD) measures at birth of boys and girls and their 95 percent confidence intervals (95% CI). The significance of anthropometric measurement differences between boys and girls is also reported. Boys' measures were significantly greater than girls ($p < 0.001$). Centiles of anthropometric measurements at birth of boys and girls are given in Table II. Table III compares our data with that of NCHS¹² and reference data for Shiraz.¹¹ The correlation between neonate anthropometric measurements at birth is given in Table IV. Anthropometric measurements were correlated to each other significantly ($p < 0.001$). However, correlation of head circumference with weight was the highest.

Anthropometric measurements at birth of our subjects were significantly lower than that of NCHS for both sexes ($p < 0.001$). Also, anthropometric measurements at birth of our subjects were significantly lower than that of Shiraz for both sexes ($p < 0.001$), the reference data at birth for Shiraz, Iran, the most developed city of southern Iran.¹¹ This trend was observed when we compared our weight data with that of Tabriz,⁸ a main developed city located in north west Iran. However, birth weight of Islamshahr (a north Tehran developing city) neonates¹⁰ were significantly lower than ours ($p < 0.001$), showing that birth weight may be considered as a good indicator of development. By using multivariate analysis the joint vector of height, weight and head circumference in boys was greater than this vector in girls ($p < 0.001$).

Birth weight of our neonates were significantly lower ($p < 0.001$) than their counterparts in Czechs and Sweden¹³ as well as USA,¹² but higher than Nigerian neonates¹⁴ for both sexes. Boys' birth weight had been significantly higher than that of girls' in our study ($p < 0.001$), which concurs with other European¹⁵⁻¹⁷ and American studies,¹² apart from Iranian studies cited earlier.^{8,10,11}

Birth weight dominates other anthropometric measurements in most studies. We have studied supine length and head circumference at birth of neonates and shown that they were highly significantly correlated with birth weight ($p < 0.001$). This justifies that in the absence

of a weighing scale and due to technical difficulties in measuring weight in developing societies, one can use head circumference at birth as an alternative valid indicator of low birth weight, as we have achieved here, which concurs with similar results for arm and chest circumferences in other studies.^{1,3}

Birth weights and head circumference at birth of neonates whose mothers had normal delivery (57%) were significantly higher ($p < 0.001$) than those whose mothers endured caesarian section (43%). However, the difference between their supine lengths was not significant in the two mentioned groups.

S.M.T. AYATOLLAHI, PhD, FSS, CStat., AND M. RAFIEE, M.S.

From the Department of Biostatistics and Epidemiology, School of Public Health, Shiraz University of Medical Sciences, Shiraz, and the Department of Biostatistics and Health, Arak University of Medical Sciences, Arak, Iran.

REFERENCES

1. Arisory AE, Sarman G: Chest and mid arm circumferences: identification of low birth weight newborns in Turkey. *J Trop Paediatr* 41: 34-37, 1995.
2. Krishnamurthy S, Joshi S: Gender differences and low birth weight with maternal smokers tobacco use in pregnancy. *J Trop Paediatr* 39: 253-54, 1993.
3. Fawcus SR, Bassett MT: Chest circumferences as an indicator of low birthweight in Zimbabwe. *J Trop Paediatr* 39: 48-9, 1993.
4. Hedayat SH, Koohestani PA, Ghassemi H: Birth weight in relation to economic status and certain maternal factors based on an Iranian sample. *Trop Geograph Med* 23: 355-64, 1971.
5. Sarram M, Saadatnejadi M: Birth weight in Shiraz (Iran) in relation to maternal socioeconomic status. *Obstet Gynecol* 30: 360-67, 1967.
6. Soheiliazad AA, Jazayeri A, Jalali M, Sadeghi R: Infants nutrition and growth from birth to two years. *J Hakim* 1(2): 117-126 [Persian], 1998.

Length, Weight and Head Circumference in Arak Infants

7. Qadamli P, Mohseni SH: Maternal risk factors and low birth weight. *Tabriz Uni Med J* 48: 41-44 [Persian], 2000.
8. Rafrat M, Mahdavi R, Milani AS, Mahboob SA: Maternal overweight in pregnancy and weight at birth. *Tabriz Uni Med J* 55: 31-37 [Persian], 2002.
9. Nikavar A, Moqimi A, Chehreie A: Sizes at birth in 1000 healthy neonates in a Tehran hospital. *Iran Uni Med J* 21: 603-607 [Persian], 2003.
10. Kaseb F, Kimiagar S, Ghaffarpour M, Walaie N: Effect of traditional food supplementation during pregnancy on maternal weight gain and birthweight. *Int J Vitam Nutr Res* 72(6): 389-93, 2002.
11. Ayatollahi SMT, Shahsawary S: Sizes at birth in Shiraz, Iran. *J Trop Paediatr* 48: 245-247, 2002.
12. NCHS, Advance data, CDC growth charts: United States. US Department of Health and Human Services, Number 314, 30 May, 2000.
13. Koupilova I, Vagero D, Leon DA: Social variation in size at birth and preterm delivery in the Czech republic and Sweden, 1989-1991. *Paediatr Perinatal Epidemiol* 12: 7-24, 1998.
14. Ebomoy E, Adetoro OO, Wickremasinghe AR: Birth weight and sociological factors in Ilorin, Nigeria. *J Biosoc Sci* 23:417-23, 1991.
15. Johnson AA, Knight E, Edwards C, Oyemade UJ, Cole OJ, Westeny OE: Dietary intakes, anthropometric measurements and pregnancy outcomes. *J Nutr* 124: 5936-5942, 1994.
16. Antal M, Regoly MA, Varsanyi H, Birol L, Sagi K, Molnar DV: Nutritional survey of pregnant women in Hungary. *International J Vit Nutr Res* 67:115-122, 1997.
17. Guinhard Costa AM, Grange G, Larroch JC, Papierni K: Sexual differences in anthropometric measurements in France. *Biol Neonate* 72: 156-64, 1997.