


Study Protocol: The Feeding Exercise Randomized Trial among Overweight and Obese Adolescents in Greece

Anna Patsopoulou¹, Georgios Rachiotis^{1*} , Antonios Katsioulis¹, Paschalis Adam Molyvdas², Christos Hadjichristodoulou¹

Received: 29 Jul 2019

Published: 21 Mar 2022

Abstract

Background: This paper describes the study protocol used in the Feeding Exercise Clinical Trial in Adolescents in the region of Larissa in Greece, a randomized controlled clinical trial, among overweight/obese adolescents.

Methods: The main aim of the study was to comparatively evaluate the effectiveness of 2 different clinical interventions among 12 to 18-year-old overweight and obese adolescents. The first group participated in an exercise program and the second group in a combined dietary and exercise program. The third group was the control group. The study was conducted between 2014 and 2015. All adolescents aged 12 to 18 years old from public schools of Larissa and also their parents asked to participate. The effects of the intervention program will be analyzed by repeated-measures analysis of variance or the Friedman test. Changes in lifestyle behaviors from the baseline to the end of the intervention will be assessed using a chi-square test for categorical variables. A Pearson or a Spearman correlation coefficient and a linear regression analysis will be performed to explore any associations between quantitative variables. The following parameters were measured among adolescents: height, weight, body mass index, waist circumference, systolic and diastolic pressure, pulse rate, dietary and exercise habits of the adolescents and their parents.

Conclusion: This is the first clinical trial in Greece investigating the impact of clinical interventions on obesity among adolescents. It is expected that the results will provide useful insights into the effectiveness of clinical interventions among overweight and obese adolescents in Greece.

Keywords: Obesity, Overweight, Adolescents, Clinical Trial, Body Mass Index, Greece

Conflicts of Interest: None declared

Funding: The study has been supported by the Post Graduate Program: "Applied Public and Environmental Hygiene" at the Medical Faculty, School of Science, University of Thessaly, Greece.

*This work has been published under CC BY-NC-SA 1.0 license.

Copyright© Iran University of Medical Sciences

Cite this article as: Patsopoulou A, Rachiotis G, Katsioulis A, Molyvdas PA, Hadjichristodoulou C. Study Protocol: The Feeding Exercise Randomized Trial among Overweight and Obese Adolescents in Greece. *Med J Islam Repub Iran.* 2022 (21 Mar);36:24. <https://doi.org/10.47176/mjiri.36.24>

Introduction

Obesity is a common but often underestimated condition of clinical and public health importance all over the world. In Europe and the United States of America, obesity and excess weight constitute an epidemic that is a serious public health hazard and a major risk factor for morbidity and mortality (1-3). Health educators are currently being asked to become involved in child overweight prevention programs (4). Obesity is a significant health

problem for adolescents and could be predictive of abnormal body mass index (BMI) among children and adults (5-8). Adolescents represent one of the most vulnerable population group with regard to nutrition and health education (9, 10). Adolescence refers to as a period of major physical and cognitive change and it has been considered as a "critical period" for the development of adult obesity; and adolescents with established overweight or obesity

Corresponding author: Dr Georgios Rachiotis, g.rachiotis@gmail.com

¹ Department of Hygiene and Epidemiology, Medical Faculty, University of Thessaly, Thessaly, Greece

² Department of Physiology, Medical Faculty, University of Thessaly, Larissa, Greece

↑What is "already known" in this topic:

Obesity among adolescents is an underestimated condition of clinical and public health importance.

→What this article adds:

This is the first clinical trial in Greece investigating the impact of clinical interventions on obesity among adolescents.

have a moderate to substantial risk of remaining so in adulthood (11). Adult diseases related to obesity are now becoming more prevalent among the youth (12, 13). The growing economic impact of obesity is also a concern (14-16). Key health and medical organizations (16, 17) in several countries have recommended that research should be conducted on the effectiveness of well-designed interventions to combat overweight and obesity in children and adolescents (18, 19).

In addition, there is evidence that over 60% of young people eat more than the recommended amount of fat, while only 34% of boys and 33% of girls consume the daily suggested servings of fruits and vegetables (17). Nearly 25% of the calories that adolescents consume are from low nutritional value foods (20). Moreover, it is estimated that the vast majority of girls and boys consume more than 10% of their calories from saturated fat (21, 22). Despite their natural tendencies, adolescents have become less physically active in comparison to those who were children 50 years ago (23). The Center for Disease Control reported that less than 50% of adolescents are physically active on a regular basis, with a sharp decline in activity from childhood. It has been suggested that 20% of adolescents are not engaged in any leisure time physical activity (24, 25). In order to improve the health of the adolescent population, effective intervention programs focused on modifying unhealthy lifestyle behaviours are needed (18, 19, 25). Regular physical activity in teenagers and adolescents has been shown to improve strength and aerobic endurance, bone mass and density, blood glucose regulation, and reduce body fat (26, 27). In Greece, various researches have been conducted on the frequency of overweight and obesity among children. On the contrary, only few studies investigated overweight and/or obesity among adolescents (28-31). In addition, to the best of our knowledge, no clinical trial has been conducted in Greece investigating the impact of clinical interventions on obesity among adolescents.

Consequently, the Feeding Exercise Clinical Trial in Adolescents (FETA) program aims to compare and evaluate the efficacy of two different clinical interventions among overweight and obese adolescents 12-18-year-old.

Methods

Study Design

The study consisted of 2 parts.

Part 1. Descriptive Study: The first part included the measurement of the following parameters among 12 to 18 year-old students in public high schools of Larissa: height, weight, BMI, waist circumference, systolic and diastolic blood pressure, and cardiac rate. In addition, information was obtained on dietary and exercise habits of the adolescents and their parents. All the overweight or obese students detected in the first phase of the study were invited to take part in the randomized control trial.

Part 2. Intervention Program (Clinical Trial): In the clinical trial, 2 different intervention groups and 1 control group were employed. The first group participated in an exercise program and the second group in a combined dietary and exercise program. The third group was the

control group. Each group comprised of 50 individuals, aged 12 to 18, of both sexes. All groups completed a 50-m speed trial and filled out a questionnaire concerning their nutritional and exercise habits, both at the start and at the completion of the program. The students participated in a 50-m speed trial. The same test took place both at the start and at the finish of the program. The implementation of the program will start with the speed trials for all the 3 groups. The first group participated in an exercise program (duration: a 3-month program, training 3 times a week for 45 minutes at a time). The second group underwent a combined training and nutrition program for 3 months, meeting 3 times a week, for 45 minutes at a time. The first 10 minutes of each session included discussion regarding dietary issues, aiming at supplying knowledge and changing habits. The remaining time was used for the exercise program. The third group, the control group, only completed the questionnaires and participated in the speed trial tests at the beginning and the completion of the program. The intervention program lasted for 3 months, with 45 minutes per week. Measurements (weight, height, BMI, and waist circumference) continued for 3 months after the completion of the program (total duration of the program: 6 months).

Inclusion criteria

All students from the 15 public middle schools in the city of Larissa in central Greece were invited to measure their weight and height to calculate their BMI. All the overweight or obese students were invited to take part in the study.

Exclusion Criteria

Those who had an organic cause for their obesity and those who received any medication that could interfere with growth or weight control were excluded from the study.

Participants

To detect a 0.28 standardized effect size from the baseline to 6 months in the BMI score, with a power of 80% at 2-sided 5% level of significance and assuming a dropout rate of 20%, a minimum sample size of 65 participants per group was required. Analyses were conducted on a modified intention-to-treat basis, which included all randomized participants who had a baseline and at least 1 post-baseline completed questionnaire, with the last observation carried forward.

Outcome Measures

The following parameters were measured pre- and postintervention.

(i) Adiposity: Weight will be measured by the use of children barefoot and wearing light clothing, with Tanita HD646 scales (Tanita Corporation of America Inc) to 0.1 kg. In addition, the parameter of height will be measured using the stretch stature method and PE87 portable stadiometers (Mentone Educational Centre). Nonextensible steel tapes will be used to assess waist circumference, which is measured at the mid-point level between the low-

er costal border and the iliac crest. Anthropometric measures are conducted using the International Society for the Advancement of Kinanthropometry procedures (32). With regard to the BMI, it will be calculated via measuring weight and height. Moreover, we will use the Cole scale to establish a definition for children's overweight and obesity (33). Last, the waist circumference will be measured by the use of a tape measure.

(ii) Metabolic Profile Measures: Systolic and diastolic blood pressure and cardiac rate will be measured using an automated blood pressure monitor.

(iii) Dietary Intake: Two questionnaires will be administered to the students: the modified version of the Parental Authority Questionnaire (PAQ) (34, 35) and the modified version of Parent-Initiated Motivational Climate Questionnaire-2 (PIMCQ-2) (36). Another questionnaire (modified version of the Family Eating and Activity Habits Questionnaire (FEAHQ) (37-39) will be administered to parents and children to investigate the dietary intake of children. The modified version of the PAQ includes 30 items for every parent, presented in a 5-level Likert scale. The first 10 items of the data describe parental authority behaviour examples, the next 10 are of parental authoritarian behaviour, and the last 10 are of parental permissiveness. This questionnaire is based on the work of Baumrind (34) and Buri (35) in order to study the 3 types of parental authority which are (a) permissive, (b) authoritarian, and (c) authoritative. The modified version of the PIMCQ-2 is a 36-item questionnaire containing 3 subscales: Learning/Enjoyment Climate, Worry Conductive Climate, and Success Without Effort Climate. In addition, we investigated the relationship between parental style and perceived motivational climate of parents through the use of a questionnaire based on an idea presented by White and Duda in 1992.

Program Evaluation: Over a 6-month period, the followings were evaluated in each group:

1. Performance in the speed trial
2. Height, weight, BMI, waist circumference, and heart rate
3. Dietary intake using the modified version of the Family Eating and Activity Habits Questionnaire by Golan.

Ethical considerations

The Medical School of University of Thessaly and Ministry of Education and Religious Affairs, Sport and Culture has approved the protocol of this study. Written informed consent from all parents was obtained before their enrolment into the study.

Data analysis: Epi Info software will be used for data entry and SPSS software for the statistical analysis. Qualitative variables will be presented as absolute (N) and relative frequencies (%). Quantitative variables will be presented as mean with standard deviation or median with interquartile range. Kolmogorov-Smirnov test and Levene test will be used to test the normality and the equality of variances of the quantitative data, respectively. An analysis of variance (ANOVA) or the Kruskal-Wallis test will be used to analyze differences in age, body weight, and BMI among participants before the interven-

tion. A One-way ANOVA, with Bonferroni corrections for multiple comparisons, will be conducted to explore differences among groups at the baseline for parents and children's data. The effects of the intervention program will be analyzed by the repeated measures ANOVA or the Friedman test. Changes in life style behaviours from the baseline to the end of the intervention will be assessed using a chi-square test for categorical variables. A Pearson or a Spearman correlation coefficient and a linear regression analysis will be performed to explore any associations between quantitative variables. The Cronbach alpha coefficient will be used to estimate the internal consistency of the PAQ, the PIMCQ, and the FEAHQ. A p value < .05 will be considered as statistically significant.

Quality Control

Monitoring the quality of the FETA program is considered to be essential to ensure a robust study and to maintain internal validity. Several procedures are employed to ensure that the quality of the study will be optimal. These procedures are described below.

(i) Assessor Blinding: Data collection personnel will be blind to participants' group allocation. Children and their parents will be asked not to inform the data collection personnel of their group allocation status.

(ii) Training: A trained physical education teacher and accredited practising dietitians will run the project. Data collection personnel who are involved in the assessments will be trained by full-time study members before the assessments.

Ethics

The protocol of the study has been approved by the General Assembly of the Medical School of University of Thessaly and also by the Ministry of Education and Religious Affairs, Sport and Culture. Written informed consent from all parents will be obtained before their enrolment into the study.

Discussion

This paper describes the protocol of the Feeding Exercise Randomised Clinical Trial in Adolescents (FETA) in the region of Larissa in Greece, a randomized controlled trial, among overweight/obese adolescents. The FETA program consists of 2 parts. The first part of the project (descriptive study) is expected to provide useful information about the prevalence and the risk factors of obesity among adolescents in the region of Thessaly, Greece. The second part consists of a clinical trial, the first clinical trial among obese adolescents in Greece, which is expected to provide useful information on the effectiveness of clinical interventions among overweight and obese adolescents in Greece.

Acknowledgment

This study was part of a PhD dissertation at the Faculty of Medicine, School of Health Science, University of Thessaly, Greece University of Thessaly, Greece. We express our sincere gratitude to the participants in the study

and their families. In addition, we acknowledge the contribution of Dr Ioannis Karydis.

Conflict of Interests

The authors declare that they have no competing interests.

References

- Fryar CD, Carroll MD, Ogden CL. Prevalence of Overweight, Obesity, and Severe Obesity Among Children and Adolescents Aged 2–19 Years: United States, 1963–1965 Through 2015–2016. Division of Health and Nutrition Examination Surveys, NCHS Health E-Stats 2018.
- Abarca-Gómez L, Abdeen ZA, Hamid ZA, Abu-Rmeileh NM, Acosta-Cazares B, Acuin C, et al. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet*. 2017;390(10113):2627–42.
- Spinelli A, Buoncrisiano M, Kovacs VA, Yngve A, Spiroski I, Obreja G, et al. Prevalence of Severe Obesity among Primary School Children in 21 European Countries. *Obes Facts*. 2019;12 (2):244–258.
- Knightsmith P, Treasure J, Schmidt U. Spotting and supporting eating disorders in school: recommendations from school staff. *Health Educ Res*. 2013;28(6):1004–1013.
- World Health Organization. New global estimates of child and adolescent obesity released on World Obesity Day. 2017. <https://www.who.int/end-childhood-obesity/news/new-estimate-child-adolescent-obesity/en/>
- Rokholm B, Baker JL, Sorensen TI. The levelling off of the obesity epidemic since the year 1999 – a review of evidence and perspectives. *Obes Rev*. 2010;11(12):835–46.
- Fleming MT, Robinson M, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the global burden of disease study 2013. *Lancet*. 2014;384(9945):766–781.
- Withrow D, Alter D. The economic burden of obesity worldwide: a systematic review of the direct costs of obesity. *Obes Rev*. 2011;12(2):131–141.
- Bibiloni MdElMar, Pons A, Tur JA. Prevalence of Overweight and Obesity in Adolescents: A Systematic Review. *ISRN Obes*. 2013;2013:392747.
- Global database on the Implementation of Nutrition Action (GINA) 2018. Geneva: World Health Organization (<http://www.who.int/nutrition/gina/en/>, accessed 30 January 2018).
- Shrewsbury VA, O'Connor J, Steinbeck KS, Stevenson K, Lee A, Hill AJ, Kohn MR, Shah S, Torvaldsen S, Baur LA. A randomised controlled trial of a community-based healthy lifestyle program for overweight and obese adolescents: the Loozit study protocol. *BMC Public Health*. 2009;9:119.
- Biro FM, Wien M. Childhood obesity and adult morbidities. *The Am J Clin Nutr*. 2010;91(5):1499S–1505S.
- Reinert KRS, Po'e EK, Barkin SL. The relationship between executive function and obesity in children and adolescents: a systematic literature review. *J Obes*. 2013;2013:820956.
- Tremmel M, Gerdtham Ulf-G, Nilsson PM, Saha S. Economic Burden of Obesity: A Systematic Literature Review. *Int J Environ Res Public Health*. 2017;14(4): 435.
- Specchia ML, Veneziano MA, Cadeddu C, Ferriero AM, Mancuso A, Ianuale C, et al. Economic impact of adult obesity on health systems: A systematic review. *Eur J Public Health*. 2015;25(2):255–262.
- World Health Organization Obesity 2016. [(accessed on 4 October 2016)]; Available online: <http://www.who.int/topics/obesity/en/>
- Centers for Disease Control and Prevention Overweight & Obesity 2016. [(accessed on 4 October 2016)]; Available online: <https://www.cdc.gov/obesity/adult/defining.html>.
- Mayr HL, Cohen F, Isenring E, Soenen S; Project GRIT Team, Marshall S. Multidisciplinary lifestyle intervention in children and adolescents - results of the project GRIT (Growth, Resilience, Insights, Thrive) pilot study. *BMC Pediatr*. 2020;20(1):174.
- Partridge SR, Raeside R, Singleton AC, Hyun K, Latham Z, Grunseit A, Steinbeck K, Chow C, Redfern J. Text message behavioral intervention for teens on eating, physical activity and social wellbeing (TEXTBITES): Protocol for a Randomized Controlled Trial. *JMIR Res Protoc*. 2020;9(2):e16481.
- Centers for Disease Control and Prevention Healthy People 2020. (accessed in December 2010). Available online: https://www.cdc.gov/nchs/healthy_people/hp2020.htm
- López-Olmedo N, Carriquiry AL, Rodríguez-Ramírez S, Ramírez-Silva I, Espinosa-Montero J, Hernández-Barrera L, et al. Usual Intake of Added Sugars and Saturated Fats Is High while Dietary Fiber Is Low in the Mexican Population. *J Nutr*. 2016;146(9):1856S–1865S.
- Pes GM, Tolu F, Poulain M, Canelada A, Errigo A, Dore MP. The Eating Habits of School Children from the Longevity Blue Zone of Sardinia: a Positive Model. *J Nutr Food Sci*. 2015;5:5.
- Kohl HW III, Cook HD. Physical Activity and Physical Education: Relationship to Growth, Development, and Health. Educating the Student Body: Taking Physical Activity and Physical Education to School. Committee on Physical Activity and Physical Education in the School Environment; Food and Nutrition Board; Institute of Medicine; Washington (DC): National Academies Press (US); 2013.
- Centers for Disease Control and Prevention Physical activity for everyone. 2011 <http://www.cdc.gov/physicalactivity/everyone/glossary/index.html>.
- Booth FW, Roberts CK, La MJ. Lack of exercise is a major cause of chronic diseases. *Compr Physiol*. 2012;2(2):1143–1211.
- Csigé I, Ujvárosy D, Szabó Z, Lőrincz I, Paragh G, Harangi M, et al. The Impact of Obesity on the Cardiovascular System. *J Diabetes Res*. 2018.
- Barroso TA, Marins LB, Alves R, Gonçalves ACS, Barroso SG, Rocha GDS. Association of central obesity with the incidence of cardiovascular diseases and risk factors. *Int J Cardiovasc Sci*. 2017;30:5.
- Jelastopulu E, Kallianezos P, Merekioulas G, Alexopoulos EC, Sapountzi-Krepia D. Prevalence and risk factors of excess weight in school children in West Greece. *Nurs Health Sci*. 2012;14(3): 372–80.
- Kosti RI, Panagiotakos DB, Mihos CC, Alevizos A, Zampelas A, Mariolis A, Tountas Y. Dietary habits, physical activity and prevalence of overweight/obesity among adolescents in Greece: the Vyronas study. *Med Sci Monit*. 2007;13(10):437–44.
- Krassas GE, Tzotzas T, Tsametis C, Konstantinidis T. Prevalence and trends in overweight and obesity among children and adolescents in Thessaloniki, Greece. *J Pediatr Endocrinol Metab*. 2001;14(5):1319–26.
- Manios Y, Biribilis M, Moschonis G, Biribilis G, Mougios V, Lionis C, et al. “Healthy Growth Study” group. Childhood Obesity Risk Evaluation based on perinatal factors and family sociodemographic characteristics: CORE index. *Eur J Pediatr*. 2013;172(4): 551–5.
- International standards for anthropometric assessment. International Society for the Advancement of Kinanthropometry ISAK; 2001.
- Cole JT, Bellizzi CM, Flegal MC, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 2000;320(7244):1240.
- Baumrind D. Current patterns of parental authority. *Dev Psychol*. 1971;4(1):2–4.
- Buri JR. Parental Authority Questionnaire. *J Pers Assess*. 1991;57(1):110–119.
- White SA, Duda JL, Hart S. An exploratory examination of the Parent-initiated Motivational Climate Questionnaire. *Percept Mot Skills*. 1992;75(3 pt 1):875–80.
- Golan M. Fifteen years of the Family Eating and Activity Habits Questionnaire (FEAHQ): an update and review. *Pediatr Obes*. 2014;9(2):92–101.
- Golan M, Weizman A. Reliability and validity of the Family Eating and Activity Habits. *Eur J Clin Nutr*. 1998;52:771–777.
- Golan M. Parents as agents of change in childhood obesity - from research to practise. *Pediatr Obes*. 2006;1(2):66–76.