

Breast cancer risk assessment in Iranian women by Gail model

Ghazal Panahi, MD.¹, Hossein Shabahang, MD.², Helen Sahebghalam, MD.³

Department of Surgery, Islamic Azad University of Medical Sciences, Mashhad Branch, Mashhad, Iran.

Abstract

Background: Due to the high incidence of breast cancer and the effect of its early diagnosis on decreasing morbidity and mortality, we used the Gail model to study breast cancer risk in Iranian women.

Methods: This study was done in a simple randomized way. Participants were 2000 Iranian women older than 35 years old. The questionnaire consisted of demographic data such as age, race (optional) marriage status, level of education and standard questions of the Gail model. Gathered data were given in <http://bcra.nci.nih.gov/brc>. The breast cancer risk was calculated within the next 5 years and within the 90 years life span. The statistical analysis was done by SPSS software.

Results: Mean age of women in the study was 47.95 years. 50% of women had their menarche at/older than fourteen years of age. 50% of women had first childbirth in their twenties. 87% had a negative family history of breast cancer. 94% had no history of breast biopsy. Of the remaining 6%, no tissue atypia was reported. In the present study, breast cancer risk within 5 years was 0.92% and the breast cancer risk within 90 years of life was 9.14%. 7% showed risk of more than 1.67% in the Gail model. The age ranged between 55-65 years in this high risk group.

Conclusion: In our study of breast cancer risk in Iranian women, breast cancer risk was lower than the control group in the Gail model estimate for 5 years and a 90-year life span. (0.92% versus 1.02%, 9.14% versus 11.21%). The differences were statistically significant ($p < 0.001$)

Keywords: breast cancer, risk assessment, Gail model, high risk.

Introduction

Breast cancer comprises about 33% of cancers in women. It is also the most common cause of cancer-related death in women 40-44 years old [1,2]. Breast cancer occurs in 1 out of 8 women. So breast cancer risk assessment and the recognition of high risk individuals in the society, prompted researchers to use risk assessment models such as the Gail model [1-4].

Recognition of high risk individuals of breast

cancer leads to implementation for social policies for screening, prevention, early diagnosis and less radical treatment modalities that in themselves lead to increased 5-year survival rates in high risk persons [5,6]. In multiple studies, the validity and reliability of the Gail model for breast cancer risk assessment has been proven. We, therefore, used the Gail model in our study [5,8].

Methods

This study was done through a simple ran-

1. **Corresponding author**, Assistant Professor of General Surgery, Department of General Surgery, Arya Hospital, Eastern Golestan St. Mashhad, Iran. Tel.: +98511 7261823, Cell: +98915 3159558, Fax.: +98511 2232522, email: drghazalpanahi@yahoo.com.

2. Assistant Professor of General Surgery, Department of General Surgery, Mashhad University of Medical Sciences.

3. Assistant Professor of Radiology, Department of Radiology, Islamic Azad University of Medical Sciences, Mashhad Branch, Mashhad, Iran.

domized process. There were 2000 women who participated in the study who were older than 35. The questionnaire consisted of demographic data such as age, race (optional), marriage status, level of education and standard questions of the Gail model. Then the gathered data of each participant were given to <http://bcra.nci.nih.gov/brc>.

The breast cancer risk was calculated within the next 5 years and within 90 years of life span. Statistical analysis was done by SPSS software. The variables were analysed by T, χ^2 and Fischer tests. In high risk individuals with an index of equal or more than 1.67 percent of breast cancer risk assessed by the Gail model, strict recommendation for regular breast examination, mammography, and consultation for chemoprevention was given.

Results

The mean age of women participating in the study was 47.95 years. About 50% of women had their menarche at/older than fourteen years of age. About 50% of women had their first childbirth in their twenties. 87% of women had negative family history of breast cancer. 94% of participants had no history of breast biopsy. Of the remaining 6%, no tissue atypia had been reported.

In the present study, breast cancer risk within 5 years was 0.92 percent and the breast cancer risk within 90 years of life was 9.14 percent. Seven percent of participants showed a risk of more than 1.67 percent in the Gail model. The age ranged between 55-65 years in this high risk group.

Discussion

This study was performed to assess the breast cancer risk in Iranian women older than 35. Due to the proven reliability and validity of the Gail model, this model was used in the present study [5,8], which takes into account factors such as age, age at menarche, breast biopsy, age at 1st childbirth and the presence of breast cancer in

Age (yrs)	No. (%) of participants
35-45	999 (50)
46-55	640 (32)
55-65	255 (12.8)
>65	106 (5.2)
Total	2000 (100)
Education	
None	93 (4.6)
High school qualified or lower	918 (46)
University qualified	634 (31.7)
Not specified	355 (17.7)
Total	2000 (100)
Occupation	
None	1024 (51.2)
Employed	584 (29.2)
Self-employed	10 (0.5)
Retired	57 (2.9)
Not specified	325 (16.2)
Total	2000 (100)
Race (optional)	- (--)

Table 1. Demographic data of participating women.

Age (yrs)	No. (%) of participants
35-45	999 (50)
46-55	640 (32)
55-65	255 (12.8)
>65	106 (5.2)
Mean age	47 ± 9.5 years
Age at menarche	
<12	144 (8)
12-13	834 (41)
14 ≤	1022 (51)
Breast cancer in 1st degree-relatives	
No. history	1739 (87)
1	217 (10.8)
2	41 (2.1)
3	3 (0.1)
Age at 1st live birth (yrs)	
No pregnancy	41 (2.1)
<20	1032 (51.6)
20-24	555 (27.7)
25-29	282 (14.1)
30<	90 (4.5)
Breast biopsy	
No biopsy	1878 (94)
Once	94 (5)
Twice	13 (0.5)
3 times	3 (0.1)
5 times	12 (0.4)
Atypia reported	3 (2.38)

Table 2. Gail model representations in Iranian women in the study.

Breast cancer risk assessed	In the current study	Standard of Gail model	Mean difference	t	df	P value
5-year risk of breast cancer	0.92%	1.02%	0.10%	3.98	3998	P<0.001
90-year risk of breast cancer	9.14%	11.21%	2.07%	19.72	3998	P<0.001

Table 3. Breast cancer risk assessed by Gail model in this study.

first degree relatives of the participating woman. In this model the cumulative risk of breast cancer is calculated in the corresponding decade of life.

Statistics show an increasing incidence of breast cancer in the younger age in Iranian women [9,10]. Although in most cases in our study the risk of breast cancer within 5 years and within the 90 years life span was lower than the standard controls in the Gail model, those who had a Gail score of equal or more than 1.67 percent were regarded as high risk

The high risk group of breast cancer risk assessment were seriously recommended to have regular breast examinations, mammography [11,12], chemoprevention [13-15] and in some situations, prophylactic mastectomy was discussed too [16].

In spite of the high acceptance of mammography in our study, the participating women in the high risk group had less compliance towards tamoxifen as a chemopreventive drug.

We believe that similar studies can be employed to adopt relevant policies in emphasizing clinical breast examinations, screening for breast cancer and chemoprevention with tamoxifen to decrease the incidence of advanced breast cancer in developing countries.

References

1. Guinee VF. Epidemiology of breast cancer. In: Bland KI, Copeland EM, editors. *The breast: comprehensive management of benign and malignant diseases*. Philadelphia: WB Saunders; 1998. pp.339.
2. Jemal A, Murray T, Samuels A, et al. Cancer statistics. *CA Cancer J Clin* 2003. 53 (5): 157.

3. Jay RH, Lippman ME, Morrow M, Kent C. *Diseases of breast*. 3rd ed. Vol 1. 2004 : 103-105, 115-119, 363-372.

4. Ries LAG, Eisner MP, Kosary CL, et al. *Seer Cancer Statistics Review*, National Cancer Institute. Bethesda, MD. 2005. [Retrieved at: <http://seer.cancer.gov/csr>]

5. Spiegelamn D, Colditz GA, Hunter D, et al. Validations of the Gail et al model for predicting individual breast cancer risk. *J Natl Cancer Inst* 1994. 86: 600-7.

6. Bondy ML, Lustbader ED, Halabi S, et al. Validation of a breast cancer risk assessment model in women with a positive family history. *J Natl Cancer Inst* 1994. 86: 620-5

7. Rockhill B, Spiegelman D, Byrne C, Hunter DJ, Colditz GA. Validation of the Gail et al model of breast cancer risk prediction and implication for chemoprevention. *J Natl Cancer Inst* 2001. 93 (5): 334-5.

8. National Cancer Institute. Breast cancer risk assessment tool. Retrieved at: <http://www.bcr.nci.nih.gov/brc>

9. Yavari P, Mosavizadeh M, Sadrol-Heftazi B, Mehraabi Y. Reproductive characteristics and the risk of breast cancer: a case control study in Iran. *Asian Pac J Cancer Prev* 2005. 69 (3): 370-5

11. Kerlikowske k, Grady D, Rubin SM, et al. Efficacy of screening mammography. A meta-analysis. *JAMA* 1995; 273 : 149-54.

12. Kopans DB. An overview of the breast cancer screening controversy. *J Natl Cancer Inst Monogr* 1997. 22: 1-3.

13. Fisher B, Costantino JP, Wickerham DL, et al. Tamoxifen for prevention of breast cancer: report of the National Surgical Adjuvant Breast and Bowel project p-1 study. *J Natl Cancer Inst* 1998; 90: 1371-88.

14. Powles T, Eeles R, Ashley S, et al. Interim analysis of the incidence of breast cancer in the Royal Marsden Hospital tamoxifen randomised chemoprevention trial. *The lancet* 1998. 352: 98-101.

15. Gail MH, Costantino JP, Bryant J, et al. Weighing the risks and benefits of tamoxifen treatment for preventing breast cancer. *J Natl Cancer Inst* 1999; 91: 1829-46.

16. Scheuer L, Kauff N, Robson M. et al. Outcome of preventive surgery and screening for breast and ovarian cancer in BRCA mutation carriers. *J Clin Oncol* 2002; 20: 1260-8.