

DEVELOPMENT OF A METHOD FOR THE DETERMINATION OF 4-HYDROXY-3-METHOXY-MANDELIC ACID (VANILLYLMANDELIC ACID) IN URINE BY PAPER CHROMATOGRAPHY

DURDI QUJEQ, Ph.D., AND Q. USEFNIA PASHA, M.D.

From the Department of Biochemistry and Biophysics, Babol University of Medical Sciences, Babol, I.R. Iran.

ABSTRACT

We have developed a simple and precise paper chromatographic method for the determination of 4-hydroxy-3-methoxymandelic acid (VMA) in urine. Concentrations of VMA in patients with neuroblastoma were increased in comparison to controls. The linearity was excellent in the concentration range tested. The within-assay coefficient of variation for control and patient urine was less than 2.2%. The recovery was in the range of 97.9-99.4%. Results from testing urine samples of controls and patients with neuroblastoma suggest that this method is a reliable and convenient system for quantification of VMA in urine and can be used in the mass screening of neuroblastoma in infants. Sample preparation requires minimal time and the entire procedure is completed within 5 h.

MJIRI, Vol. 14, No. 3, 267-270, 2000.

Keywords: Vanillylmandelic acid, paper chromatography, neuroblastoma.

INTRODUCTION

4-Hydroxy-3-methoxymandelic acid (VMA) is one of the major catecholamine metabolites normally excreted in urine. The determination of urinary VMA is of importance for the diagnosis of neuroblastoma, pheochromocytoma and Parkinson's disease.^{1,2} Knowledge of the excretion pattern of VMA in urine is an effective aid in the diagnosis of catecholamine secreting tumors such as pheochromocytoma, neuroblastoma and ganglioneuroma.³ Neuroblastoma is one of the few malignant tumors that excrete unambiguous markers for diagnosis. Neuroblastoma, a malignant tumor of infancy and childhood, has a relatively good prognosis, even with disseminated disease.⁴ Neuroblastoma is the most common solid tumor of childhood with an incidence of 1 in 7000 children under the age of 5 years. It arises from cells of the sympathetic nervous system and has the characteristic of secreting dopamine and its major metabolite homovanillic acid, in excess.⁵ Recently, a great deal of progress has been made concerning the use of VMA in the evaluation of psychiatric patients and VMA has been used to monitor chronic lead exposure and response to medication during the treatment of Parkinson's disease.⁶ Many procedures have been

applied to the assay of VMA including thin layer and paper chromatography,^{7,8} electrophoresis,⁹ spectrophotometry,¹⁰ gas chromatography,¹¹ gas chromatography-mass spectrometry,¹² and high-performance liquid chromatography.¹³⁻¹⁶ However, these methods are difficult to perform quantitatively for use in routine clinical chemistry laboratories.

Our aim in the present work was to separate VMA from other urinary metabolites and devise a simple, low-cost, accurate, and precise method for monitoring urinary VMA concentrations. The assay presented here is a simple procedure for quantifying VMA by paper chromatography. This method is applicable for use in routine clinical chemistry laboratories.

MATERIAL AND METHODS

Pure VMA, citrate, sodium azide, sodium sulfate, Whatman paper, formic acid, P-nitroaniline, ethyl acetate, phenolic acid, ether, ethanol, acetone and formamide were obtained from Sigma (St. Louis, Mo., U.S.A.). Other reagents were purchased from Merck (Darmstadt, F.R.G.).

Standard preparation: A stock solution (500 µg/mL of VMA) was prepared by diluting 50 mg of VMA to 100 mL

Determination of VMA Levels in Urine

roblastoma patients. After testing more urine samples, particularly those from infants suffering from neuroblastoma, we believe that this paper chromatographic method can be used effectively in the mass screening of neuroblastoma in infants.

REFERENCES

1. Agbaba D, Stojanov M, Rajacic S, Zivanov-Stakie D, Majkie-Singh N: Densitometric determination of urinary 4-hydroxy-3-methoxymandelic acid (vanillylmandelic acid). *Clin Chem* 39 (12): 2500-2503, 1993.
2. Nebinger P, Koel M: Automated analysis of urinary 3-hydroxy-mandelic acid using ion-pair chromatography and fluorimetric detection. *Journal of Chromatography* 434: 43-49, 1988.
3. Ni P, Guyon F, Caude M, Rosset R: Automatic determination of urinary 4-hydroxy-3-methoxymandelic (vanillylmandelic) acid by liquid chromatography with electrochemical detection. *Journal of Chromatography* 424: 255-262, 1988.
4. Tokuda T, Tokieda T: Simultaneous determination of vanillylmandelic acid, homovanillic acid and creatinine in urine by reversed-phase ion-pair chromatography. *Journal of Chromatography* 530: 418-423, 1990.
5. Shi RZ, Ho YP, Yeung JHK, Yu OR PM, Wah TOKK, Lau MWM, Arumanayagam M: Development of an enzyme-linked immunosorbent assay with monoclonal antibody for quantification of homovanillic acid in human urine samples. *Clinical Chemistry* 44 (8): 1674-1679, 1998.
6. Binder SR, Sivorinovsky G: Measurement of urinary vanillylmandelic acid and homovanillic acid by high-performance liquid chromatography with electrochemical detection following extraction by ion-exchange and ion-moderated partition. *Journal of Chromatography* 336: 173-188, 1984.
7. Annina JS, Lipson M, Williams LA: Determination of 3-methoxy-4-hydroxymandelic acid (VMA) in urine by thin-layer chromatography. *Clin Chem* 11: 905-913, 1965.
8. Vahidi A, Roberts HR, San Filippo Jr J, Sankar DV: Paper chromatographic quantitation of 4-hydroxy-3-methoxy-mandelic acid (VMA) in urine. *Clin Chem* 17: 903-907, 1971.
9. Hermann GA: The determination of urinary 3-methoxy-4-hydroxymandelic (vanillylmandelic) acid by means of electrophoresis with cellulose acetate membrane. *Am J Clin Pathol* 41: 373-379, 1964.
10. Pasiano JJ, Crout JR, Abraham D: The determination of 3-methoxy-4-hydroxymandelic (vanillylmandelic) acid in urine. *Clin Chim Acta* 7: 285-291, 1962.
11. Brewster MA, Berry DH, Moriarty M: Urinary 3-methoxy-4-hydroxyphenylacetic-9 (homovanillic) and 3-methoxy-4-hydroxymandelic (vanillylmandelic) acids: gas-liquid chromatographic methods and experience with 13 cases of neuroblastoma. *Clin Chem* 23 (12): 2247-2249, 1977.
12. Seviour JI, McGill AC, Dale G: Method of measurement of urinary homovanillic acid and vanillylmandelic acid by gas chromatography-mass spectrometry suitable for neuroblastoma screening. *Journal of Chromatography* 432: 273-277, 1988.
13. Eriksson BM, Persson BA, Wikstrom M: Determination of urinary vanillylmandelic acid by direct and coupled-column chromatography with electrochemical detection. *Journal of Chromatography* 527: 11-19, 1990.
14. Lindgren K, Rodopoulos N: Determination of vanillylmandelic acid with ion-pair chromatography and fluorescence detection. *Clin Chem* 32 (4): 693-696, 1986.
15. Imai Y, Ito S, Maruta K, Fujita K: Simultaneous determination of catecholamines and serotonin by liquid chromatography, after treatment with boric acid gel. *Clin Chem* 34 (3): 528-530, 1988.
16. Kawaguchi S, Hirachi N, Fukamachi M: New analyzer for the determination of urinary vanillylmandelic acid, homovanillic acid and creatinine. *Journal of Chromatography* 567: 11-19, 1991.
17. Sunderman Jr FW, Cleveland PD, Law NC, Sunderman FW: A method for the determination of 3-methoxy-4-hydroxymandelic acid, vanillylmandelic acid, for the diagnosis of pheochromocytoma. *Am J Clin Pathol* 34: 293-312, 1960.
18. Oishi S: Random one-hour urine catecholamine and 4-hydroxy-3-methoxy-mandelic acid assays for diagnosis of pheochromocytoma. *Clin Chim Acta* 103: 335-342, 1980.
19. Yoshida A, Yoshika M, Yamazaki T: Urinary levels of vanillylmandelic acid and homovanillic acid determined by high-speed liquid chromatography. *Clin Chim Acta* 73: 315-320, 1976.
20. Nishi M, Miyake H, Takeda T, Hanai J, Fujita K, Ichimiya H, Tanaka T, Hiramata T, Ishikawa YK, Udoh T, Azuma H, Takase A: The relationship between homovanillic/vanillylmandelic acid ratios and prognosis in neuroblastoma. *Oncol Rep* 5 (3): 631-633, 1998.
21. Frostad B, Tani E, Kogner P, Maeda S, Bjork O, Skoog L: The clinical use of fine needle aspiration cytology for diagnosis and management of children with neuroblastic tumours. *Eur J Cancer* 34 (4): 529-536, 1998.
22. Chan SC, Chan KL, Peh WC: Clinics in diagnostic imaging, adrenal neuroblastoma. *Singapore Med J* 38 (11): 504-508, 1997.